



**HAL**  
open science

## New insights into the reduction systems of plastidial thioredoxins point out the unique properties of thioredoxin z from Arabidopsis

Anne-Sophie Bohrer, Vincent Massot, Gilles Innocenti, Jean-Philippe Reichheld, Emmanuelle Issakidis-Bourguet, H el ene Vanacker

### ► To cite this version:

Anne-Sophie Bohrer, Vincent Massot, Gilles Innocenti, Jean-Philippe Reichheld, Emmanuelle Issakidis-Bourguet, et al.. New insights into the reduction systems of plastidial thioredoxins point out the unique properties of thioredoxin z from Arabidopsis. *Journal of Experimental Botany*, 2012, 63 (18), pp.6315-6323. 10.1093/jxb/ers283 . hal-02118559

**HAL Id: hal-02118559**

**<https://univ-perp.hal.science/hal-02118559>**

Submitted on 28 Nov 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

1 **New insights into the reduction systems of plastidial thioredoxins point out the**  
2 **unique properties of thioredoxin z from Arabidopsis<sup>1</sup>**

3

4 Anne-Sophie Bohrer<sup>2</sup>, Vincent Massot<sup>2</sup>, Gilles Innocenti, Jean-Philippe Reichheld<sup>3</sup>,  
5 Emmanuelle Issakidis-Bourguet\* and H el ene Vanacker

6

7 Institut de Biologie des Plantes, UMR CNRS 8618, Saclay Plant Sciences, Univ  
8 Paris-Sud, 91405, Orsay cedex, France; <sup>3</sup>Laboratoire G enome et D eveloppement des  
9 Plantes, Univ Perpignan, UMR CNRS 5096, 66860, Perpignan cedex, France  
10

11 <sup>1</sup> This work was supported by MENRT grants to A-S. B. and V. M. and the Saclay  
12 Plant Sciences program (SPS, ANR-10-LABX-40)

13 <sup>2</sup> These two authors equally contributed to this work

14 \* Corresponding author; e-mail [emmanuelle.issakidis-bourguet@u-psud.fr](mailto:emmanuelle.issakidis-bourguet@u-psud.fr), phone +  
15 33 1 69 15 33 37, fax + 33 1 69 15 34 23

16

17 Date of submission: 12<sup>th</sup> of July 2012, revised 12<sup>th</sup> of September 2012

18 Number of tables and figures: 1 table and 6 figures

1 **ABSTRACT**

2 In plants, thioredoxins (TRX) constitute a large protein disulfide oxido-reductase  
3 family comprising ten plastidial members in *Arabidopsis thaliana* (Arabidopsis)  
4 subdivided in five types. The f and m-types regulate enzymes involved mainly in  
5 carbon metabolism whereas the x, y and z types have an antioxidant function. The  
6 reduction of TRXm and f in chloroplasts is performed in the light by  
7 ferredoxin:thioredoxin reductase (FTR) that uses photosynthetically reduced  
8 ferredoxin (Fd) as a reductant. The reduction system of Arabidopsis TRXx, y and z  
9 has never been demonstrated. Recently, a gene encoding an atypical plastidial  
10 NADPH-dependent TRX reductase (NTRC) was found. In the present study, gene  
11 expression analysis revealed that both reductases are expressed in all organs of  
12 Arabidopsis and could potentially serve as electron donors to plastidial TRX. This  
13 ability was tested *in vitro* either with purified NTRC in presence of NADPH or with  
14 a light-driven reconstituted system comprising thylakoids and purified Fd and FTR.  
15 Our results demonstrate that FTR reduces the x and y TRX isoforms but not the  
16 recently identified TRXz. Moreover, we show that NTRC cannot be an efficient  
17 alternative reducing system neither for TRXz nor for the other plastidial TRX. Our  
18 data reveal that TRXf, m, x and y, known as redox regulators in the chloroplast, have  
19 also the ability to reduce, *in vitro*, TRXz. Overall, the present study points out the  
20 unique properties of TRXz among plastidial TRX.

21 Key words: Thioredoxin, FTR, NTRC, Redox signaling, Plastid, Arabidopsis

## 1 **INTRODUCTION**

2 Thioredoxins (TRX) belong to a large multigenic family of oxidoreductases found in  
3 all free living organisms. These small soluble proteins (12-14 kDa) share a  
4 characteristic fold, a conserved active site WCG/PPC sequence, and a low redox  
5 potential that confers on them strong reductive properties. The two cysteines of the  
6 active site are involved in thiol/disulfide exchanges with their target proteins.

7 The first plant thioredoxins (m and f-types) were identified as regulators of  
8 chloroplastic enzymes related to carbon metabolism (Buchanan, 1980). Genomic  
9 revolution revealed that plant TRX isoforms are encoded by multi-gene families  
10 (Meyer et al., 2005; Nuruzzaman et al., 2008). In *Arabidopsis thaliana* (Arabidopsis),  
11 twenty TRX isoforms are found, divided into three subgroups: cytosolic,  
12 mitochondrial and plastidial. Sequence similarities and intron positions revealed that  
13 plastidial isoforms comprised nine members in Arabidopsis, subdivided into four  
14 types, the f, m, x and y-types (Mestres-Ortega and Meyer, 1999; Meyer et al., 2005),  
15 with four isoforms in the m-type, 2 in the f-type, 2 in the y-type, 1 in the x-type.  
16 Recently, two independent studies identified a tenth plastidial thioredoxin, TRXz  
17 (Arsova et al., 2010; Meng et al., 2010).

18 This multiplicity of TRX isoforms raised the question of their redundancy or  
19 possible functional specialization. Complementation studies on a TRX-deficient yeast  
20 strain hypersensitive to oxidative stress showed that Arabidopsis TRXf, m and x  
21 displayed different efficiencies (Issakidis-Bourguet et al., 2001). Afterwards,  
22 biochemical *in vitro* studies revealed specificity of chloroplastic TRX towards their  
23 target enzymes. Indeed, TRXf and m redox regulate the activity of enzymes mainly  
24 involved in the primary metabolism, whereas the more recently identified isoforms x,

1 y and z can serve as hydrogen donors for antioxidant enzymes such as peroxiredoxins  
2 (PRX), suggesting their role in oxidative stress defence (Collin et al., 2003, 2004;  
3 Lamkemeyer et al., 2006; Née et al., 2009; Chibani et al., 2011). Moreover, recent  
4 proteomic studies revealed a multiplicity (nearly 100) of putative chloroplastic targets  
5 involved in a wide range of processes (Michelet et al., 2006; Lemaire et al., 2007;  
6 Viera Dos Santos et al., 2007; Valerio et al. 2010). New putative TRX targets were  
7 also found in isolated organelles from wheat endosperm (Balmer et al., 2006) and in  
8 *Arabidopsis* root extracts (Marchand et al., 2010).

9       Once oxidized, TRX can be re-reduced by specific reductases. In the light, the  
10 photosynthetic electron transfer chain reduces ferredoxin (Fd) which can donate  
11 electrons to various enzymes including Fd:TRX reductase (FTR). This key enzyme of  
12 the Fd/TRX regulatory system has been shown to reduce efficiently TRX<sub>m</sub> and f in a  
13 reconstituted light activation system (Droux et al., 1987). In the cytosol and  
14 mitochondria of plant cells, TRX are reduced by an NADPH-dependent TRX  
15 reductase (NTR) (Jacquot et al., 1994; Laloï et al., 2001; Reichheld et al., 2005). A  
16 new gene encoding an atypical plastidial NTR (named NTRC) was identified in  
17 *Arabidopsis*, rice and *Medicago truncatula* (Serrato et al., 2004; Pérez-Ruiz et al.,  
18 2006; Alkhalifioui et al., 2007). It encodes a bipartite protein containing an NTR  
19 domain fused to a TRX domain. NTRC was shown to be an efficient redox system for  
20 chloroplast protection against oxidative damage and proposed as a possible alternative  
21 plastidial TRX reducing system (Serrato et al., 2004; Pérez-Ruiz et al., 2006).

22       The recent finding of three new plastidial TRX types (x, y and z) as well as the  
23 presence of plastidial TRX isoforms in non-photosynthetic organs (Collin et al., 2004;  
24 Balmer et al., 2006; Traverso et al., 2008) raised the question of their reducing  
25 system(s). One can wonder whether NTRC can reduce plastidial TRX in non-

1 photosynthetic tissues, or be a reducing system for the x, y and z isoforms. A direct  
2 biochemical test can assess the ability of a given reductase to reduce a given TRX.  
3 However, the interaction is possible only if the two partners share the same spatial and  
4 temporal localization. Thus, in the present work, the expression patterns of the ten  
5 genes encoding the plastidial TRX from Arabidopsis together with the four genes  
6 encoding the plastidial TRX reductase subunits were first examined using quantitative  
7 RT-PCR. Gene expression profiles revealed that both FTR and NTRC could  
8 potentially serve as reductants for all TRX in plastids. This ability was tested directly  
9 *in vitro* using NTRC in presence of NADPH, or FTR in a light-driven reconstituted  
10 Fd/TRX system. Here we demonstrate that FTR can reduce TRXx and y but not  
11 TRXz. Moreover, we show that NTRC cannot be an alternative reducing system for  
12 plastidial TRX, but that reduced plastidial thioredoxins are able to reduce TRXz *in*  
13 *vitro*.

## 1 **MATERIALS AND METHODS**

### 2 **Reagents**

3 All biochemical reagents were purchased from Sigma-Aldrich, unless otherwise  
4 mentioned.

5

### 6 **Plant Material**

7 *Arabidopsis thaliana* wild-type (Columbia ecotype) were grown on soil in the  
8 greenhouse under natural growth conditions (20°C day/18°C night temperature, 60%  
9 day/55% night hygrometry, light intensity 80-100  $\mu\text{E m}^{-2} \text{s}^{-1}$  ) and watered every 2  
10 days.

11 Peas (*Pisum sativum* var. merveille de Kelvedon) were grown in the greenhouse on  
12 vermiculite, watered daily and harvested 3 weeks after sowing.

13

### 14 **RNA Isolation and cDNA Synthesis**

15 Total RNA were isolated from 100 mg of frozen plant material (except for seeds:  
16 30 mg) using the NucleoSpin® RNA Plant kit from Macherey-Nagel, according to  
17 manufacturer's instructions. For RNA extraction from dry seeds 5% (w/v)  
18 polyvinylpyrrolidone was added to the extraction buffer. The reverse transcription  
19 step was performed with up to 1  $\mu\text{g}$  of total RNA, using the SuperScript III First-  
20 Strand Synthesis System for RT-PCR kit (Invitrogen) with oligo(dT) primer.

21

## 1      **Quantitative RT-PCR Analysis**

2      Gene expression profiling was conducted by real-time quantitative RT-PCR using  
3      gene-specific primer pairs and procedures detailed in Table S1.

4

## 5      **Expression, Production and Purification of Recombinant Proteins**

6      Proteins and related purification procedures are described in Table S2.

7

## 8      **2-Cys PRX Activity Assays**

9      PRX activity was assayed in 30 mM Tris-HCl buffer, pH 7.9, in the presence of 20  
10     μM 2-Cys PRXA from Arabidopsis, 400 μM DTT, 600 μM H<sub>2</sub>O<sub>2</sub> and 10 μM TRX.  
11     Remaining hydrogen peroxide was quantified colorimetrically at 560 nm using  
12     Peroxoquant reagent (Perbio Science). Alternatively, 2-Cys PRX activity was  
13     determined as oxidation of NADPH followed at 340 nm in a reaction mixture  
14     containing 100 mM phosphate buffer, pH 7.2, 10 mM EDTA, 1 mM H<sub>2</sub>O<sub>2</sub>, 10 μM  
15     PRX, 200 nM NTRC and 150 μM NADPH.

16

## 17     **Radiolabeling of SH groups in TRX z**

18     A sample of TRXz preparation (60μg) was alkylated with [<sup>14</sup>C]iodoacetamide  
19     (IAM) (0.5 MBeq/μmol, 2 mM), without treatment, or after an oxidizing (trans-4,5-  
20     dihydroxy-1,2-dithiane (DTTox) 10 mM, 15 min) or a reducing treatment  
21     (dithiotreitol (DTTred) 100 mM, 15 min). After trichloroacetic acid precipitation and  
22     separation by SDS-PAGE (Droux et al., 1987), proteins were Coomassie blue stained



1 and labeling was visualized by exposure of the dried gel to a low energy imaging  
2 screen, for two weeks (personal molecular Imager FX system, BioRad).

3

#### 4 **Reduction of TRX<sub>x, y</sub> and z by the Fd/FTR System**

5 The reconstituted light activation system comprised, in a total volume of 50  $\mu$ L,  
6 washed thylakoids equivalent to 10-20  $\mu$ g chlorophyll, prepared as described in Droux  
7 et al. (1987), 100 mM Tris-HCl, pH 7.9, 2-5  $\mu$ M Fd, 10  $\mu$ M FTR, 25  $\mu$ M TRX and 5  
8  $\mu$ M subunit mutant NADP-MDH. Aliquots were withdrawn as a function of time of  
9 illumination ( $120 \mu\text{E m}^{-2} \text{s}^{-2}$ ) and, for TRX<sub>x</sub> and y, tested for NADP-MDH activation  
10 level. For TRX<sub>z</sub> (NADP-MDH omitted) thiol redox status was estimated by redox  
11 western. Alternatively to the light-dependent system, 10  $\mu$ M TRX<sub>z</sub> was incubated for  
12 15 min in the presence of 20  $\mu$ M NADPH, 40 nM FNR and 1  $\mu$ M FTR and 1  $\mu$ M Fd  
13 according to Chibani et al., (2011). Free thiol groups were alkylated for 25 min, with  
14 25 mM N-ethyl maleimide (NEM) and 30 mM iodoacetic acid (IAA). Alkylated  
15 proteins were subjected to non-reducing SDS-PAGE and the reduction state of TRX<sub>z</sub>  
16 was evaluated from its monomer/dimer status by Western blot.

17

#### 18 **Chemical Activation of NADP-Malate Dehydrogenase**

19  $\Delta$ 15-NADP-MDH (shortened by 15 amino acids at the N-terminus) whose  
20 biochemical characteristics are identical to those of the full-length wild-type MDH  
21 (Johansson et al., 1999) and mutant C207A/C365A/C377A NADP-MDH were used.  
22 Enzymes (5  $\mu$ M) were incubated at room temperature, with 25  $\mu$ M TRX plus 1 mM  
23 DTT, or 5 or 100  $\mu$ M TRX<sub>z</sub> plus 100 mM DTT or DTT alone. The enzymatic activity

1 was measured in 100 mM Tris-HCl, pH 7.9, 180  $\mu$ M NADPH, 750  $\mu$ M oxaloacetate.  
2 Disappearance of NADPH was followed spectrophotometrically at 340 nm.

3

#### 4 **Immunodetection of TRX z by Western Blot**

5 Samples were separated by non-reducing 12% SDS-PAGE and transferred onto  
6 nitrocellulose membrane (iBlot Gel Transfer Device, Invitrogen). Detection was  
7 performed using either a commercial anti *Strep*-tag II monoclonal antibody  
8 conjugated to horseradish peroxidase (IBA) at a dilution of 1:10,000 or a primary  
9 specific anti-TRXz antibody at a dilution of 1:2,500 and a secondary horseradish  
10 peroxidase conjugated antibody at a dilution of 1:50,000, and revealed by enhanced  
11 chemiluminescence (ECL, GE Healthcare). The specificity of the anti-TRXz antibody  
12 was checked towards all the other plastidial TRX by Western blot (Figure S1).

13

#### 14 **Reduction of plastidial TRX by NTRC and NTRA tested** 15 **spectrophotometrically**

16 The ability of Arabidopsis cytosolic NTRA or plastidial NTRC to reduce TRX was  
17 measured by the 5,5'-dithio-bis (2-nitrobenzoic acid) (DTNB, Pierce) reduction assay.  
18 The reaction mixture (1 mL) contained 20 nM NTRC or NTRA, 150  $\mu$ M NADPH, 1  
19 mM DNTB and 10  $\mu$ M TRX in 100 mM NaPO<sub>4</sub>, pH 8.0 and 1 mM EDTA. All  
20 measurements were carried out at 25°C by monitoring the increase in absorbance at  
21 412 nm caused by the release of thionitrobenzoate (TNB<sup>-</sup>).

22

1        **Reduction of TRXz by the GSH/GRX system, NTRC or plastidial TRX**

2        TRXz was pre-oxidized by 10 mM DTT<sub>ox</sub> during 20 min. DTT<sub>ox</sub> was removed on  
3        a PD Spin Trap G-25 column (GE Healthcare). Oxidized TRXz (10 μM) was  
4        incubated for 20 min at room temperature with various reductants: either 1, 5 or 10  
5        mM reduced glutathione (GSH) or 300 μM NADPH, 6 μg/mL bakers yeast  
6        glutathione reductase (GR), 5 mM GSH and 10 μM chloroplastic glutaredoxin  
7        GRXS16. NTRC (2 μM) was tested with 500 μM NADPH. Plastidial TRX (f1, m1, x  
8        or y1; 10 μM) were also tested in presence of 1 mM DTT. Redox state of TRXz was  
9        evaluated from its monomer/dimer status by redox western.

## 1 **RESULTS**

2

### 3 **Transcript levels reveal that both NTRC and FTR are expressed along with** 4 **plastidial TRX, including in non-photosynthetic organs**

5 Using real-time quantitative PCR the presence of mRNAs encoding plastidial TRX  
6 was detected in all organs of Arabidopsis at different levels depending on the isoform  
7 (Figure 1A). Globally, genes encoding m-type TRX (except m3) and TRX-x were  
8 more abundantly expressed, especially in leaves and young seedlings. Genes coding  
9 for TRXf1 and f2 and TRXy2, though expressed at a lower level, showed similar  
10 expression profiles. The gene coding for TRXz was expressed at the lowest level and  
11 its expression was restricted to leaves, flowers and young siliques. Noteworthy,  
12 Attrxm3 and Attrxy1 expression patterns differed from those of other plastidial  
13 Arabidopsis trx genes, being predominantly expressed in non-photosynthetic tissues *e.*  
14 *g.* mature siliques, roots and dry seeds. As previously suggested (Collin et al., 2004)  
15 the gene encoding TRXy1 is the most strongly expressed in seeds among all plastidial  
16 Arabidopsis TRX. Overall, these results are consistent with transcriptomic data  
17 obtained from the “Arabidopsis eFP Browser” (Figure S2).

18 The expression levels of the genes encoding the variable subunits FTRA1 and  
19 FTRA2 of FTR (Keryer et al., 2004) were close to the one encoding NTRC (Figure  
20 1B). The gene encoding the catalytic FTRB subunit was expressed in all organs tested  
21 including non-photosynthetic organs. FTRB transcripts were quite abundant as  
22 compared to FTRA1 and FTRA2. Expression profiles of both genes encoding FTR  
23 variable subunits were more strongly expressed in leaves, young seedlings and mature

1 siliques. Similarly, NTRC showed a wide expression profile with a higher level in  
2 green tissues.

3 These findings prompted us to test both FTR and NTRC in their ability to reduce  
4 plastidial TRX. TRX<sub>x</sub> and <sub>y</sub> have been characterized in our laboratory (Collin et al.,  
5 2003, 2004) and showed specific antioxidant functions. However, their reduction  
6 system was still unknown as well as the one of Arabidopsis TRX<sub>z</sub>.

7

### 8 **TRX<sub>z</sub> is a functional oxidoreductase with target specificities**

9 Among the well-known TRX-dependent chloroplastic enzymes, NADP-MDH is  
10 probably the most studied enzyme (Ruelland and Miginiac-Maslow, 1999). It is  
11 devoid of activity in its oxidized state and gets activated in presence of reduced TRX<sub>f</sub>  
12 or <sub>m</sub>, except <sub>m3</sub> (Collin et al., 2003). We expressed TRX<sub>z</sub> in its mature form (N-  
13 terminal transit peptide removed) in *E. coli* as a *Strep-tagged* protein. It failed to  
14 activate wild-type sorghum NADP-MDH in the presence of DTT (data not shown).

15 Activation of NADP-MDH involves the reduction of two regulatory disulfides  
16 located in sequence extensions at the N- and C-terminus (Issakidis et al., 1994). We  
17 engineered a mutant form of NADP-MDH where only the N-terminal disulfide  
18 remains (C207A/C365A/C377A mutant, Ruelland et al., 1997). This mutant has a  
19 basal activity and is more easily activated, due to a less negative redox potential. It is  
20 activated by plastidial TRX isoforms that do not activate the wild type enzyme, such  
21 as TRX<sub>x</sub> and <sub>y</sub> (Collin et al, 2003; 2004). Unlike TRX<sub>m1</sub> used as a positive control,  
22 TRX<sub>z</sub> proved inefficient (even at 100 μM) towards mutant NADP-MDH, in the  
23 presence of a high (100 mM) DTT concentration necessary to reach a complete  
24 reduction of this TRX. Thus, TRX<sub>z</sub> is unable to activate NADP-MDH (Table 1).

1 We also tested the capacity of TRXz to mediate proton donation to 2-Cys PRX by  
2 following the reduction of H<sub>2</sub>O<sub>2</sub> by PRX in the presence of DTT. 2-Cys PRX is  
3 known to be reduced efficiently by TRXx, to a lesser extent by TRXy, and not by  
4 TRXm3 (Collin et al., 2003; 2004). Figure 2 shows that, indeed TRXx is the best  
5 reductant for 2-Cys PRX, Trxm3 being quite inefficient. TRXz was as efficient as  
6 TRXy1 validating that our preparation is redox active. Thus, unlike popular TRXz  
7 (Chibani et al., 2011), Arabidopsis TRXz can donate electrons to homologous 2-Cys  
8 PRX.

9

#### 10 **FTR reduces efficiently TRXx and y but cannot reduce TRXz**

11 FTR is the known reductant of plastidial TRXf and m but its capacity to reduce the  
12 more recently identified Arabidopsis TRX isoforms x, y and z was never studied. We  
13 tested the capacity of FTR to reduce plastidial Arabidopsis TRX in a reconstituted  
14 system comprising illuminated pea thylakoids, Arabidopsis ferredoxin 2 (Fd) and  
15 spinach FTR (Droux et al., 1987). In this system, TRXx, y1 and y2 were able to  
16 activate mutant NADP-MDH with efficiencies comparable to TRXf1 (Figure 3). In  
17 the dark, or when a component of the activation mixture was omitted, no NADP-  
18 MDH activation was observed (Figure 3 and data not shown). These data provide  
19 evidence that TRXx and y can be efficiently reduced by FTR. Since TRXz is not able  
20 to activate mutant NADP-MDH, its reduction was followed by redox western. Indeed,  
21 we observed that TRXz forms dimers when oxidized and gets monomerized upon  
22 reduction by DTT. We checked that monomerization was correlated with the  
23 appearance of free thiols on the protein using radiolabeling with the thiol-alkylating  
24 agent iodoacetamide (<sup>14</sup>C-IAM). A strong and specific labeling was observed for the

1 monomer in its reduced form, most probably corresponding to the reduction of the  
2 active-site disulfide (Figure 4). As there was no labeling of the dimer, it should  
3 correspond to oxidized protein. This hypothesis is strengthened by the absence of  
4 labeling of the oxidized monomer, corresponding to the initial redox state of our  
5 TRXz purified preparation and after a treatment with DTTox (Figure 4). This  
6 observation allowed us to use the monomerization/dimerization behavior of the  
7 protein as a test of reduction/oxidation. We tested TRXz reduction by FTR in the  
8 reconstituted light system described above (Figure 5). Upon illumination, no  
9 monomerization of TRXz was observed. In the same experiment, the light-driven  
10 reduction of TRXf1 (evidenced by NADP-MDH activation) proved efficient (data not  
11 shown). Inability of FTR to reduce TRXz was confirmed using an *in vitro*  
12 reconstituted “dark” system consisting of NADPH, FNR, Fd and FTR (Figure S5B,  
13 lane 9). These results provide evidence that FTR can be the physiological reductase of  
14 plastidial TRXx and y, with comparable efficiencies to TRXf. Noteworthy; TRXz is  
15 the first plastidial TRX which is not reduced by their classical reductase.

#### 16 **NTRC is not an efficient alternative reducing system for plastidial TRX**

17 Most plastidial TRX genes are expressed in green tissues but, as mentioned above,  
18 some isoforms, such as TRXy1, are predominantly present in non-photosynthetic  
19 tissues where the mRNA corresponding to NTRC could be detected. Moreover, as  
20 FTR cannot reduce TRXz, NTRC could possibly be its physiological substitute. We  
21 produced and purified the Arabidopsis recombinant NTRC. Its functionality was  
22 confirmed by its capacity to mediate an efficient degradation of H<sub>2</sub>O<sub>2</sub> by 2-Cys PRXA  
23 (Figure S3). We then assayed the reduction of the five types of plastidial TRX by  
24 NTRC using DTNB as a final electron acceptor, as this is a useful tool for measuring  
25 TRX oxido-reductase capacity. At 20 nM, NTRC alone, probably due to its own TRX

1 domain, displays a TRX reductase activity with a  $k_{\text{cat}}$  value ( $\text{s}^{-1}$ ) of  $4.47 \pm 0.59$ . This  
2 activity increases 2-3 fold in presence of plastidial TRX (Figure S4A). However, this  
3 reduction rate corresponds to the basal rate observed with the cytosolic NTRA for  
4 TRXf1 that is five-fold lower than the reduction rate of the cytosolic TRXh3 (Figure  
5 S4B). Moreover, when the ability of NTRC to reduce TRXz was tested by following  
6 its monomerization by redox western, no increase of the monomeric form at the  
7 expense of the dimers was observed (Figure S5A). The faint intermediate molecular  
8 weight bands, more abundant in aged preparations (data not shown), most probably  
9 correspond to the dimer's conformational variants as they are recognized by the anti-  
10 TRXz antibody and disappear upon reduction. We conclude that TRXz cannot be  
11 efficiently reduced by NTR.

12

### 13 **TRXz is not reduced by the GSH/GRX system**

14 Previous data have demonstrated that a subgroup of cytosolic TRX, which are not  
15 reduced by NTR, can be reduced by glutaredoxin (GRX) which is itself reduced by  
16 glutathione (GSH) (Gelhaye et al., 2003). In this system, known as the GSH/GRX  
17 system, the transmission of the redox signal is mediated from NADPH to GSH *via*  
18 glutathione reductase (GR). None of TRXf, m, x and y are reduced by GSH  
19 (unpublished results). However, Chibani et al. (2011) have recently found close values  
20 for the redox midpoint potentials ( $E_m$ ) of TRXz and GSH (Noctor, 2006). The ability  
21 of TRXz to be reduced by GR, GSH and/or a plastidial GRX (GRXS16;  
22 Bandyopadhyay et al., 2008) was tested by following its reduction through its  
23 monomerization. No monomerization of TRXz occurred with any of these



1 components (Fig. S5B). Thus we conclude that none of the components of the  
2 GSH/GRX system can reduce TRXz *in vitro*.

3

#### 4 **Plastidial TRX can reduce TRXz**

5 The  $E_m$  value for the catalytic disulfide of TRXz was found to be -311 mV at pH  
6 7.9 (Chibani et al., 2011), which is less negative than the  $E_m$  values of the other  
7 plastidial TRX ranging from -365 to -335mV (Collin et al., 2003; 2004). Therefore,  
8 we investigated the ability of other plastidial TRX to reduce TRXz (Figure 6). While 1  
9 mM DTT could not directly reduce TRXz, in presence of TRXf or m, the amount of  
10 monomeric TRXz was largely increased at the expense of the dimeric form (Figure  
11 6A). In comparison, TRXx and y exhibited a significant, though lower, efficiency  
12 (Figure 6B). These data demonstrate that TRXz can be reduced by other plastidial  
13 TRX *in vitro*.

## 1        **DISCUSSION**

2        The main goal of the present work was to test *in vitro* the ability and specificity of  
3        plastidial reductases to reduce the various plastidial TRX. As a preliminary, we show,  
4        using quantitative RT-PCR, that the ten genes encoding plastidial TRX together with  
5        the three genes encoding FTR subunits and the gene encoding NTRC were expressed  
6        in the same organs, at the same time (Figure 1) and thus could be partners. These data  
7        will need to be further completed with a study at the protein level and gene expression  
8        analyses in different cell types. The detection of a messenger RNA suggesting the  
9        presence of the translated corresponding protein, these findings prompted us to test  
10       both FTR and NTRC in their capacities to reduce plastidial TRX and particularly the  
11       more recently identified TRX<sub>x</sub>, y and z isoforms from Arabidopsis.

12       FTR is the well-known reductase for TRX<sub>f</sub> and m isoforms, TRX<sub>x</sub>, y and z were  
13       found as efficient proton donors to peroxiredoxins (Collin et al., 2003, 2004; present  
14       work: Figure 2) but the upstream reduction of TRX by following PRX activity in a  
15       reconstituted Fd/FTR/TRX light activation system cannot be easily tested. Indeed,  
16       reactive oxygen species (ROS) production by illuminated thylakoids interferes with  
17       the activity test. However, the ability of FTR in reducing TRX<sub>x</sub> and y could be tested  
18       by following the activation of an engineered mutant NADP-MDH enzyme which can  
19       be reduced by these TRX isoforms (Figure 3). Our results demonstrate that TRX<sub>x</sub> and  
20       y are reduced by FTR with efficiencies comparable to those of the m and f-type  
21       isoforms (Droux et al., 1987). Thus, in the light, FTR is clearly the reducing system  
22       for plastidial TRX<sub>f</sub>, m, x and y in Arabidopsis. However, the recent finding of f and  
23       m-TRX in non-photosynthetic organs raised the question of their reducing system in  
24       heterotrophic plastids (Traverso et al., 2008). Moreover, the presence of a complete  
25       FTR system in amyloplasts was demonstrated (Balmer et al., 2006). In roots, FNR

1 reduces type III Fd which is more easily reduced than leaf isoforms, as it displays a  
2 less negative  $E_m$  (Hanke et al., 2004). This property favors the electron transfer from  
3 NADPH to FTR via FNR and Fd. The functionality of this system was already tested  
4 by König et al. (2002) for TRXf and m but still needs to be tested for other plastidial  
5 TRX isoforms, especially the ones expressed in non-photosynthetic tissue such as  
6 TRXm3 and y1.

7 The ability of FTR in reducing TRXz could be assessed in the reconstituted light  
8 system by following the reduction state of TRXz through its monomerization. Indeed,  
9 TRXz could not activate NADP-MDH, either wild-type or mutant, but the labeling  
10 experiment (Figure 4) showing that only the reduced monomer can be derivatized  
11 offered us a simple way to observe the reduction of this TRX. Our data show that FTR  
12 cannot reduce TRXz in the light (Figure 5) in contrast with all the other plastidial  
13 TRX. This finding is not consistent with the work of Chibani et al. (2011) where a  
14 heterologous NADPH-driven FTR reduction system was used. These authors assessed  
15 the reduction of poplar TRXz by a shift of migration during non-reducing SDS-PAGE  
16 following the alkylation with methoxyl-PEG maleimide. A shift of *ca.* 15 kDa, instead  
17 of 4 kDa as expected for the alkylation of two thiol groups, was obtained. This form  
18 could correspond to the dimer we observed in our experiments. A  
19 dimerization/monomerization upon oxidation/reduction was previously observed for  
20 TRXf which contains a third Cys residue responsible for this behavior (Michelet et al.,  
21 2005). In the mature Arabidopsis TRXz, no additional Cys residue is found, indicating  
22 that dimerization is linked to the specific properties of this isoform.

23 Noteworthy, in our light activation experiment, when TRXz was mixed with  
24 thylakoid membranes, no monomeric form was observed at all (Figure 5), even in the  
25 dark, while our TRXz preparations always appeared as a mixture of monomers and

1 dimers, even in the NADPH/FNR/FTR/Fd reconstituted system (Figure S4).  
2 Moreover, the amount of TRXz seemed to be lower in the dark (time 0) and to  
3 increase upon illumination (Figure 5). Therefore, TRXz might interact (directly or  
4 indirectly) with thylakoid membranes in the dark and could be released in the light.  
5 Such a phenomenon was recently reported for FNR (Benz et al., 2010) and would  
6 deserve a more detailed study.

7 The NTR domain of NTRC being active in the reduction of the TRX domain, it  
8 has been proposed as a possible alternative plastidial TRX reducing system (Serrato et  
9 al., 2004). In the present work, we tested Arabidopsis NTRC with homologous TRX  
10 isoforms belonging to the five plastidial types. We found that, *in vitro*, NTRC cannot  
11 reduce efficiently any plastidial TRX. This confirms the results obtained by others  
12 (Serrato et al., 2004; Pérez-Ruiz et al., 2006; Chibani et al., 2011) testing heterologous  
13 proteins.

14 Our data show that FTR is the only reductase capable to reduce TRXf, m, x  
15 and y in plastids and reveal that even though TRXz is expressed mainly in green  
16 tissues, it is not reduced by any known plastidial TRX reduction system. Interestingly,  
17 we found that other plastidial TRX can reduce TRXz. This is thermodynamically  
18 feasible as these TRX have a redox midpoint potential ( $E_m$ ) less negative than the  $E_m$   
19 of TRXz (Collin et al., 2003; 2004; Chibani et al., 2011). This is the first example, to  
20 our knowledge, of a TRX reducing another TRX. Overall, these biochemical data  
21 underscoring that TRXz stands out from the other plastidial isoforms are consistent  
22 with the severe albino phenotype of the Arabidopsis *trxz* knock-out mutant (Arsova et  
23 al., 2010; Meng et al., 2010) contrasting with the mutants of the other plastidial TRX  
24 that do not exhibit a visible phenotype under standard growth conditions (Laugier et  
25 al., 2012).

1 TRXz was previously identified in a proteomic study as a component of  
2 transcriptionally active chromosome in plastids (pTAC; Pfalz et al., 2006) and more  
3 recently as a subunit of a complex formed with the plastid-encoded RNA polymerase  
4 (PEP; Schröter et al., 2010). *trxz* mutant or silenced plants are affected in the  
5 expression of the chloroplast genome, particularly the genes transcribed by the PEP  
6 (Arsova et al., 2010; Schröter et al., 2010). This strongly suggests the implication of  
7 TRXz in the transcription of the plastome. TRXz interacts with several targets,  
8 including 2 Cys-PRX, FNR and two proteins with similarities to plastid fructokinase-  
9 like (FLN) proteins (Schröter et al., 2010) previously identified as a component of  
10 pTAC (Pfalz et al., 2006) but with no fructokinase activity (Arsova et al., 2010). It has  
11 been shown that TRXz acts as an adaptor protein between the receptor-like protein Cf-  
12 9 and the Ser/Thr kinase ACIK1 in the cytosol of tomato (Nekrasov et al., 2006).  
13 Recently, Schröter et al. (2010) proposed that TRXz could play a similar adaptor role  
14 as a subunit of the PEP complex for regulation of plastid gene expression. TRXz  
15 would be an additional key player of photosynthetic gene expression (Dietz and  
16 Pfannschmidt, 2011).

17 Overall, this points to a unique role of TRXz, which indeed possesses TRX  
18 activity. TRXz appears as an important sensor of the redox status of the chloroplast  
19 being potentially oxidized by 2-Cys PRX (sensing reactive oxygen species) and  
20 reduced by other TRX (sensing light indirectly) revealing its potential important role  
21 in the redox homeostasis of the chloroplast.

## **Supplementary material**

Supplementary data are available at *JXB* online.

Table S1: List of specific primers used for RT-qPCR and experimental procedure.

Table S2: Summary of the different proteins used and purification procedures.

Figure S1: Anti-TRXz antibody specificity.

Figure S2: Expression patterns of plastidial thioredoxins and reductases from the « Arabidopsis eFP browser » website.

Figure S3: Reduction of H<sub>2</sub>O<sub>2</sub> by 2-Cys PRX using NTRC as an electron donor.

Figure S4: Reduction of plastidial TRX by NTRs.

Figure S5: Reduction of TRXz by various reductants.

## **Acknowledgments**

The authors are grateful to Dr N. Rouhier for providing GRXS16 protein, to Dr M. Zaffagnini for technical advices and scientific discussions and to Dr M. Miginiac-Maslow for support, helpful suggestions and critical reading of the manuscript.

## REFERENCES

- Alkhalfioui F, Renard M, Montrichard F.** 2007. Unique properties of NADP-thioredoxin reductase C in legumes. *Journal of Experimental Botany* **58**: 969-978
- Arsova B, Hoja U, Wimmelbacher M, Greiner E, Ustun S, Melzer M, Petersen K, Lein W, Bornke F.** 2010. Plastidial thioredoxin z interacts with two fructokinase-like proteins in a thiol-dependent manner: evidence for an essential role in chloroplast development in *Arabidopsis* and *Nicotiana benthamiana*. *Plant Cell* **22**: 1498-1515
- Balmer Y, Vensel WH, Cai N, Manieri W, Schürmann P, Hurkman WJ, Buchanan BB.** 2006. A complete ferredoxin/thioredoxin system regulates fundamental processes in amyloplasts. *The Proceedings of the National Academy in Sciences USA* **103**: 2988-2993
- Bandyopadhyay S, Gama F, Molina-Navarro MM, Gualberto JM, Claxton R, Naik SG, Huynh BH, Herrero E, Jacquot J-P, Johnson M K, Rouhier N.** 2008. Chloroplast monothiol glutaredoxins as scaffold proteins for the assembly and delivery of [2Fe-2S] clusters. *The EMBO Journal* **27**: 1122-1133
- Benz JP, Lintala M, Soll J, Mulo P, Bolter B.** 2010. A new concept for ferredoxin NADP(H) oxidoreductase binding to plant thylakoids. *Trends in Plant Science* **15**: 608-613
- Boyer ME, Wang CW, Swartz JR.** 2006. Simultaneous expression and maturation of the iron-sulfur protein ferredoxin in a cell-free system. *Biotechnology and Bioengineering* **94**: 128-138
- Broin M, Cuine S, Eymery F, Rey P.** 2002. The plastidic 2-cysteine peroxiredoxin is a target for a thioredoxin involved in the protection of the photosynthetic apparatus against oxidative damage. *Plant Cell* **14**: 1417-1432
- Buchanan BB.** 1980. Role of light in the regulation of chloroplast enzymes. *Annual Review of Plant Physiology and Plant Molecular Biology* **31**: 341-374
- Chibani K, Tarrago L, Schürmann P, Jacquot J-P, Rouhier N.** 2011. Biochemical properties of poplar thioredoxin z. *FEBS Letters* **585**: 1077-1081
- Collin V, Issakidis-Bourguet E, Marchand C, Hirasawa M, Lancelin JM, Knaff DB, Miginiac-Maslow M.** 2003. The *Arabidopsis* plastidial thioredoxins - New functions and new insights into specificity. *The Journal of Biological Chemistry* **278**: 23747-23752
- Collin V, Lamkemeyer P, Miginiac-Maslow M, Hirasawa M, Knaff DB, Dietz K-J, Issakidis-Bourguet E.** 2004. Characterization of plastidial thioredoxins from *Arabidopsis* belonging to the new y-type. *Plant Physiology* **136**: 4088-4095



**Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible W-R.** 2005. Genome-wide identification of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiology* **139**: 5-17

**Dietz K-J and Pfannschmidt T.** 2011. Novel regulators in photosynthetic redox control of plant metabolism and gene expression. *Plant Physiology* **155**: 1477-1485

**Droux M, Miginiac-Maslow M, Jacquot JP, Gadal P, Crawford NA, Kosower NS, Buchanan BB.** 1987. Ferredoxin-thioredoxin reductase - a catalytically active dithiol group links photoreduced ferredoxin to thioredoxin functional in photosynthetic enzyme regulation. *Archives of Biochemistry and Biophysics* **256**: 372-380

**Gelhaye E, Rouhier N, Jacquot JP.** 2003. Evidence for a subgroup of thioredoxin h that requires GSH/Grx for its reduction. *FEBS Letters* **555**: 443-448

**Hanke GT, Kimata-Arigo Y, Taniguchi I, Hase T.** 2004. A post genomic characterization of arabidopsis ferredoxins. *Plant Physiology* **134**: 255-264

**Issakidis E, Saarinen M, Decottignies P, Jacquot JP, Crépin C, Gadal P, Miginiac-Maslow M.** 1994. Identification and characterization of the 2nd regulatory disulfide bridge of recombinant sorghum leaf NADP-malate dehydrogenase. *The Journal of Biological Chemistry* **269**: 3511-3517

**Issakidis-Bourguet E, Mouaheb N, Meyer Y, Miginiac-Maslow M.** 2001. Heterologous complementation of yeast reveals a new putative function for chloroplast m-type thioredoxin. *The Plant Journal* **25**: 127-135

**Jacquot J-P, Rivera-Madrid R, Marinho P, Kollarova M, Le Maréchal P, Miginiac-Maslow M, Meyer Y.** 1994. *Arabidopsis thaliana* NADPH thioredoxin reductase cDNA characterization and expression of the recombinant protein in *Escherichia-coli*. *Journal of Molecular Biology* **235**: 1357-1363

**Johansson K, Ramaswamy S, Saarinen M, Lemaire-Chamley M, Issakidis-Bourguet E, Miginiac-Maslow M, Eklund H.** 1999. Structural basis for light activation of a chloroplast enzyme: The structure of sorghum NADP-malate dehydrogenase in its oxidized form. *Biochemistry* **38**: 4319-4326

**Keryer E, Collin V, Lavergne D, Lemaire SD, Issakidis-Bourguet E.** 2004. Characterization of Arabidopsis T-DNA mutants for the variable subunit of ferredoxin:thioredoxin reductase. *Photosynthesis Research* **79**: 265-274

**König J, Baier M, Horling F, Kahmann U, Harris G, Schürmann P, Dietz K-J.** 2002. The plant-specific function of 2-Cys peroxiredoxin-mediated detoxification of peroxides in the redox-hierarchy of photosynthetic electron flux. *The Proceedings of the National Academy in Sciences USA* **99**: 5738-5743

**Laloi C, Rayapuram N, Chartier Y, Grienenberger JM, Bonnard G, Meyer Y.** 2001. Identification and characterization of a mitochondrial thioredoxin system in plants. *The Proceedings of the National Academy in Sciences USA* **98**: 14144-14149

**Lamkemeyer P, Laxa M, Collin V, Li W, Finkemeier I, Schottler MA, Holtkamp V, Tognetti VB, Issakidis-Bourguet E, Kandlbinder A, Weis E, Miginiac-Maslow M, Dietz K-J.** 2006. Peroxiredoxin Q of *Arabidopsis thaliana* is attached to the thylakoids and functions in context of photosynthesis. *Plant Journal* **45**: 968-981

**Laugier E, Tarrago L, Courteille A, Innocenti G, Eymery F, Rumeau D, Issakidis-Bourguet E, Rey P.** 2012. Involvement of thioredoxin  $\gamma 2$  in the preservation of leaf methionine sulfoxide reductase capacity and growth under high light. *Plant Cell and Environment*. doi: 10.1111/PCE.12005

**Lemaire SD, Michelet L, Zaffagnini M, Massot V, Issakidis-Bourguet E.** 2007. Thioredoxins in chloroplasts. *Current Genetics* **51**: 343-365

**Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* **25**: 402-408

**Marchand C, Vanacker H, Collin V, Issakidis-Bourguet E, Le Maréchal P, Decottignies P.** 2010. Thioredoxins targets in *Arabidopsis* roots. *Proteomics* **10**: 2418-2428

**Meng L, Wong JH, Feldman LJ, Lemaux PG, Buchanan BB.** 2010. A membrane associated thioredoxin required for plant growth moves from cell to cell, suggestive of a role in intercellular communication. *The Proceedings of the National Academy in Sciences USA* **107**: 3900-3905

**Mestres-Ortega D, Meyer Y.** 1999. The *Arabidopsis thaliana* genome encodes at least four thioredoxins m and a new prokaryotic-like thioredoxin. *Gene* **240**: 307-316

**Meyer Y, Reichheld JP, Vignols F.** 2005. Thioredoxins in *Arabidopsis* and other plants. *Photosynthesis Research* **86**: 419-433

**Michelet L, Zaffagnini M, Massot V, Keryer E, Vanacker H, Miginiac-Maslow M, Issakidis-Bourguet E, Lemaire SD.** 2006. Thioredoxins, glutaredoxins, and glutathionylation: new crosstalks to explore. *Photosynthesis Research* **89**: 225-245

**Michelet L, Zaffagnini M, Marchand C, Collin V, Decottignies P, Tsan P, Lancelin JM, Trost P, Miginiac-Maslow M, Noctor G, Lemaire SD.** 2005. Glutathionylation of chloroplast thioredoxin f is a redox signaling mechanism in plants. *The Proceedings of the National Academy in Sciences USA* **102**: 16478-16483

**Née G, Zaffagnini M, Trost P, Issakidis-Bourguet E.** 2009. Redox regulation of chloroplastic glucose-6-phosphate dehydrogenase: A new role for f-type thioredoxin. *FEBS Letters* **583**: 2827-2832

**Nekrasov V, Ludwig AA, Jones JDG.** 2006. CITRX thioredoxin is a putative adaptor protein connecting Cf-9 and the ACIK1 protein kinase during the Cf-9/Avr9-induced defence response. *FEBS Letters* **580**: 4236-4241

**Noctor G.** 2006. Metabolic signalling in defence and stress: the central roles of soluble redox couples. *Plant, Cell & Environment* **29**: 409-425

**Nuruzzaman M, Gupta M, Zhang C, Wang L, Xie W, Xiong L, Zhang Q, Lian X.** 2008. Sequence and analysis of the thioredoxin gene family in rice. *Molecular Genetics & Genomics* **280**: 139-151

**Pfalz J, Liere K, Kandlbinder A, Dietz K-J, Oelmüller R.** 2006. pTAC2, -6 and -12 are components of the transcriptionally active plastid chromosome that are required for plastid gene expression. *Plant Cell* **18**: 176-197

**Pérez-Ruiz JM, Spinola MC, Kirchsteiger K, Moreno J, Sahrawy M, Cejudo FJ.** 2006. Rice NTRC is a high-efficiency redox system for chloroplast protection against oxidative damage. *Plant Cell* **18**: 2356-2368

**Reichheld J-P, Meyer E, Khafif M, Bonnard G, Meyer Y.** 2005. AtNTRB is the major mitochondrial thioredoxin reductase in *Arabidopsis thaliana*. *FEBS Letters* **579**: 337-342

**Rozen S, Skaletsky HJ.** 2000. Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds) *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press, Totowa, NJ, pp 365-386

**Ruelland E, Lemaire Chamley M, Le Marechal P, Issakidis-Bourguet E, Djukic N, Miginiac-Maslow M.** 1997. An internal cysteine is involved in the thioredoxin-dependent activation of sorghum leaf NADP-malate dehydrogenase. *The Journal of Biological Chemistry* **272**: 19851-19857

**Ruelland E, Miginiac-Maslow M.** 1999. Regulation of chloroplast enzyme activities by thioredoxins: activation or relief from inhibition? *Trends in Plant Science* **4**: 136-141

**Schröter Y, Steiner S, Matthai K, Pfannschmidt T.** 2010. Analysis of oligomeric protein complexes in the chloroplast sub-proteome of nucleic acid-binding proteins from mustard reveals potential redox regulators of plastid gene expression. *Proteomics* **10**: 2191-2204

**Serrato AJ, Pérez-Ruiz JM, Spinola MC, Cejudo FJ.** 2004. A novel NADPH thioredoxin reductase, localized in the chloroplast, which deficiency causes hypersensitivity to abiotic stress in *Arabidopsis thaliana*. *The Journal of Biological Chemistry* **279**: 43821-43827

**Traverso JA, Vignols F, Cazalis R, Serrato AJ, Pulido P, Sahrawy M, Meyer Y, Cejudo FJ, Chueca A.** 2008. Immunocytochemical localization of *Pisum sativum* TRXs f and m in non-photosynthetic tissues. *Journal of Experimental Botany* **59**: 1267-1277

**Valerio C, Costa A, Marri L, Issakidis-Bourguet E, Pupillo P, Trost P, Sparla F.** 2010. Thioredoxin-regulated  $\beta$ -amylase (BAM1) triggers diurnal starch

degradation in guard cells, and in mesophyll cells under osmotic stress. *Journal of Experimental Botany* **62**: 545-555

**Vieira Dos Santos C, Laugier E, Tarrago L, Massot V, Issakidis-Bourguet E, Rouhier N, Rey P.** 2007. Specificity of thioredoxins and glutaredoxins as electron donors to two distinct classes of Arabidopsis plastidial methionine sulfoxide reductases B. *FEBS Letters* **581**: 4371-4376

## Table 1

**Table 1.** Activation test of mutant NADP-MDH by TRXz.

Mutant NADP-MDH (0.9  $\mu\text{g}$ ) was incubated with 100 mM DTT and various concentrations of TRXz. Activation by TRXm1 was used as positive control. Activity was measured spectrophotometrically at 340 nm through the oxaloacetate-dependent disappearance of NADPH. Activation treatments were made at room temperature for 20 min in presence of DTT (100 mM) alone or with TRX (5  $\mu\text{M}$  unless otherwise mentioned).

	<b>Initial rates of reduction</b> ( $\mu\text{moles oxidized NADPH}$ $\cdot\text{mg}^{-1}\text{ NADP-MDH}\cdot\text{sec}^{-1}$ )
DTT alone	$0.71 \pm 0.06$
TRX m1	$1.36 \pm 0.09$
TRX z	$0.73 \pm 0.07$
100 $\mu\text{M}$ TRX z	$0.79 \pm 0.10$

## Legends for figures

**Figure 1.** Expression of plastidial TRX and TRX reductases in Arabidopsis.

RT-qPCR analyses were performed on total RNA isolated from young seedlings, leaves, flowers, young and mature siliques, seeds and roots. Transcript levels are the result of three independent experiments and were normalized by the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen, 2001) using the PP2A gene as standard (Czechowski et al., 2005). A, expression patterns of plastidial TRX genes, *trxf1* transcripts level in leaves used as calibrator. B, expression patterns of *ntrc* and *ftc* subunits genes, *ntrc* transcripts level in leaves used as calibrator.

**Figure 2.** Efficiency of TRXz in the reduction of 2-Cys PRX compared to other plastidial TRX.

The enzyme 2-Cys PRXA (20  $\mu$ M) was incubated in the presence of 400  $\mu$ M DTT, 600  $\mu$ M H<sub>2</sub>O<sub>2</sub> and 10  $\mu$ M TRX. As a control, 2-Cys PRX was incubated without TRX (DTT). Data are the mean of 3 independent experiments. Error bars correspond to standard deviation.

**Figure 3.** Activation of NADP-MDH by plastidial TRXf, x and y in the light, in a Fd/TRX reconstituted system.

Mutant NADP-MDH was illuminated at 120  $\mu$ E m<sup>-2</sup> s<sup>-1</sup> in the presence of freshly prepared thylakoids (equivalent to 20  $\mu$ g chlorophyll), 5  $\mu$ M ferredoxin, 2  $\mu$ M FTR, and 25  $\mu$ M TRXf1 (f1, black circles), TRXy1 (y1, black diamonds), TRXy2 (y2,

black triangles) or TRXx (x, black squares). TRXy2 was also tested in the dark (open triangles). Experiments were reproduced 3 times, a representative set of data is shown.

**Figure 4.** Redox and oligomerization state of TRXz.

TRXz was alkylated with  $^{14}\text{C}$ -IAM after an oxidizing (DTTox 10 mM) or a reducing treatment (DTTred 100 mM). The samples were separated by non-reducing SDS-PAGE. Proteins and radiolabeling were respectively visualized by Coomassie blue staining and exposure of the dried gel to a low energy imaging screen.

**Figure 5.** Reduction test of TRXz by FTR in a reconstituted light-driven Fd/TRX system.


Experimental conditions were as described in legend of Fig. 3. As controls, oxidized TRXz (10  $\mu\text{M}$ ) was incubated with (lanes 2 and 3) or without (lane 1) 100 mM DTT. For the reduction test of TRXz by FTR, TRXz (25  $\mu\text{M}$ ) was incubated in the presence of thylakoids, 5  $\mu\text{M}$  Fd and 10  $\mu\text{M}$  FTR in the light. Aliquots were withdrawn periodically, the proteins were alkylated, subjected to non-reducing SDS-PAGE, electrotransferred onto nitrocellulose membrane and probed with anti *Strep*-tag antibody. Experiment was reproduced 4 times, a representative experiment is shown.

**Figure 6.** Reduction of TRXz by other plastidial TRX.

TRXz (10  $\mu$ M) was incubated without (lane 1) or with DTT (1, 10 or 100 mM, lane 2, 7, 8 respectively) or in presence of 10  $\mu$ M of another plastidial TRX (f1, m1: panel A; x, y1: panel B) without (lane 3, 5) or with DTT (1 mM, lane 4, 6), for 15 min at room temperature. Proteins were then alkylated and subjected to non-reducing SDS-PAGE, electrotransferred onto nitrocellulose membrane and probed with an anti-TRXz antibody. Experiments were reproduced four times, a representative experiment is shown.



**Figure 1**



**Figure 2**







Figure 3



**Figure 4**




**Figure 5**




**Figure 6**


**A**



**B**



## Figure S1



**Figure S1.** Anti-TRX z antibody specificity. Anti TRX z antibody was tested against all the plastidial (f1, 2; m1, 2, 3, 4; x; y1, 2; z), a mitochondrial (o3) and a cytosolic (h3) TRX. After separation by SDS-PAGE, proteins were electrotransferred onto nitrocellulose membrane and probed with the anti-TRXz antibody.


Figure S2



**Figure S2.** Expression pattern of the *trx1*, *f2*, *m1*, *m2*, *m3*, *m4*, *x*, *y1*, *y2* and *z* and the *ntrc*, *ftra1* and *ftrb* genes from *Arabidopsis thaliana*. Data were obtained from the « Arabidopsis eFP browser » (<http://bbc.botany.utoronto.ca/efp/cgi-bin/efpWeb.cgi>). *ftra2* could not be analyzed due to the absence of the corresponding specific probe on Affymetrix Arabidopsis microarray.




**Figure S3**



**Figure S3.** Reduction of hydrogen peroxide by 2-Cys PRX using NTRC as an electron donor.

Arabidopsis NTRC (200 nM) was incubated in presence of 150 μM NADPH and its capacity as an electron donor to 2-Cys PRX A (20 μM) was tested through its capacity to reduce H<sub>2</sub>O<sub>2</sub> (1 mM). Reaction was spectrophotometrically monitored by the disappearance of the reduced cofactor at 340 nm. Data were normalized to the initial OD. Open squares: NTRC alone, open circles: NTRC plus 10 μM 2-Cys PRX, black triangles: 2-Cys PRX without NTRC. Experiments were reproduced 3 times, a representative set of data is shown.


**Figure S4**



**Figure S4.** Reduction test of plastidial TRX by NADPH-dependent TRX reductases.

NTRC (20 nM) or NTRA (20 nM) was incubated in the presence of 150  $\mu M$  NADPH and 1 mM DTNB with 10  $\mu M$  TRX. The reaction of DTNB with free thiols was followed spectrophotometrically at 412 nm through the appearance of  $TNB^-$  upon time. As controls, we tested TRXf1 in the absence of NADPH and NTRC or NTRA; the basal activity of NTRC or NTRA was measured in presence of NADPH without TRX. Data are the mean of 3 independent experiments. Error bars correspond to standard deviation.

## Figure S5



**Figure S5.** Reduction of TRXz by NTRC, GSH, GRX, GR or FTR studied by redox western. **A-** Reduction by NTRC: As a control for reduction, oxidized TRXz (10  $\mu$ M; lane 1) was fully reduced with 100 mM DTT (lane 2). Reduction of TRXz by NTRC was tested in the absence (lane 4) or the presence (lane 6) of NADPH.

**B-** Reduction by GSH, GRX, GR or FTR: As a control for reduction, oxidized TRXz (10  $\mu$ M; lane 1) was incubated with 10 or 100 mM DTT (lanes 2 and 3). Reduction of TRXz was tested either with reduced glutathione (GSH) (1, 5 or 10 mM; lanes 4 to 6) or 300  $\mu$ M NADPH, 6  $\mu$ g/mL baker's yeast glutathione reductase (GR), 5 mM GSH and 10  $\mu$ M of chloroplastic glutaredoxin GRXS16 (lane 7) or 300  $\mu$ M NADPH and 6  $\mu$ g/mL GR (lane 8) or 20  $\mu$ M NADPH, 40 nM FNR, 1  $\mu$ M Fd and 1  $\mu$ M FTR (lane 9).

After a 15 min treatment at room temperature, proteins were alkylated and subjected to non-reducing SDS-PAGE, electrotransferred onto nitrocellulose membrane and probed either with the anti-TRXz specific antibody (panel A) or with the anti-Strep-tag antibody (panel B). The reduction of TRXz was followed through its monomerization. Experiments were reproduced 3 times, a representative experiment is shown.

1 **New insights into the reduction systems of plastidial thioredoxins point out the**  
2 **unique properties of thioredoxin z from Arabidopsis<sup>1</sup>**

3

4 Anne-Sophie Bohrer<sup>2</sup>, Vincent Massot<sup>2</sup>, Gilles Innocenti, Jean-Philippe Reichheld<sup>3</sup>,  
5 Emmanuelle Issakidis-Bourguet\* and H el ene Vanacker

6

7 Institut de Biologie des Plantes, UMR CNRS 8618, Saclay Plant Sciences, Univ  
8 Paris-Sud, 91405, Orsay cedex, France; <sup>3</sup>Laboratoire G enome et D eveloppement des  
9 Plantes, Univ Perpignan, UMR CNRS 5096, 66860, Perpignan cedex, France  
10

11 <sup>1</sup> This work was supported by MENRT grants to A-S. B. and V. M. and the Saclay  
12 Plant Sciences program (SPS, ANR-10-LABX-40)

13 <sup>2</sup> These two authors equally contributed to this work

14 \* Corresponding author; e-mail [emmanuelle.issakidis-bourguet@u-psud.fr](mailto:emmanuelle.issakidis-bourguet@u-psud.fr), phone +  
15 33 1 69 15 33 37, fax + 33 1 69 15 34 23

16

17 Date of submission: 12<sup>th</sup> of July 2012, revised 12<sup>th</sup> of September 2012

18 Number of tables and figures: 1 table and 6 figures

1 **ABSTRACT**

2 In plants, thioredoxins (TRX) constitute a large protein disulfide oxido-reductase  
3 family comprising ten plastidial members in *Arabidopsis thaliana* (Arabidopsis)  
4 subdivided in five types. The f and m-types regulate enzymes involved mainly in  
5 carbon metabolism whereas the x, y and z types have an antioxidant function. The  
6 reduction of TRXm and f in chloroplasts is performed in the light by  
7 ferredoxin:thioredoxin reductase (FTR) that uses photosynthetically reduced  
8 ferredoxin (Fd) as a reductant. The reduction system of Arabidopsis TRXx, y and z  
9 has never been demonstrated. Recently, a gene encoding an atypical plastidial  
10 NADPH-dependent TRX reductase (NTRC) was found. In the present study, gene  
11 expression analysis revealed that both reductases are expressed in all organs of  
12 Arabidopsis and could potentially serve as electron donors to plastidial TRX. This  
13 ability was tested *in vitro* either with purified NTRC in presence of NADPH or with  
14 a light-driven reconstituted system comprising thylakoids and purified Fd and FTR.  
15 Our results demonstrate that FTR reduces the x and y TRX isoforms but not the  
16 recently identified TRXz. Moreover, we show that NTRC cannot be an efficient  
17 alternative reducing system neither for TRXz nor for the other plastidial TRX. Our  
18 data reveal that TRXf, m, x and y, known as redox regulators in the chloroplast, have  
19 also the ability to reduce, *in vitro*, TRXz. Overall, the present study points out the  
20 unique properties of TRXz among plastidial TRX.

21 Key words: Thioredoxin, FTR, NTRC, Redox signaling, Plastid, Arabidopsis

## 1 **INTRODUCTION**

2 Thioredoxins (TRX) belong to a large multigenic family of oxidoreductases found in  
3 all free living organisms. These small soluble proteins (12-14 kDa) share a  
4 characteristic fold, a conserved active site WCG/PPC sequence, and a low redox  
5 potential that confers on them strong reductive properties. The two cysteines of the  
6 active site are involved in thiol/disulfide exchanges with their target proteins.

7 The first plant thioredoxins (m and f-types) were identified as regulators of  
8 chloroplastic enzymes related to carbon metabolism (Buchanan, 1980). Genomic  
9 revolution revealed that plant TRX isoforms are encoded by multi-gene families  
10 (Meyer et al., 2005; Nuruzzaman et al., 2008). In *Arabidopsis thaliana* (Arabidopsis),  
11 twenty TRX isoforms are found, divided into three subgroups: cytosolic,  
12 mitochondrial and plastidial. Sequence similarities and intron positions revealed that  
13 plastidial isoforms comprised nine members in Arabidopsis, subdivided into four  
14 types, the f, m, x and y-types (Mestres-Ortega and Meyer, 1999; Meyer et al., 2005),  
15 with four isoforms in the m-type, 2 in the f-type, 2 in the y-type, 1 in the x-type.  
16 Recently, two independent studies identified a tenth plastidial thioredoxin, TRXz  
17 (Arsova et al., 2010; Meng et al., 2010).

18 This multiplicity of TRX isoforms raised the question of their redundancy or  
19 possible functional specialization. Complementation studies on a TRX-deficient yeast  
20 strain hypersensitive to oxidative stress showed that Arabidopsis TRXf, m and x  
21 displayed different efficiencies (Issakidis-Bourguet et al., 2001). Afterwards,  
22 biochemical *in vitro* studies revealed specificity of chloroplastic TRX towards their  
23 target enzymes. Indeed, TRXf and m redox regulate the activity of enzymes mainly  
24 involved in the primary metabolism, whereas the more recently identified isoforms x,

1 y and z can serve as hydrogen donors for antioxidant enzymes such as peroxiredoxins  
2 (PRX), suggesting their role in oxidative stress defence (Collin et al., 2003, 2004;  
3 Lamkemeyer et al., 2006; Née et al., 2009; Chibani et al., 2011). Moreover, recent  
4 proteomic studies revealed a multiplicity (nearly 100) of putative chloroplastic targets  
5 involved in a wide range of processes (Michelet et al., 2006; Lemaire et al., 2007;  
6 Viera Dos Santos et al., 2007; Valerio et al. 2010). New putative TRX targets were  
7 also found in isolated organelles from wheat endosperm (Balmer et al., 2006) and in  
8 *Arabidopsis* root extracts (Marchand et al., 2010).

9       Once oxidized, TRX can be re-reduced by specific reductases. In the light, the  
10 photosynthetic electron transfer chain reduces ferredoxin (Fd) which can donate  
11 electrons to various enzymes including Fd:TRX reductase (FTR). This key enzyme of  
12 the Fd/TRX regulatory system has been shown to reduce efficiently TRX<sub>m</sub> and f in a  
13 reconstituted light activation system (Droux et al., 1987). In the cytosol and  
14 mitochondria of plant cells, TRX are reduced by an NADPH-dependent TRX  
15 reductase (NTR) (Jacquot et al., 1994; Laloï et al., 2001; Reichheld et al., 2005). A  
16 new gene encoding an atypical plastidial NTR (named NTRC) was identified in  
17 *Arabidopsis*, rice and *Medicago truncatula* (Serrato et al., 2004; Pérez-Ruiz et al.,  
18 2006; Alkhalifioui et al., 2007). It encodes a bipartite protein containing an NTR  
19 domain fused to a TRX domain. NTRC was shown to be an efficient redox system for  
20 chloroplast protection against oxidative damage and proposed as a possible alternative  
21 plastidial TRX reducing system (Serrato et al., 2004; Pérez-Ruiz et al., 2006).

22       The recent finding of three new plastidial TRX types (x, y and z) as well as the  
23 presence of plastidial TRX isoforms in non-photosynthetic organs (Collin et al., 2004;  
24 Balmer et al., 2006; Traverso et al., 2008) raised the question of their reducing  
25 system(s). One can wonder whether NTRC can reduce plastidial TRX in non-

1 photosynthetic tissues, or be a reducing system for the x, y and z isoforms. A direct  
2 biochemical test can assess the ability of a given reductase to reduce a given TRX.  
3 However, the interaction is possible only if the two partners share the same spatial and  
4 temporal localization. Thus, in the present work, the expression patterns of the ten  
5 genes encoding the plastidial TRX from Arabidopsis together with the four genes  
6 encoding the plastidial TRX reductase subunits were first examined using quantitative  
7 RT-PCR. Gene expression profiles revealed that both FTR and NTRC could  
8 potentially serve as reductants for all TRX in plastids. This ability was tested directly  
9 *in vitro* using NTRC in presence of NADPH, or FTR in a light-driven reconstituted  
10 Fd/TRX system. Here we demonstrate that FTR can reduce TRXx and y but not  
11 TRXz. Moreover, we show that NTRC cannot be an alternative reducing system for  
12 plastidial TRX, but that reduced plastidial thioredoxins are able to reduce TRXz *in*  
13 *vitro*.



## 1 **MATERIALS AND METHODS**

### 2 **Reagents**

3 All biochemical reagents were purchased from Sigma-Aldrich, unless otherwise  
4 mentioned.

5

### 6 **Plant Material**

7 *Arabidopsis thaliana* wild-type (Columbia ecotype) were grown on soil in the  
8 greenhouse under natural growth conditions (20°C day/18°C night temperature, 60%  
9 day/55% night hygrometry, light intensity 80-100  $\mu\text{E m}^{-2} \text{s}^{-1}$  ) and watered every 2  
10 days.

11 Peas (*Pisum sativum* var. merveille de Kelvedon) were grown in the greenhouse on  
12 vermiculite, watered daily and harvested 3 weeks after sowing.

13

### 14 **RNA Isolation and cDNA Synthesis**

15 Total RNA were isolated from 100 mg of frozen plant material (except for seeds:  
16 30 mg) using the NucleoSpin® RNA Plant kit from Macherey-Nagel, according to  
17 manufacturer's instructions. For RNA extraction from dry seeds 5% (w/v)  
18 polyvinylpyrrolidone was added to the extraction buffer. The reverse transcription  
19 step was performed with up to 1  $\mu\text{g}$  of total RNA, using the SuperScript III First-  
20 Strand Synthesis System for RT-PCR kit (Invitrogen) with oligo(dT) primer.

21

## 1      **Quantitative RT-PCR Analysis**

2      Gene expression profiling was conducted by real-time quantitative RT-PCR using  
3      gene-specific primer pairs and procedures detailed in Table S1.

4

## 5      **Expression, Production and Purification of Recombinant Proteins**

6      Proteins and related purification procedures are described in Table S2.

7

## 8      **2-Cys PRX Activity Assays**

9      PRX activity was assayed in 30 mM Tris-HCl buffer, pH 7.9, in the presence of 20  
10      $\mu\text{M}$  2-Cys PRXA from Arabidopsis, 400  $\mu\text{M}$  DTT, 600  $\mu\text{M}$   $\text{H}_2\text{O}_2$  and 10  $\mu\text{M}$  TRX.  
11     Remaining hydrogen peroxide was quantified colorimetrically at 560 nm using  
12     Peroxoquant reagent (Perbio Science). Alternatively, 2-Cys PRX activity was  
13     determined as oxidation of NADPH followed at 340 nm in a reaction mixture  
14     containing 100 mM phosphate buffer, pH 7.2, 10 mM EDTA, 1 mM  $\text{H}_2\text{O}_2$ , 10  $\mu\text{M}$   
15     PRX, 200 nM NTRC and 150  $\mu\text{M}$  NADPH.

16

## 17     **Radiolabeling of SH groups in TRX z**

18     A sample of TRXz preparation (60 $\mu\text{g}$ ) was alkylated with [ $^{14}\text{C}$ ]iodoacetamide  
19     (IAM) (0.5 MBeq/ $\mu\text{mol}$ , 2 mM), without treatment, or after an oxidizing (trans-4,5-  
20     dihydroxy-1,2-dithiane (DTTox) 10 mM, 15 min) or a reducing treatment  
21     (dithiotreitol (DTTred) 100 mM, 15 min). After trichloroacetic acid precipitation and  
22     separation by SDS-PAGE (Droux et al., 1987), proteins were Coomassie blue stained

1 and labeling was visualized by exposure of the dried gel to a low energy imaging  
2 screen, for two weeks (personal molecular Imager FX system, BioRad).

3

#### 4 **Reduction of TRX<sub>x</sub>, y and z by the Fd/FTR System**

5 The reconstituted light activation system comprised, in a total volume of 50  $\mu$ L,  
6 washed thylakoids equivalent to 10-20  $\mu$ g chlorophyll, prepared as described in Droux  
7 et al. (1987), 100 mM Tris-HCl, pH 7.9, 2-5  $\mu$ M Fd, 10  $\mu$ M FTR, 25  $\mu$ M TRX and 5  
8  $\mu$ M subunit mutant NADP-MDH. Aliquots were withdrawn as a function of time of  
9 illumination ( $120 \mu\text{E m}^{-2} \text{s}^{-2}$ ) and, for TRX<sub>x</sub> and y, tested for NADP-MDH activation  
10 level. For TRX<sub>z</sub> (NADP-MDH omitted) thiol redox status was estimated by redox  
11 western. Alternatively to the light-dependent system, 10  $\mu$ M TRX<sub>z</sub> was incubated for  
12 15 min in the presence of 20  $\mu$ M NADPH, 40 nM FNR and 1  $\mu$ M FTR and 1  $\mu$ M Fd  
13 according to Chibani et al., (2011). Free thiol groups were alkylated for 25 min, with  
14 25 mM N-ethyl maleimide (NEM) and 30 mM iodoacetic acid (IAA). Alkylated  
15 proteins were subjected to non-reducing SDS-PAGE and the reduction state of TRX<sub>z</sub>  
16 was evaluated from its monomer/dimer status by Western blot.

17

#### 18 **Chemical Activation of NADP-Malate Dehydrogenase**

19  $\Delta$ 15-NADP-MDH (shortened by 15 amino acids at the N-terminus) whose  
20 biochemical characteristics are identical to those of the full-length wild-type MDH  
21 (Johansson et al., 1999) and mutant C207A/C365A/C377A NADP-MDH were used.  
22 Enzymes (5  $\mu$ M) were incubated at room temperature, with 25  $\mu$ M TRX plus 1 mM  
23 DTT, or 5 or 100  $\mu$ M TRX<sub>z</sub> plus 100 mM DTT or DTT alone. The enzymatic activity

1 was measured in 100 mM Tris-HCl, pH 7.9, 180  $\mu$ M NADPH, 750  $\mu$ M oxaloacetate.  
2 Disappearance of NADPH was followed spectrophotometrically at 340 nm.

3

#### 4 **Immunodetection of TRX z by Western Blot**

5 Samples were separated by non-reducing 12% SDS-PAGE and transferred onto  
6 nitrocellulose membrane (iBlot Gel Transfer Device, Invitrogen). Detection was  
7 performed using either a commercial anti *Strep*-tag II monoclonal antibody  
8 conjugated to horseradish peroxidase (IBA) at a dilution of 1:10,000 or a primary  
9 specific anti-TRXz antibody at a dilution of 1:2,500 and a secondary horseradish  
10 peroxidase conjugated antibody at a dilution of 1:50,000, and revealed by enhanced  
11 chemiluminescence (ECL, GE Healthcare). The specificity of the anti-TRXz antibody  
12 was checked towards all the other plastidial TRX by Western blot (Figure S1).

13

#### 14 **Reduction of plastidial TRX by NTRC and NTRA tested** 15 **spectrophotometrically**

16 The ability of Arabidopsis cytosolic NTRA or plastidial NTRC to reduce TRX was  
17 measured by the 5,5'-dithio-bis (2-nitrobenzoic acid) (DTNB, Pierce) reduction assay.  
18 The reaction mixture (1 mL) contained 20 nM NTRC or NTRA, 150  $\mu$ M NADPH, 1  
19 mM DNTB and 10  $\mu$ M TRX in 100 mM NaPO<sub>4</sub>, pH 8.0 and 1 mM EDTA. All  
20 measurements were carried out at 25°C by monitoring the increase in absorbance at  
21 412 nm caused by the release of thionitrobenzoate (TNB<sup>-</sup>).

22

1        **Reduction of TRXz by the GSH/GRX system, NTRC or plastidial TRX**

2        TRXz was pre-oxidized by 10 mM DTTox during 20 min. DTTox was removed on  
3        a PD Spin Trap G-25 column (GE Healthcare). Oxidized TRXz (10  $\mu$ M) was  
4        incubated for 20 min at room temperature with various reductants: either 1, 5 or 10  
5        mM reduced glutathione (GSH) or 300  $\mu$ M NADPH, 6  $\mu$ g/mL bakers yeast  
6        glutathione reductase (GR), 5 mM GSH and 10  $\mu$ M chloroplastic glutaredoxin  
7        GRXS16. NTRC (2  $\mu$ M) was tested with 500  $\mu$ M NADPH. Plastidial TRX (f1, m1, x  
8        or y1; 10  $\mu$ M) were also tested in presence of 1 mM DTT. Redox state of TRXz was  
9        evaluated from its monomer/dimer status by redox western.

## 1 **RESULTS**

2

### 3 **Transcript levels reveal that both NTRC and FTR are expressed along with** 4 **plastidial TRX, including in non-photosynthetic organs**

5 Using real-time quantitative PCR the presence of mRNAs encoding plastidial TRX  
6 was detected in all organs of Arabidopsis at different levels depending on the isoform  
7 (Figure 1A). Globally, genes encoding m-type TRX (except m3) and TRX-x were  
8 more abundantly expressed, especially in leaves and young seedlings. Genes coding  
9 for TRXf1 and f2 and TRXy2, though expressed at a lower level, showed similar  
10 expression profiles. The gene coding for TRXz was expressed at the lowest level and  
11 its expression was restricted to leaves, flowers and young siliques. Noteworthy,  
12 Attrxm3 and Attrxy1 expression patterns differed from those of other plastidial  
13 Arabidopsis trx genes, being predominantly expressed in non-photosynthetic tissues *e.*  
14 *g.* mature siliques, roots and dry seeds. As previously suggested (Collin et al., 2004)  
15 the gene encoding TRXy1 is the most strongly expressed in seeds among all plastidial  
16 Arabidopsis TRX. Overall, these results are consistent with transcriptomic data  
17 obtained from the “Arabidopsis eFP Browser” (Figure S2).

18 The expression levels of the genes encoding the variable subunits FTRA1 and  
19 FTRA2 of FTR (Keryer et al., 2004) were close to the one encoding NTRC (Figure  
20 1B). The gene encoding the catalytic FTRB subunit was expressed in all organs tested  
21 including non-photosynthetic organs. FTRB transcripts were quite abundant as  
22 compared to FTRA1 and FTRA2. Expression profiles of both genes encoding FTR  
23 variable subunits were more strongly expressed in leaves, young seedlings and mature

1 siliques. Similarly, NTRC showed a wide expression profile with a higher level in  
2 green tissues.

3 These findings prompted us to test both FTR and NTRC in their ability to reduce  
4 plastidial TRX. TRX<sub>x</sub> and <sub>y</sub> have been characterized in our laboratory (Collin et al.,  
5 2003, 2004) and showed specific antioxidant functions. However, their reduction  
6 system was still unknown as well as the one of Arabidopsis TRX<sub>z</sub>.

7

### 8 **TRX<sub>z</sub> is a functional oxidoreductase with target specificities**

9 Among the well-known TRX-dependent chloroplastic enzymes, NADP-MDH is  
10 probably the most studied enzyme (Ruelland and Miginiac-Maslow, 1999). It is  
11 devoid of activity in its oxidized state and gets activated in presence of reduced TRX<sub>f</sub>  
12 or <sub>m</sub>, except <sub>m3</sub> (Collin et al., 2003). We expressed TRX<sub>z</sub> in its mature form (N-  
13 terminal transit peptide removed) in *E. coli* as a *Strep-tagged* protein. It failed to  
14 activate wild-type sorghum NADP-MDH in the presence of DTT (data not shown).

15 Activation of NADP-MDH involves the reduction of two regulatory disulfides  
16 located in sequence extensions at the N- and C-terminus (Issakidis et al., 1994). We  
17 engineered a mutant form of NADP-MDH where only the N-terminal disulfide  
18 remains (C207A/C365A/C377A mutant, Ruelland et al., 1997). This mutant has a  
19 basal activity and is more easily activated, due to a less negative redox potential. It is  
20 activated by plastidial TRX isoforms that do not activate the wild type enzyme, such  
21 as TRX<sub>x</sub> and <sub>y</sub> (Collin et al, 2003; 2004). Unlike TRX<sub>m1</sub> used as a positive control,  
22 TRX<sub>z</sub> proved inefficient (even at 100 μM) towards mutant NADP-MDH, in the  
23 presence of a high (100 mM) DTT concentration necessary to reach a complete  
24 reduction of this TRX. Thus, TRX<sub>z</sub> is unable to activate NADP-MDH (Table 1).

1 We also tested the capacity of TRXz to mediate proton donation to 2-Cys PRX by  
2 following the reduction of H<sub>2</sub>O<sub>2</sub> by PRX in the presence of DTT. 2-Cys PRX is  
3 known to be reduced efficiently by TRXx, to a lesser extent by TRXy, and not by  
4 TRXm3 (Collin et al., 2003; 2004). Figure 2 shows that, indeed TRXx is the best  
5 reductant for 2-Cys PRX, Trxm3 being quite inefficient. TRXz was as efficient as  
6 TRXy1 validating that our preparation is redox active. Thus, unlike popular TRXz  
7 (Chibani et al., 2011), Arabidopsis TRXz can donate electrons to homologous 2-Cys  
8 PRX.

9

#### 10 **FTR reduces efficiently TRXx and y but cannot reduce TRXz**

11 FTR is the known reductant of plastidial TRXf and m but its capacity to reduce the  
12 more recently identified Arabidopsis TRX isoforms x, y and z was never studied. We  
13 tested the capacity of FTR to reduce plastidial Arabidopsis TRX in a reconstituted  
14 system comprising illuminated pea thylakoids, Arabidopsis ferredoxin 2 (Fd) and  
15 spinach FTR (Droux et al., 1987). In this system, TRXx, y1 and y2 were able to  
16 activate mutant NADP-MDH with efficiencies comparable to TRXf1 (Figure 3). In  
17 the dark, or when a component of the activation mixture was omitted, no NADP-  
18 MDH activation was observed (Figure 3 and data not shown). These data provide  
19 evidence that TRXx and y can be efficiently reduced by FTR. Since TRXz is not able  
20 to activate mutant NADP-MDH, its reduction was followed by redox western. Indeed,  
21 we observed that TRXz forms dimers when oxidized and gets monomerized upon  
22 reduction by DTT. We checked that monomerization was correlated with the  
23 appearance of free thiols on the protein using radiolabeling with the thiol-alkylating  
24 agent iodoacetamide (<sup>14</sup>C-IAM). A strong and specific labeling was observed for the



1 monomer in its reduced form, most probably corresponding to the reduction of the  
2 active-site disulfide (Figure 4). As there was no labeling of the dimer, it should  
3 correspond to oxidized protein. This hypothesis is strengthened by the absence of  
4 labeling of the oxidized monomer, corresponding to the initial redox state of our  
5 TRXz purified preparation and after a treatment with DTTox (Figure 4). This  
6 observation allowed us to use the monomerization/dimerization behavior of the  
7 protein as a test of reduction/oxidation. We tested TRXz reduction by FTR in the  
8 reconstituted light system described above (Figure 5). Upon illumination, no  
9 monomerization of TRXz was observed. In the same experiment, the light-driven  
10 reduction of TRXf1 (evidenced by NADP-MDH activation) proved efficient (data not  
11 shown). Inability of FTR to reduce TRXz was confirmed using an *in vitro*  
12 reconstituted “dark” system consisting of NADPH, FNR, Fd and FTR (Figure S5B,  
13 lane 9). These results provide evidence that FTR can be the physiological reductase of  
14 plastidial TRXx and y, with comparable efficiencies to TRXf. Noteworthy; TRXz is  
15 the first plastidial TRX which is not reduced by their classical reductase.

#### 16 **NTRC is not an efficient alternative reducing system for plastidial TRX**

17 Most plastidial TRX genes are expressed in green tissues but, as mentioned above,  
18 some isoforms, such as TRXy1, are predominantly present in non-photosynthetic  
19 tissues where the mRNA corresponding to NTRC could be detected. Moreover, as  
20 FTR cannot reduce TRXz, NTRC could possibly be its physiological substitute. We  
21 produced and purified the Arabidopsis recombinant NTRC. Its functionality was  
22 confirmed by its capacity to mediate an efficient degradation of H<sub>2</sub>O<sub>2</sub> by 2-Cys PRXA  
23 (Figure S3). We then assayed the reduction of the five types of plastidial TRX by  
24 NTRC using DTNB as a final electron acceptor, as this is a useful tool for measuring  
25 TRX oxido-reductase capacity. At 20 nM, NTRC alone, probably due to its own TRX

1 domain, displays a TRX reductase activity with a  $k_{\text{cat}}$  value ( $\text{s}^{-1}$ ) of  $4.47 \pm 0.59$ . This  
2 activity increases 2-3 fold in presence of plastidial TRX (Figure S4A). However, this  
3 reduction rate corresponds to the basal rate observed with the cytosolic NTRA for  
4 TRXf1 that is five-fold lower than the reduction rate of the cytosolic TRXh3 (Figure  
5 S4B). Moreover, when the ability of NTRC to reduce TRXz was tested by following  
6 its monomerization by redox western, no increase of the monomeric form at the  
7 expense of the dimers was observed (Figure S5A). The faint intermediate molecular  
8 weight bands, more abundant in aged preparations (data not shown), most probably  
9 correspond to the dimer's conformational variants as they are recognized by the anti-  
10 TRXz antibody and disappear upon reduction. We conclude that TRXz cannot be  
11 efficiently reduced by NTR.

12

### 13 **TRXz is not reduced by the GSH/GRX system**

14 Previous data have demonstrated that a subgroup of cytosolic TRX, which are not  
15 reduced by NTR, can be reduced by glutaredoxin (GRX) which is itself reduced by  
16 glutathione (GSH) (Gelhaye et al., 2003). In this system, known as the GSH/GRX  
17 system, the transmission of the redox signal is mediated from NADPH to GSH *via*  
18 glutathione reductase (GR). None of TRXf, m, x and y are reduced by GSH  
19 (unpublished results). However, Chibani et al. (2011) have recently found close values  
20 for the redox midpoint potentials ( $E_m$ ) of TRXz and GSH (Noctor, 2006). The ability  
21 of TRXz to be reduced by GR, GSH and/or a plastidial GRX (GRXS16;  
22 Bandyopadhyay et al., 2008) was tested by following its reduction through its  
23 monomerization. No monomerization of TRXz occurred with any of these

1 components (Fig. S5B). Thus we conclude that none of the components of the  
2 GSH/GRX system can reduce TRXz *in vitro*.

3

#### 4 **Plastidial TRX can reduce TRXz**

5 The  $E_m$  value for the catalytic disulfide of TRXz was found to be -311 mV at pH  
6 7.9 (Chibani et al., 2011), which is less negative than the  $E_m$  values of the other  
7 plastidial TRX ranging from -365 to -335mV (Collin et al., 2003; 2004). Therefore,  
8 we investigated the ability of other plastidial TRX to reduce TRXz (Figure 6). While 1  
9 mM DTT could not directly reduce TRXz, in presence of TRXf or m, the amount of  
10 monomeric TRXz was largely increased at the expense of the dimeric form (Figure  
11 6A). In comparison, TRXx and y exhibited a significant, though lower, efficiency  
12 (Figure 6B). These data demonstrate that TRXz can be reduced by other plastidial  
13 TRX *in vitro*.

## 1        **DISCUSSION**

2        The main goal of the present work was to test *in vitro* the ability and specificity of  
3        plastidial reductases to reduce the various plastidial TRX. As a preliminary, we show,  
4        using quantitative RT-PCR, that the ten genes encoding plastidial TRX together with  
5        the three genes encoding FTR subunits and the gene encoding NTRC were expressed  
6        in the same organs, at the same time (Figure 1) and thus could be partners. These data  
7        will need to be further completed with a study at the protein level and gene expression  
8        analyses in different cell types. The detection of a messenger RNA suggesting the  
9        presence of the translated corresponding protein, these findings prompted us to test  
10       both FTR and NTRC in their capacities to reduce plastidial TRX and particularly the  
11       more recently identified TRX<sub>x</sub>, y and z isoforms from Arabidopsis.

12       FTR is the well-known reductase for TRX<sub>f</sub> and m isoforms, TRX<sub>x</sub>, y and z were  
13       found as efficient proton donors to peroxiredoxins (Collin et al., 2003, 2004; present  
14       work: Figure 2) but the upstream reduction of TRX by following PRX activity in a  
15       reconstituted Fd/FTR/TRX light activation system cannot be easily tested. Indeed,  
16       reactive oxygen species (ROS) production by illuminated thylakoids interferes with  
17       the activity test. However, the ability of FTR in reducing TRX<sub>x</sub> and y could be tested  
18       by following the activation of an engineered mutant NADP-MDH enzyme which can  
19       be reduced by these TRX isoforms (Figure 3). Our results demonstrate that TRX<sub>x</sub> and  
20       y are reduced by FTR with efficiencies comparable to those of the m and f-type  
21       isoforms (Droux et al., 1987). Thus, in the light, FTR is clearly the reducing system  
22       for plastidial TRX<sub>f</sub>, m, x and y in Arabidopsis. However, the recent finding of f and  
23       m-TRX in non-photosynthetic organs raised the question of their reducing system in  
24       heterotrophic plastids (Traverso et al., 2008). Moreover, the presence of a complete  
25       FTR system in amyloplasts was demonstrated (Balmer et al., 2006). In roots, FNR

1 reduces type III Fd which is more easily reduced than leaf isoforms, as it displays a  
2 less negative  $E_m$  (Hanke et al., 2004). This property favors the electron transfer from  
3 NADPH to FTR via FNR and Fd. The functionality of this system was already tested  
4 by König et al. (2002) for TRXf and m but still needs to be tested for other plastidial  
5 TRX isoforms, especially the ones expressed in non-photosynthetic tissue such as  
6 TRXm3 and y1.

7 The ability of FTR in reducing TRXz could be assessed in the reconstituted light  
8 system by following the reduction state of TRXz through its monomerization. Indeed,  
9 TRXz could not activate NADP-MDH, either wild-type or mutant, but the labeling  
10 experiment (Figure 4) showing that only the reduced monomer can be derivatized  
11 offered us a simple way to observe the reduction of this TRX. Our data show that FTR  
12 cannot reduce TRXz in the light (Figure 5) in contrast with all the other plastidial  
13 TRX. This finding is not consistent with the work of Chibani et al. (2011) where a  
14 heterologous NADPH-driven FTR reduction system was used. These authors assessed  
15 the reduction of poplar TRXz by a shift of migration during non-reducing SDS-PAGE  
16 following the alkylation with methoxyl-PEG maleimide. A shift of *ca.* 15 kDa, instead  
17 of 4 kDa as expected for the alkylation of two thiol groups, was obtained. This form  
18 could correspond to the dimer we observed in our experiments. A  
19 dimerization/monomerization upon oxidation/reduction was previously observed for  
20 TRXf which contains a third Cys residue responsible for this behavior (Michelet et al.,  
21 2005). In the mature Arabidopsis TRXz, no additional Cys residue is found, indicating  
22 that dimerization is linked to the specific properties of this isoform.

23 Noteworthy, in our light activation experiment, when TRXz was mixed with  
24 thylakoid membranes, no monomeric form was observed at all (Figure 5), even in the  
25 dark, while our TRXz preparations always appeared as a mixture of monomers and

1 dimers, even in the NADPH/FNR/FTR/Fd reconstituted system (Figure S4).  
2 Moreover, the amount of TRXz seemed to be lower in the dark (time 0) and to  
3 increase upon illumination (Figure 5). Therefore, TRXz might interact (directly or  
4 indirectly) with thylakoid membranes in the dark and could be released in the light.  
5 Such a phenomenon was recently reported for FNR (Benz et al., 2010) and would  
6 deserve a more detailed study.

7       The NTR domain of NTRC being active in the reduction of the TRX domain, it  
8 has been proposed as a possible alternative plastidial TRX reducing system (Serrato et  
9 al., 2004). In the present work, we tested Arabidopsis NTRC with homologous TRX  
10 isoforms belonging to the five plastidial types. We found that, *in vitro*, NTRC cannot  
11 reduce efficiently any plastidial TRX. This confirms the results obtained by others  
12 (Serrato et al., 2004; Pérez-Ruiz et al., 2006; Chibani et al., 2011) testing heterologous  
13 proteins.

14       Our data show that FTR is the only reductase capable to reduce TRXf, m, x  
15 and y in plastids and reveal that even though TRXz is expressed mainly in green  
16 tissues, it is not reduced by any known plastidial TRX reduction system. Interestingly,  
17 we found that other plastidial TRX can reduce TRXz. This is thermodynamically  
18 feasible as these TRX have a redox midpoint potential ( $E_m$ ) less negative than the  $E_m$   
19 of TRXz (Collin et al., 2003; 2004; Chibani et al., 2011). This is the first example, to  
20 our knowledge, of a TRX reducing another TRX. Overall, these biochemical data  
21 underscoring that TRXz stands out from the other plastidial isoforms are consistent  
22 with the severe albino phenotype of the Arabidopsis *trxz* knock-out mutant (Arsova et  
23 al., 2010; Meng et al., 2010) contrasting with the mutants of the other plastidial TRX  
24 that do not exhibit a visible phenotype under standard growth conditions (Laugier et  
25 al., 2012).

1 TRXz was previously identified in a proteomic study as a component of  
2 transcriptionally active chromosome in plastids (pTAC; Pfalz et al., 2006) and more  
3 recently as a subunit of a complex formed with the plastid-encoded RNA polymerase  
4 (PEP; Schröter et al., 2010). *trxz* mutant or silenced plants are affected in the  
5 expression of the chloroplast genome, particularly the genes transcribed by the PEP  
6 (Arsova et al., 2010; Schröter et al., 2010). This strongly suggests the implication of  
7 TRXz in the transcription of the plastome. TRXz interacts with several targets,  
8 including 2 Cys-PRX, FNR and two proteins with similarities to plastid fructokinase-  
9 like (FLN) proteins (Schröter et al., 2010) previously identified as a component of  
10 pTAC (Pfalz et al., 2006) but with no fructokinase activity (Arsova et al., 2010). It has  
11 been shown that TRXz acts as an adaptor protein between the receptor-like protein Cf-  
12 9 and the Ser/Thr kinase ACIK1 in the cytosol of tomato (Nekrasov et al., 2006).  
13 Recently, Schröter et al. (2010) proposed that TRXz could play a similar adaptor role  
14 as a subunit of the PEP complex for regulation of plastid gene expression. TRXz  
15 would be an additional key player of photosynthetic gene expression (Dietz and  
16 Pfannschmidt, 2011).

17 Overall, this points to a unique role of TRXz, which indeed possesses TRX  
18 activity. TRXz appears as an important sensor of the redox status of the chloroplast  
19 being potentially oxidized by 2-Cys PRX (sensing reactive oxygen species) and  
20 reduced by other TRX (sensing light indirectly) revealing its potential important role  
21 in the redox homeostasis of the chloroplast.

## **Supplementary material**

Supplementary data are available at *JXB* online.

Table S1: List of specific primers used for RT-qPCR and experimental procedure.

Table S2: Summary of the different proteins used and purification procedures.

Figure S1: Anti-TRXz antibody specificity.

Figure S2: Expression patterns of plastidial thioredoxins and reductases from the « Arabidopsis eFP browser » website.

Figure S3: Reduction of H<sub>2</sub>O<sub>2</sub> by 2-Cys PRX using NTRC as an electron donor.

Figure S4: Reduction of plastidial TRX by NTRs.

Figure S5: Reduction of TRXz by various reductants.



## **Acknowledgments**

The authors are grateful to Dr N. Rouhier for providing GRXS16 protein, to Dr M. Zaffagnini for technical advices and scientific discussions and to Dr M. Miginiac-Maslow for support, helpful suggestions and critical reading of the manuscript.

## REFERENCES

- Alkhalifioui F, Renard M, Montrichard F.** 2007. Unique properties of NADP-thioredoxin reductase C in legumes. *Journal of Experimental Botany* **58**: 969-978
- Arsova B, Hoja U, Wimmelbacher M, Greiner E, Ustun S, Melzer M, Petersen K, Lein W, Bornke F.** 2010. Plastidial thioredoxin z interacts with two fructokinase-like proteins in a thiol-dependent manner: evidence for an essential role in chloroplast development in *Arabidopsis* and *Nicotiana benthamiana*. *Plant Cell* **22**: 1498-1515
- Balmer Y, Vensel WH, Cai N, Manieri W, Schürmann P, Hurkman WJ, Buchanan BB.** 2006. A complete ferredoxin/thioredoxin system regulates fundamental processes in amyloplasts. *The Proceedings of the National Academy in Sciences USA* **103**: 2988-2993
- Bandyopadhyay S, Gama F, Molina-Navarro MM, Gualberto JM, Claxton R, Naik SG, Huynh BH, Herrero E, Jacquot J-P, Johnson M K, Rouhier N.** 2008. Chloroplast monothiol glutaredoxins as scaffold proteins for the assembly and delivery of [2Fe-2S] clusters. *The EMBO Journal* **27**: 1122-1133
- Benz JP, Lintala M, Soll J, Mulo P, Bolter B.** 2010. A new concept for ferredoxin NADP(H) oxidoreductase binding to plant thylakoids. *Trends in Plant Science* **15**: 608-613
- Boyer ME, Wang CW, Swartz JR.** 2006. Simultaneous expression and maturation of the iron-sulfur protein ferredoxin in a cell-free system. *Biotechnology and Bioengineering* **94**: 128-138
- Broin M, Cuine S, Eymery F, Rey P.** 2002. The plastidic 2-cysteine peroxiredoxin is a target for a thioredoxin involved in the protection of the photosynthetic apparatus against oxidative damage. *Plant Cell* **14**: 1417-1432
- Buchanan BB.** 1980. Role of light in the regulation of chloroplast enzymes. *Annual Review of Plant Physiology and Plant Molecular Biology* **31**: 341-374
- Chibani K, Tarrago L, Schürmann P, Jacquot J-P, Rouhier N.** 2011. Biochemical properties of poplar thioredoxin z. *FEBS Letters* **585**: 1077-1081
- Collin V, Issakidis-Bourguet E, Marchand C, Hirasawa M, Lancelin JM, Knaff DB, Miginiac-Maslow M.** 2003. The *Arabidopsis* plastidial thioredoxins - New functions and new insights into specificity. *The Journal of Biological Chemistry* **278**: 23747-23752
- Collin V, Lamkemeyer P, Miginiac-Maslow M, Hirasawa M, Knaff DB, Dietz K-J, Issakidis-Bourguet E.** 2004. Characterization of plastidial thioredoxins from *Arabidopsis* belonging to the new y-type. *Plant Physiology* **136**: 4088-4095

**Czechowski T, Stitt M, Altmann T, Udvardi MK, Scheible W-R.** 2005. Genome-wide identification of superior reference genes for transcript normalization in Arabidopsis. *Plant Physiology* **139**: 5-17

**Dietz K-J and Pfannschmidt T.** 2011. Novel regulators in photosynthetic redox control of plant metabolism and gene expression. *Plant Physiology* **155**: 1477-1485

**Droux M, Miginiac-Maslow M, Jacquot JP, Gadal P, Crawford NA, Kosower NS, Buchanan BB.** 1987. Ferredoxin-thioredoxin reductase - a catalytically active dithiol group links photoreduced ferredoxin to thioredoxin functional in photosynthetic enzyme regulation. *Archives of Biochemistry and Biophysics* **256**: 372-380

**Gelhaye E, Rouhier N, Jacquot JP.** 2003. Evidence for a subgroup of thioredoxin h that requires GSH/Grx for its reduction. *FEBS Letters* **555**: 443-448

**Hanke GT, Kimata-Arigo Y, Taniguchi I, Hase T.** 2004. A post genomic characterization of arabidopsis ferredoxins. *Plant Physiology* **134**: 255-264

**Issakidis E, Saarinen M, Decottignies P, Jacquot JP, Crépin C, Gadal P, Miginiac-Maslow M.** 1994. Identification and characterization of the 2nd regulatory disulfide bridge of recombinant sorghum leaf NADP-malate dehydrogenase. *The Journal of Biological Chemistry* **269**: 3511-3517

**Issakidis-Bourguet E, Mouaheb N, Meyer Y, Miginiac-Maslow M.** 2001. Heterologous complementation of yeast reveals a new putative function for chloroplast m-type thioredoxin. *The Plant Journal* **25**: 127-135

**Jacquot J-P, Rivera-Madrid R, Marinho P, Kollarova M, Le Maréchal P, Miginiac-Maslow M, Meyer Y.** 1994. *Arabidopsis thaliana* NADPH thioredoxin reductase cDNA characterization and expression of the recombinant protein in *Escherichia-coli*. *Journal of Molecular Biology* **235**: 1357-1363

**Johansson K, Ramaswamy S, Saarinen M, Lemaire-Chamley M, Issakidis-Bourguet E, Miginiac-Maslow M, Eklund H.** 1999. Structural basis for light activation of a chloroplast enzyme: The structure of sorghum NADP-malate dehydrogenase in its oxidized form. *Biochemistry* **38**: 4319-4326

**Keryer E, Collin V, Lavergne D, Lemaire SD, Issakidis-Bourguet E.** 2004. Characterization of Arabidopsis T-DNA mutants for the variable subunit of ferredoxin:thioredoxin reductase. *Photosynthesis Research* **79**: 265-274

**König J, Baier M, Horling F, Kahmann U, Harris G, Schürmann P, Dietz K-J.** 2002. The plant-specific function of 2-Cys peroxiredoxin-mediated detoxification of peroxides in the redox-hierarchy of photosynthetic electron flux. *The Proceedings of the National Academy in Sciences USA* **99**: 5738-5743

**Laloi C, Rayapuram N, Chartier Y, Grienenberger JM, Bonnard G, Meyer Y.** 2001. Identification and characterization of a mitochondrial thioredoxin system in plants. *The Proceedings of the National Academy in Sciences USA* **98**: 14144-14149

**Lamkemeyer P, Laxa M, Collin V, Li W, Finkemeier I, Schottler MA, Holtkamp V, Tognetti VB, Issakidis-Bourguet E, Kandlbinder A, Weis E, Miginiac-Maslow M, Dietz K-J.** 2006. Peroxiredoxin Q of *Arabidopsis thaliana* is attached to the thylakoids and functions in context of photosynthesis. *Plant Journal* **45**: 968-981

**Laugier E, Tarrago L, Courteille A, Innocenti G, Eymery F, Rumeau D, Issakidis-Bourguet E, Rey P.** 2012. Involvement of thioredoxin  $\gamma 2$  in the preservation of leaf methionine sulfoxide reductase capacity and growth under high light. *Plant Cell and Environment*. doi: 10.1111/PCE.12005

**Lemaire SD, Michelet L, Zaffagnini M, Massot V, Issakidis-Bourguet E.** 2007. Thioredoxins in chloroplasts. *Current Genetics* **51**: 343-365

**Livak KJ, Schmittgen TD.** 2001. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods* **25**: 402-408

**Marchand C, Vanacker H, Collin V, Issakidis-Bourguet E, Le Maréchal P, Decottignies P.** 2010. Thioredoxins targets in *Arabidopsis* roots. *Proteomics* **10**: 2418-2428

**Meng L, Wong JH, Feldman LJ, Lemaux PG, Buchanan BB.** 2010. A membrane associated thioredoxin required for plant growth moves from cell to cell, suggestive of a role in intercellular communication. *The Proceedings of the National Academy in Sciences USA* **107**: 3900-3905

**Mestres-Ortega D, Meyer Y.** 1999. The *Arabidopsis thaliana* genome encodes at least four thioredoxins m and a new prokaryotic-like thioredoxin. *Gene* **240**: 307-316

**Meyer Y, Reichheld JP, Vignols F.** 2005. Thioredoxins in *Arabidopsis* and other plants. *Photosynthesis Research* **86**: 419-433

**Michelet L, Zaffagnini M, Massot V, Keryer E, Vanacker H, Miginiac-Maslow M, Issakidis-Bourguet E, Lemaire SD.** 2006. Thioredoxins, glutaredoxins, and glutathionylation: new crosstalks to explore. *Photosynthesis Research* **89**: 225-245

**Michelet L, Zaffagnini M, Marchand C, Collin V, Decottignies P, Tsan P, Lancelin JM, Trost P, Miginiac-Maslow M, Noctor G, Lemaire SD.** 2005. Glutathionylation of chloroplast thioredoxin f is a redox signaling mechanism in plants. *The Proceedings of the National Academy in Sciences USA* **102**: 16478-16483

**Née G, Zaffagnini M, Trost P, Issakidis-Bourguet E.** 2009. Redox regulation of chloroplastic glucose-6-phosphate dehydrogenase: A new role for f-type thioredoxin. *FEBS Letters* **583**: 2827-2832

**Nekrasov V, Ludwig AA, Jones JDG.** 2006. CITRX thioredoxin is a putative adaptor protein connecting Cf-9 and the ACIK1 protein kinase during the Cf-9/Avr9-induced defence response. *FEBS Letters* **580**: 4236-4241

**Noctor G.** 2006. Metabolic signalling in defence and stress: the central roles of soluble redox couples. *Plant, Cell & Environment* **29**: 409-425

**Nuruzzaman M, Gupta M, Zhang C, Wang L, Xie W, Xiong L, Zhang Q, Lian X.** 2008. Sequence and analysis of the thioredoxin gene family in rice. *Molecular Genetics & Genomics* **280**: 139-151

**Pfalz J, Liere K, Kandlbinder A, Dietz K-J, Oelmüller R.** 2006. pTAC2, -6 and -12 are components of the transcriptionally active plastid chromosome that are required for plastid gene expression. *Plant Cell* **18**: 176-197

**Pérez-Ruiz JM, Spinola MC, Kirchsteiger K, Moreno J, Sahrawy M, Cejudo FJ.** 2006. Rice NTRC is a high-efficiency redox system for chloroplast protection against oxidative damage. *Plant Cell* **18**: 2356-2368

**Reichheld J-P, Meyer E, Khafif M, Bonnard G, Meyer Y.** 2005. AtNTRB is the major mitochondrial thioredoxin reductase in *Arabidopsis thaliana*. *FEBS Letters* **579**: 337-342

**Rozen S, Skaletsky HJ.** 2000. Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds) *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press, Totowa, NJ, pp 365-386

**Ruelland E, Lemaire Chamley M, Le Marechal P, Issakidis-Bourguet E, Djukic N, Miginiac-Maslow M.** 1997. An internal cysteine is involved in the thioredoxin-dependent activation of sorghum leaf NADP-malate dehydrogenase. *The Journal of Biological Chemistry* **272**: 19851-19857

**Ruelland E, Miginiac-Maslow M.** 1999. Regulation of chloroplast enzyme activities by thioredoxins: activation or relief from inhibition? *Trends in Plant Science* **4**: 136-141

**Schröter Y, Steiner S, Matthai K, Pfannschmidt T.** 2010. Analysis of oligomeric protein complexes in the chloroplast sub-proteome of nucleic acid-binding proteins from mustard reveals potential redox regulators of plastid gene expression. *Proteomics* **10**: 2191-2204

**Serrato AJ, Pérez-Ruiz JM, Spinola MC, Cejudo FJ.** 2004. A novel NADPH thioredoxin reductase, localized in the chloroplast, which deficiency causes hypersensitivity to abiotic stress in *Arabidopsis thaliana*. *The Journal of Biological Chemistry* **279**: 43821-43827

**Traverso JA, Vignols F, Cazalis R, Serrato AJ, Pulido P, Sahrawy M, Meyer Y, Cejudo FJ, Chueca A.** 2008. Immunocytochemical localization of *Pisum sativum* TRXs f and m in non-photosynthetic tissues. *Journal of Experimental Botany* **59**: 1267-1277

**Valerio C, Costa A, Marri L, Issakidis-Bourguet E, Pupillo P, Trost P, Sparla F.** 2010. Thioredoxin-regulated  $\beta$ -amylase (BAM1) triggers diurnal starch

degradation in guard cells, and in mesophyll cells under osmotic stress. *Journal of Experimental Botany* **62**: 545-555

**Vieira Dos Santos C, Laugier E, Tarrago L, Massot V, Issakidis-Bourguet E, Rouhier N, Rey P.** 2007. Specificity of thioredoxins and glutaredoxins as electron donors to two distinct classes of Arabidopsis plastidial methionine sulfoxide reductases B. *FEBS Letters* **581**: 4371-4376

## Table 1

**Table 1.** Activation test of mutant NADP-MDH by TRXz.

Mutant NADP-MDH (0.9  $\mu\text{g}$ ) was incubated with 100 mM DTT and various concentrations of TRXz. Activation by TRXm1 was used as positive control. Activity was measured spectrophotometrically at 340 nm through the oxaloacetate-dependent disappearance of NADPH. Activation treatments were made at room temperature for 20 min in presence of DTT (100 mM) alone or with TRX (5  $\mu\text{M}$  unless otherwise mentioned).

	<b>Initial rates of reduction</b> ( $\mu\text{moles oxidized NADPH}$ $\cdot\text{mg}^{-1}\text{ NADP-MDH}\cdot\text{sec}^{-1}$ )
DTT alone	$0.71 \pm 0.06$
TRX m1	$1.36 \pm 0.09$
TRX z	$0.73 \pm 0.07$
100 $\mu\text{M}$ TRX z	$0.79 \pm 0.10$

## Legends for figures

**Figure 1.** Expression of plastidial TRX and TRX reductases in Arabidopsis.

RT-qPCR analyses were performed on total RNA isolated from young seedlings, leaves, flowers, young and mature siliques, seeds and roots. Transcript levels are the result of three independent experiments and were normalized by the  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen, 2001) using the PP2A gene as standard (Czechowski et al., 2005). A, expression patterns of plastidial TRX genes, *trxf1* transcripts level in leaves used as calibrator. B, expression patterns of *ntrc* and *ftc* subunits genes, *ntrc* transcripts level in leaves used as calibrator.

**Figure 2.** Efficiency of TRXz in the reduction of 2-Cys PRX compared to other plastidial TRX.

The enzyme 2-Cys PRXA (20  $\mu\text{M}$ ) was incubated in the presence of 400  $\mu\text{M}$  DTT, 600  $\mu\text{M}$   $\text{H}_2\text{O}_2$  and 10  $\mu\text{M}$  TRX. As a control, 2-Cys PRX was incubated without TRX (DTT). Data are the mean of 3 independent experiments. Error bars correspond to standard deviation.

**Figure 3.** Activation of NADP-MDH by plastidial TRXf, x and y in the light, in a Fd/TRX reconstituted system.

Mutant NADP-MDH was illuminated at 120  $\mu\text{E m}^{-2} \text{s}^{-1}$  in the presence of freshly prepared thylakoids (equivalent to 20  $\mu\text{g}$  chlorophyll), 5  $\mu\text{M}$  ferredoxin, 2  $\mu\text{M}$  FTR, and 25  $\mu\text{M}$  TRXf1 (f1, black circles), TRXy1 (y1, black diamonds), TRXy2 (y2,



black triangles) or TRXx (x, black squares). TRXy2 was also tested in the dark (open triangles). Experiments were reproduced 3 times, a representative set of data is shown.

**Figure 4.** Redox and oligomerization state of TRXz.

TRXz was alkylated with  $^{14}\text{C}$ -IAM after an oxidizing (DTTox 10 mM) or a reducing treatment (DTTred 100 mM). The samples were separated by non-reducing SDS-PAGE. Proteins and radiolabeling were respectively visualized by Coomassie blue staining and exposure of the dried gel to a low energy imaging screen.

**Figure 5.** Reduction test of TRXz by FTR in a reconstituted light-driven Fd/TRX system.

Experimental conditions were as described in legend of Fig. 3. As controls, oxidized TRXz (10  $\mu\text{M}$ ) was incubated with (lanes 2 and 3) or without (lane 1) 100 mM DTT. For the reduction test of TRXz by FTR, TRXz (25  $\mu\text{M}$ ) was incubated in the presence of thylakoids, 5  $\mu\text{M}$  Fd and 10  $\mu\text{M}$  FTR in the light. Aliquots were withdrawn periodically, the proteins were alkylated, subjected to non-reducing SDS-PAGE, electrotransferred onto nitrocellulose membrane and probed with anti *Strep*-tag antibody. Experiment was reproduced 4 times, a representative experiment is shown.

**Figure 6.** Reduction of TRXz by other plastidial TRX.

TRXz (10  $\mu$ M) was incubated without (lane 1) or with DTT (1, 10 or 100 mM, lane 2, 7, 8 respectively) or in presence of 10  $\mu$ M of another plastidial TRX (f1, m1: panel A; x, y1: panel B) without (lane 3, 5) or with DTT (1 mM, lane 4, 6), for 15 min at room temperature. Proteins were then alkylated and subjected to non-reducing SDS-PAGE, electrotransferred onto nitrocellulose membrane and probed with an anti-TRXz antibody. Experiments were reproduced four times, a representative experiment is shown.