

Two decades of seawater acidification experiments on tropical scleractinian corals: Overview, meta-analysis and perspectives

Mathilde Godefroid, Sam Dupont, Marc Metian, Laetitia Hédouin

▶ To cite this version:

Mathilde Godefroid, Sam Dupont, Marc Metian, Laetitia Hédouin. Two decades of seawater acidification experiments on tropical scleractinian corals: Overview, meta-analysis and perspectives. Marine Pollution Bulletin, 2022, 178, pp.113552. 10.1016/j.marpolbul.2022.113552. hal-03841318

HAL Id: hal-03841318 https://univ-perp.hal.science/hal-03841318v1

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 **Title** 2 Two decades of seawater acidification experiments on tropical scleractinian corals: overview, meta-3 analysis and perspectives 4 5 **Authors** 6 Mathilde Godefroid^{1,2,*}, Sam Dupont^{3,4}, Marc Metian⁴, Laetitia Hédouin^{1,2} 7 8 **Affiliations** 9 ¹PSL Research University: EPHE-CNRS-UPVD, USR 3278 CRIOBE, BP 1013, 98729 Papetoai, Mo'orea, 10 French Polynesia 11 ²Laboratoire d'Excellence « CORAIL», Mo'orea, French Polynesia 12 ³Department of Biological and Environmental Sciences, University of Gothenburg, Kristineberg 13 Marine Research Station, Kristineberg 566, 45178 Fiskebäckskil, Sweden 14 ⁴ Radioecology Laboratory International Atomic Energy Agency (IAEA), Marine Laboratories, 4 Quai 15 Antoine 1er, 98000, Principality of Monaco 16 17 18 *Corresponding author: 19 Name: Godefroid Mathilde 20 Present address: Marine biology laboratory, 21 Université Libre de Bruxelles, 22 Av. F.D. Roosevelt, 50, CP 160/15 23 1050 Bruxelles 24 Belgium 25 (+32)489 36 70 08 Phone: 26 E-mail: godefroid.mathilde1@gmail.com

Abstract

Ocean acidification has emerged as a major concern in the last fifteen years and studies on the impacts of seawater acidification on marine organisms have multiplied accordingly. This review aimed at synthesizing the literature on the effects of seawater acidification on tropical scleractinians under laboratory-controlled conditions. We identified 141 articles (published between 1999-2021) and separated endpoints into 22 biological categories to identify global trends for mitigation and gaps in knowledge and research priorities for future investigators. The relative number of affected endpoints increased with pH intensity (particularly for endpoints associated to calcification and reproduction). When exposed to pH 7.6-7.8 (compared to higher pH), 49% of endpoints were affected. The diversity in experimental designs prevented deciphering the modulating role of coral life stages, genera or duration of exposure. Finally, important bias in research efforts included most experiments on adult corals (68.5%), in 27 out of 150 (18%) coral ecoregions and exclusively from shallow-waters.

Keywords

Ocean acidification, Tropical scleractinian, pH, Calcification, Review

Acknowledgements

- 45 We thank Jacob et al. (2020) for making their R scripts available and Dr Moreau C. for his help in
- 46 making Fig. 1. This work was funded by the Foundation de France for a project called "ACID REEFS",
- 47 by Ministère de la Transition Écologique et Solidaire and the Foundation for Research on Biodiversity
- 48 (FRB) for a project entitled "ACID REEFS". The IAEA is grateful for the support provided to its
- 49 Environment Laboratories by the Government of the Principality of Monaco.

1. Introduction

 Due to the increasing emissions of carbon dioxide into the atmosphere, the Ocean is warming and getting more acidic. Ocean acidification (OA) has been identified as a global environmental threat and included as the United Nations' Sustainable Development Goal 14.3, as well as one of the nine planetary boundaries (Jagers et al., 2019; Rockström et al., 2009). OA impacts on marine species and ecosystems are well documented, including effect on marine calcifiers, threatening coral reefs as well as broader marine ecosystems (Doney et al., 2009; Feely, 2004; Gattuso & Hansson, 2011; Hendriks et al., 2010; Jiang et al., 2019; Kroeker et al., 2010, 2013; Orr et al., 2005).

Most of the available knowledge on the effects of OA on marine organisms comes from short-term laboratory experiments on isolated organisms (Kroeker et al., 2010). Fixed-term studies have the disadvantage of potentially under/overestimate the effects of OA, as some taxa may show vulnerability/acclimatization in the long-term (Fantazzini et al., 2015). Similarly, other life stages than those considered may show different vulnerability and controlled conditions does not consider the indirect effects due to OA-driven ecological changes (Fabricius et al., 2011). In this context, knowledge acquired through field experiments taking advantage of organisms that are naturally exposed to OA (e.g. CO₂ vent systems) are very complementary as they account for the life-long acclimatization of organisms. These studies already showed that OA may change reef community composition and metabolism (Biscéré et al., 2019; Noonan et al., 2018). They also reported effects on skeletal porosity (Prada et al., 2021) and a variety of responses to OA on calcification rate, suggesting species-specific acclimatization to OA (Strahl et al., 2015).

Coral reefs have received particular attention as they are among the most severely threatened ecosystem on Earth (Pandolfi, 2003; Raven, 2005). The effects of seawater acidification on corals have been extensively studied (Chan & Connolly, 2013; Erez et al., 2011), but responses often differ between studies depending on parameters such as tested populations, species and life-cycle stages (Kawahata et al., 2019). These apparent conflicting results in the literature could also be attributed to variations in experimental designs and methodologies. In this study, we reviewed the literature that examined the effects of decreased pH on tropical scleractinian corals under laboratory-controlled conditions. A meta-analysis based on 169 experiments conducted in 141 peer-reviewed articles published since 1999 was performed, with four main objectives:

- (1) To provide a semi-quantitative description of the evolution of research efforts testing the effects of seawater acidification on tropical scleractinian corals,
- (2) To evaluate the effect of seawater acidification on coral biological functions,

- 82 (3) To investigate the modulating influence of exposure duration, coral life stages and coral genera on sensitivity to seawater acidification and,
- 84 (4) To highlight research gaps and bias.

Based on this review, we provide recommendations and perspectives for future research and an updated baseline for scientists starting in this field of research.

87

88

2. Material and methods

89 The code used in this manuscript based Jacob et al. (2020): was on 90 https://doi.org/10.5281/zenodo.3694955.

91

92

93 94

95

96

97

98

99

100

101

102

103104

105

106

107

108

2.1 Literature search

A literature search was performed using the following databases: Ocean Acidification – International Coordination Centre portal for ocean acidification biological response data, Google Scholar, Scopus (keywords such as "coral" AND "acidification"). We listed all experimental studies published before 2021, which mainly focused on the effects of experimental seawater acidification on tropical scleractinian corals. Studies were selected according to the following criteria: 1) the experiment(s) was(were) performed under laboratory-controlled conditions; 2) one treatment (at least) was designed to test the effects of seawater acidification only (without interaction with another stressor); 3) a tropical scleractinian coral species (at least) was studied; 4) an acidified treatment was compared with a control treatment (present-day conditions), with constant conditions (i.e. non-oscillating). In situ experiment(s) of the effects of seawater acidification on biological functions were not included in the database but were considered for the discussion. Higher than present-day (control) pH conditions (e.g. pre-industrial pH conditions) were not included in the database. When the statistics lacked clarity/transparency or did not allow to be adequately decomposed into the database, the article was excluded. The completeness of the results obtained was considered as satisfactory based on "snowballing" (i.e. checking citations on reference lists of relevant articles until no further relevant article could be found; Sayers, 2007).

109

110

2.2 Main data extraction

For each experiment selected in the database, the following information was extracted: Study location(s), study duration(s), coral genus and species, coral life stage(s), number of studied species, tested pHs, pH scale and endpoints.

2.3 Data organization

For analyses, six pH classes were defined: Class 1 for pH between 7.8 and 8, Class 2 for pH between 7.6 and 7.8, Class 3 for pH between 7.4 and 7.6, Class 4 for pH between 7.2 and 7.4, Class 5 for pH between 7 and 7.2, Class 6 for pH below 7. Recent studies highlight the importance of local adaptation to the present natural variability in the carbonate chemistry to define species sensitivity to OA (e.g. Vargas et al., 2017). As a consequence, the definition of pH scenarios or categories relevant in the context of OA should consider this natural variability. As diurnal and seasonal pH variability is low in coral reef environments (e.g. Cyronak et al., 2020), pH within class 1 to 6 are considered relevant in the context of OA. Similarly, duration, genus and life stage were also divided into classes. Four duration classes were defined: Class 1 for very short-term experiments (<1 week), Class 2 for short-term experiments (1-4 week(s)), Class 3 for medium-term experiments (1-3 month(s)), Class 4 for long-term experiments (>3 months); Three genus classes: Class 1 for species from the genus *Pocillopora*; and three life stage classes were chosen: Class 1 for free-living non-calcifying early life stages (Gametes, Gastrulae, Larvae), Class 2 for settled calcifying early life stages (Recruits, Primary polyps, Juveniles) and Class 3 for the adults. All classes are summarized in Table 1.

Table 1. pH, Duration, genus and life stage classes used to classify this review.

Parameters	Classes	
	Class 1	7.8-8
	Class 2	7.6-7.8
рH	Class 3	7.4-7.6
F	Class 4	7.2-7.4
	Class 5	7.0-7.2
	Class 6	<7
	Class 1	<1 week
Duration	Class 2	1-4 week(s)
	Class 3	1-3 month(s)
	Class 4	>3 months
	Class 1	Acropora
Genus	Class 2	Porites
	Class 3	Pocillopora
	Class 1	Free-living early life stages
Life stage	Class 2	Settled early life stages
	Class 3	Adults

2.4 Endpoints

For each study and experiment, all measured endpoints were identified and divided into categories based on their biological significance. When the biological significance of an endpoint was not straightforward and/or when an endpoint could have several biological meanings, we attributed it to a single category based on the focus adopted by authors. In total, eight categories, divided into 22 subcategories, were established (Table 2).

Table 2. List of the biological categories and subcategories used in the database, with one example of endpoint measured.

Biological category	Biological subcatergory	Endpoint measured ¹
Fooding and everation	Excretion	Ammonia excretion rate
Feeding and excretion	Feeding	Feeding rate
	Reproductive investment and success	Settlement rate
Fitness	Calcification	Net calcification rate
Fitness	Mortality	Total mortality
	Growth and development	Linear extension rate
Cellular stress-response	Apoptosis	p53 expression level
•	Oxidative stress	Glutathione reductase expression level
mechanisms	Other molecular actors	HSP70
	Carbohydrate	Carbohydrate content
Diamalagulas	Lipid	Total lipid content
Biomolecules	Protein	Protein content
	Energy content	Host ATP content
Desmination and	Photophysiology	Net photosynthetic rate
Respiration and	Respiration	Dark respiration rate
photophysiology	Productivity	Net productivity
Microbiome and endolithic algae	Bacterial community composition	Gammaproteobacteria content
Metabolome		Metabolite profile
Epigenetic mechanism		DNA methylation level
	Community structure	Planar cover of the community
Community	Community calcification	Light community calcification
	Respiration	Community net oxygen consumption

¹This is a non-exhaustive list of examples. See the database in Supplementary Material for the full list of endpoints measured.

For each endpoint, we reported the occurrence of a significant effect of exposure to pH as compared to the highest tested pH: YES if the endpoint was significantly affected by pH; NO elsewhere (Jacob et

al., 2020). If multiple pH treatments were used and the statistics in the study did not compare the effect of each treatment with the control, but rather reported the result as a statistical trend, a YES was attributed to an endpoint when the trend was significant. For each category and subcategory, the percentage of affected endpoints was calculated by dividing the number of affected endpoints (*i.e.* number of YES) by the total number of endpoints in this category or subcategory (Jacob et al., 2020).

154

155

156

157158

148

149

150

151

152

153

2.5 Database construction

All selected experimental studies were grouped in a common database gathering all relevant information (main data, endpoints and effects as described from 2.2 to 2.4). The database is available as Supplementary Material.

159

160

161

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

3. Results & Discussion

3.1 General trends in experimental seawater acidification research on tropical scleractinian corals

162 Literature trend

In our database, we identified 169 experiments in 141 articles (listed in Supplementary Material) that have evaluated the effects seawater acidification under laboratory-controlled conditions on tropical scleractinian corals. The six best studied regions account for 69% of the studies: Australia (15%), Japan (14%), French Polynesia (14%), Hawaii (10%), the USA (excluding Hawaii, 8%) and Taiwan (8%; Fig. 2A). Six percent of the studies were carried out on corals that have been maintained and acclimated in aquaria for years (referred to as "Aquaria acclimated"; Fig. 2A). These included the species Stylophora pistillata that were held in aquaria in Monaco (Liew et al., 2018; Reynaud et al., 2003; Tambutté et al., 2015; Venn et al., 2013, 2019, 2020; Zoccola et al., 2016), but also by other coral species obtained from commercial suppliers (Bove et al., 2020; Wijgerde et al., 2014). Altogether, these data reveal that the effects of experimental seawater acidification on tropical scleractinian corals have only been studied in 27 out of 150 coral ecoregions (18%; Veron et al., 2005, 2015). Certain parts of the world remain unexplored, with no study to date in the African reefs nor around the coasts of India or Sri Lanka, and very few studies in the coral triangle (Figure 1), despite being known as the most biodiverse area in the ocean, containing more than 76% of the world's shallow-water reef-building corals (Asaad et al., 2018). Moreover, no study has ever focused on the impact of seawater acidification on tropical mesophotic corals, even though they represent 2/3rd of the world coral reefs (Morais & Santos, 2018) and are expected to act as "deep refugia", by providing a viable reproductive source for shallow reefs after disturbance (Bongaerts et al., 2010).

Experimental seawater acidification research on tropical scleractinians is relatively recent and has increased exponentially since the 21st century (Fig. 2B). Similarly, in situ studies have only started to develop even more recently, with a majority of studies (70%) published over the five past years (Fig. 2B). Among these, 80% took advantage of coral communities living under naturally acidified waters such as submarine springs in Mexico (Crook et al., 2012, 2013; Martinez et al., 2019; Wall et al., 2019) and CO₂ vents in Papua New Guinea (Biscéré et al., 2019; Fabricius et al., 2011; Fantazini et al., 2015; Marcelino et al., 2017; Morrow et al., 2015; Noonan et al., 2016, 2018; O'Brien et al., 2018; Prada et al., 2021; Smith et al., 2016; Strahl et al., 2015; Wall et al., 2017). Other studies compared adjacent reef sites with distinct pH conditions (Barkley et al., 2015, 2017; Camp et al., 2017). Finally, 20% of studies have performed in situ controlled CO2 enrichment such as FOCE experiments (Free Ocean CO₂ Experiments; Albright et al., 2016, 2018; Doo et al., 2019; Georgiou et al., 2015; Kline et al., 2019; Srednick et al., 2020). In situ studies possess the great advantage of being more ecologically realistic than laboratory-based studies, as they account for the complexity of species interactions of the whole reef community, under natural abiotic conditions (Albright et al., 2018). Yet, naturally acidified sites are not always available and possess their own set of limitations. They are not perfect predictors of future ocean ecology owing to temporal variability in pH, spatial proximity with other populations and changes in other parameters such as seawater temperature (Prada et al., 2021; Riebesell, 2008). Moreover, the manipulation of seawater parameters in situ is technically challenging and comes at great expenses, which greatly limits its reproducibility.

pH scale

Best practices recommend to measure seawater pH using the total hydrogen ion concentration scale (Total scale; Dickson et al., 2007). Yet, only 54.4% of experiments used this scale (Figure 2I). The rest of the experiments measured pH on the NBS scale (US National Bureau of Standards, now referred to as the IUPAC scale, 29.2%) or on the seawater scale (SWS; 5.3%). A minority of experiments (3.5%) did not measure the pH but estimated its value from the other seawater carbonate chemistry parameters, via CO2sys (Pierrot et al., 2006) or the R package seacarb (Gattuso et al., 2020). Finally, a significant proportion (7.6%) did not specify how the pH was measured/calculated. Such heterogeneity in the pH scales used to measure seawater pH probably lies in the fact that it is, together with the study of acid-base reactions in seawater, one of the most confused areas of marine chemistry (Dickson, 1993). Above all, it is essential that the pH is defined on the same pH scale as

that of all acid-dissociation constants that are used with it (Riebesell et al., 2011). We highly recommend future studies to focus on measuring total scale pH following the complete procedure and best practices in SOP6a and SOP6b (Dickson et al., 2007).

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

212

213

214

Coral characteristics

The vast majority of studies have evaluated the effects of seawater acidification on adult scleractinians (68.4%; Fig. 2C). 16.7% of studies were devoted to free-living (non-calcified) early life stages (mainly larvae -13.9%-, followed by gastrulae -1.7%- and gametes -1.1%). Then, the most studied settled (calcified) early life stages were recruits (7.8%), followed by primary polyps (4.4%) and juveniles (2.8%; Fig. 2C). This focus on adult corals is a common trend also generally observed for other marine organisms (Kroeker et al., 2010). For corals, the high attention devoted to adult stages is likely due to the convenience of collection and husbandry, compared to early life stages, which require specific knowledge on spawning events and high expertise for a successful rearing. However, future studies should put greater emphasis on early life stages as they are often inferred to be more vulnerable to environmental conditions than adults (Byrne, 2011). However, as only 4.7% of studies considered multiple life history stages, the susceptibility among life stages remains largely underexplored. Multigenerational studies are thus crucial as they allow to compare the sensitivity of a species among life stages and to evaluate transgenerational carry over effects. To the best of our knowledge, very few studies have evaluated these effects with corals, as compared to other taxa (e.g. mollusks; Hettinger et al., 2012; Parker et al., 2012) or other stressors (e.g. global warming; Dixon et al., 2015). Studies on scleractinians suggested that exposure of the parents to stressful conditions had positive influence on coral offspring performances, when subsequently re-exposed (e.g. metabolic acclimation for P. damicornis, Putnam & Gates, 2015; greater larval settlement and survivorship for P. damicornis, Putnam, 2021). Yet, this likely vary between species (e.g. S. pistillata, Bellworthy et al., 2018). Multigenerational studies are thus required to investigate parental (carryover) effects, to decipher whether the exposure to seawater acidification in adults enhance offspring tolerance, and to predict the future of coral reefs more accurately (Putnam et al., 2016, 2021).

Altogether, three genera stand for 65.1% of the experiments, with the most studied being *Acropora* (30.4%), followed by *Porites* (18.3%) and *Pocillopora* (16.3%; Fig. 2D). The most studied species from the database is *Acropora millepora*. Other well studied genera are *Montipora* (7.9%), *Stylophora* (6.3%) and *Seriatopora* (4.6%), followed by studies that have evaluated the effects of seawater acidification on the coral community (3.8%; Fig. 2D). The genera *Siderastrea*, *Pavona* and *Psammocora* each accounted for 1.7, 1.7 and 1.3% of the database, respectively. Finally, 6.7% of

experiments were performed on less studied genera (<1%, referred to as "Others"; Fig. 2D). These include species from the genera *Cyphastrea*, *Duncanopsammia*, *Favia*, *Favites*, *Fungia*, *Galaxea*, *Goniopora*, *Isopora*, *Leptastrea*, *Litophyllon*, *Madracis*, *Montastrea*, *Mussismila*, *Orbicella*, *Pseudodiploria* and *Turbinaria* (Fig. 2D). Overall, these data show that a small number of coral species are favoured in experimental seawater acidification experiments. As previously reported for heat stress experiments on corals (review by McLachlan et al., 2020), this may be due to the facility of working on highly studied species (*i.e.* with the availability of their genome), the presence of these species within each region or to logistical constraints (e.g. permits, easiness of maintenance in captivity). This finding also means that many species, and so several functional traits which are likely to drive differential responses to acidification, remain uninvestigated.

Experimental design

The majority of experiments (63.9%) considered a single species of coral (Fig. 2E). The remaining experiments used two to four (30.8%) or five to eight species (3.0%) and only 2.3% evaluated the effects of seawater acidification on the coral community (Fig. 2E). Undoubtedly, it limits and oversimplifies our understanding of the effects of OA on the biologically complex assemblage characterizing coral reefs. On top of that, species interactions can also decrease the predictability in species response to seawater acidification, since different pairings and spatial organization of corals might significantly influence the observed endpoints (Brien et al., 2016; Evensen et al., 2015, 2016). For these reasons, designing experiments based on the local species assemblage in terms of diversity, abundance and spatial organization, should be prioritized in future investigations.

Furthermore, most experiments lasted between 1 and 4 week(s) (39.6%), with the other experiments lasting less than a week (26.4%), one to three month(s) (24.7%) and more than three months (9.3%), for a minority (Fig. 2F). This trend is likely explained by the constraints associated with long-term studies, in terms of time, logistical or even financial limitations. Nevertheless, the duration of exposure may influence coral sensitivity to seawater acidification. For instance, corals are able to counteract the negative effects of acidification for a specific period through energy-demanding processes, until reaching a time-dependent threshold beyond which it will no longer be able to counter these effects (Castillo et al., 2014; Schoepf et al., 2013). For these reasons, short-term studies should be avoided, and long-term experimental studies prioritized, when possible. In that aspect, field experimentations/observations at CO₂ vents have the benefit to provide natural long-term studies (e.g. Biscéré et al., 2019; Fabricius et al., 2011; Martinez et al., 2019; Morrow et al., 2015; Noonan et al., 2018; Noonan & Fabricius, 2016; Prada et al., 2021; Strahl et al., 2015).

Endpoints

Most experiments worked on the effects of seawater acidification on the fitness (57.3%), respiration and photophysiology (23.9%) and biomolecules (7.1%) of tropical scleractinians (Fig. 2G). The other endpoints studied were involved in cellular stress response mechanism(s) (4.3%), community response(s) (3.5%), microbiome and endolithic algae composition (2.0%), epigenetic effect(s) (0.8%) and/or feeding and excretion (0.8%; Fig. 2G). Among those that have worked on the effects of seawater acidification on coral fitness, calcification was the most studied endpoint (52.6%), followed by growth and development (25.5%), reproductive investment and success (10.9%) and mortality (10.9%; Fig. 2H).

3.2 Biological effects of pH on tropical scleractinian corals

General trends

Overall, the relative number of affected endpoints increases with decreasing pH (Fig. 3). Twenty-six percent of endpoints were affected between pH 7.8 and 8, 46.2% between pH 7.6 and 7.8, 48.3% between pH 7.4 and 7.6, 46.3% between pH 7.2 and 7.4, 96.4% between pH 7.0 and 7.2 and 100% below pH 7.0. However, it should be noted that the number of available articles and data is not the same for each pH category. Studies have mainly focused on pH 7.8-7.6 (n endpoints=608), followed by pH 8.0-7.8 (n=519), pH 7.6-7.4 (n=203), pH 7.4-7.2 (n=162), pH 7.2-7.0 (n=56) and below 7.0 (n=4). Similarly, differences in experimental designs within each pH category is potentially a strong confounding factor, mixing results from experiments of multiple duration of exposure, coral lifestages, coral species, among others. This may particularly limit our interpretations for the less-studied pH categories.

Fitness

The decrease in pH and $[CO_3^{2-}]$ through OA is believed to increase the energetic cost of calcification for marine calcifiers (Gattuso & Hansson, 2011). Known as one of the biggest threats related to OA, calcification, the ability to build and maintain calcium carbonate structures, was among the most studied parameters (52.6% of studies). Results showed that 43.4% of endpoints related to calcification were affected for pH 7.8-8.0 (n=152), 55.3% for pH 7.6-7.8 (n=244), 66.7% for pH 7.4-7.6 (n=96), 85.7% for pH 7.2-7.4 (n=42), 96.9% for pH 7.0-7.2 (n=33) and 100% for pH below 7.0 (n=2) (Fig. 3). Similarly, growth and development (*i.e.* coral linear extension, biomass, larval size, recruits

diameter, etc.) were increasingly affected with decreasing pH, with 29.5% of affected endpoints for pH 7.8-8.0 (n=44), 44.4% for pH 7.6-7.8 (n=63), 61.1% for pH 7.4-7.6 (n=18), 73.7% for pH 7.2-7.4 (n=19), 88.9% for pH 7.0-7.2 (n=9) and 100% for pH below 7.0 (n=1). Fewer studies worked on the effects of seawater acidification on reproductive investment and success (10.9%). Yet, they showed that 50.0% of reproductive endpoints were affected for pH 7.8-8.0 (n=20), 75.0% for pH 7.6-7.8 (n=16), 100% for pH 7.2-7.4 (n=1) as well as for pH below 7 (n=1). Noticeable effects on reproduction included effects on sperm flagellar motility (Morita et al., 2010), time to gastrulation (Chua et al., 2013) and larval settlement (Albright et al., 2010; Doropoulos et al., 2012; Jiang et al., 2020; Pitts et al., 2020; Viyakarn et al., 2015). Finally, effects of seawater acidification on coral mortality were only showed under pH 7.6-7.8 (37.5% of affected endpoints, n=24). Mortality on adult colonies included partial mortality (local tissue loss; Bahr et al., 2016; Kavousi et al., 2016) and total mortality (death of the colonies; Yang et al., 2020). Studies also showed mortality of coral larvae (Cumbo et al., 2013; Doropoulos et al., 2012) and recruits (Dufault et al., 2012, 2013) under seawater acidification.

Respiration and photophysiology

Measurement of the coral host respiration rate is often used as an indicator of the metabolism's state/activity level. Moreover, the proper functioning of the algal symbiont photosynthetic activity is fundamental as it provides an energy source as photosynthates to the host (Muscatine et al., 1981). As the host, coral has some control over the photosynthetic activity of the algal symbionts (by contributing to the low pH of the symbiosome via H*-ATPase, Barott et al., 2015), maintaining this vital function is thus critical. The endpoints related to coral host respiration rate and algal symbiont photosynthetic activity were poorly affected by seawater acidification, regardless of the pH level (Fig. 3). Despite this, studies on the effects of seawater acidification on photosynthetic processes gave ambiguous results, with multiple studies showing an increased photosynthetic rate under low pH (Anderson et al., 2019; Biscéré et al., 2019; Langdon, 2005). Response heterogeneity could be linked to strain-specific responses by distinct *Symbiodinium sp.* (Brading et al., 2011; Graham & Sanders, 2016) or even attributed to differences in experimental parameters such as light or nutrients, which both contributes to photosynthesis (Chauvin et al., 2011). It is expected that the effect of seawater acidification on the coral's metabolism is small as compared to the effect of increasing temperature (Agostini et al., 2013; Cumbo et al., 2013; Edmunds & Burgess, 2016; Van der Zande et al., 2020).

Biomolecules

Quantification of biomolecules such as lipids, carbohydrates and proteins is often used as an indicator of energetic reserves of organisms. As such, their depletion under stressful conditions is often related to increased metabolic demands in ectotherms (Deutsch et al., 2008). Endpoints related to lipid, protein and carbohydrate metabolisms seemed to remain relatively stable under decreased pH (Fig. 3). Yet, for lipids, 55.6% of endpoints were affected for pH 7.6-7.8 (n=9) and 33.3% for pH 7.4-7.6 (n=6). Reduction of lipid content may show that maintenance of coral performances and calcification under seawater acidification increase energetic costs, which is met through catabolism of energy reserves. The subcategory "energy content" (here, the sum of lipids, carbohydrates and proteins) showed to be greatly affected at pH 7.6-7.8, but this requires further attention as it has only been studied in one experiment (Wall et al., 2017).

Cellular stress-response mechanism

Changes in the environment, like OA, are likely to drive cellular stress response, a universally conserved mechanism used to protect macromolecules within cells from the potential damage that OA may cause (Kaniewska et al., 2012). Here, a number of studies indicated that endpoints involved in cellular stress-response mechanisms were affected by seawater acidification (Fig. 3). In the subcategory "other molecular actors", 19.2% of endpoints were affected for pH 7.8-8.0 (n=26), 16.7% for pH 7.6-7.8 (n=6), 0% for pH 7.4-7.6 (n=2) and 100% for pH 7.2-7.4 (n=1). These were attributed to genes involved in innate immunity (Kaniewska et al., 2012) and heat shock response (Liew et al., 2018; Moya et al., 2015; Putnam et al., 2013). Thirty-one percent of endpoints associated with oxidative stress were affected at pH 7.8-8.0 (n=26) and 36.6% at pH 7.6-7.8 (n=41). These were essentially affecting the activity level of antioxidant enzymes (Bielmyer-Fraser et al., 2018; Jiang et al., 2019; Soriano-Santiago et al., 2013) or genes involved in the antioxidant response (Kaniewska et al., 2012; Moya et al., 2015). These later studies also indicated up/down regulation of anti-apoptotic genes, as showed with 50% of endpoints affected under pH 7.8-8.0 (n=2) and 100% under pH 7.6-7.8 (n=9).

Other biological category

Coral reefs form complex network of interactions between species. Each of its component plays a role and contributes to the overall equilibrium of the reef as well as its resilience to environmental change. Working on the coral community is therefore necessary as it allows considering the interactions between coral species and therefore provides more realistic responses. Yet, few studies

have experimentally worked on the effects of seawater acidification on coral community, likely owing to the difficulty of having a realistic community in an aquarium. For this aspect, we suggest to refer to the numerous *in situ* studies, which naturally consider the coral community. Endpoints indicated that community structure was changed under pH 7.6-8.0 (100% of affected endpoints, n=2; Edmunds et al., 2019, 2020).

As acidifying conditions affect corals, their energy requirements increase and must be compensated in order to sustain the proper biological functioning of the coral. This may be studied by looking at feeding rate (the input of energy *via* heterotrophy) and excretion rate (ammonium ion concentration; the output of energy through protein catabolism) of corals. Towle et al. (2015) indicated that coral can buffer OA effects by increasing feeding rates under pH 7.8-8.0 and Edmunds & Wall (2014) showed that catabolism of proteins and amino acids is accelerated by OA (pH 7.8-8.0), supporting that OA may perturb protein metabolism.

Phenotypic plasticity has the potential to facilitate rapid acclimatization of corals, a requirement in view of the rapid rate of change in the environment, driven by human activities. Mechanisms of rapid acclimatization may include changes in microbial community, with 50% of endpoints affected for pH 7.8-8.0 (n=4), 50% for pH 7.6-7.8 (n=4), 50% for pH 7.6-7.4 (n=2) and 87.5% for pH 7.2-7.4 (n=16) (Meron et al., 2011; Webster et al., 2013, 2016; Yang et al., 2020; Zhou et al., 2016). Another potential mechanism is epigenetics (*i.e.* the heritable postsynthesis modification of DNA or DNA-associated proteins, without a change in the DNA sequence itself, Feil & Fraga, 2012), with 50% of endpoints affected for pH 7.6-7.8 (n=2) and 100% for pH 7.2-7.4 (n=1) (Liew et al., 2018; Putnam et al., 2016). Finally, metabolomic profiling is a useful molecular tool that allows quantifying the intermediates and products of many biochemical processes. Putnam et al. (2016) found lower amount of metabolites under OA, suggesting a general suppression of metabolic activities. These techniques have started to be used on corals recently and deserve more attention in future investigations.

Gene expression

Understanding the fitness consequences of observed changes in endpoints is not always straightforward. This is particularly evident for changes in gene expression. An over or downregulations of a given gene does not implicitly translate into a positive or negative effect on protein level, function or fitness (Feder & Walser, 2005; Koussounadis et al., 2015; Greenbaum et al., 2003). Moreover, the intensity of an observed change at the gene expression level that is biologically significant is often poorly understood. For example, a 200% change in the expression of genes

involved in calcification does not necessarily translate into a 200% change in calcification. For these reasons, we did not use the gene expression data to infer biological impacts of seawater acidification.

3.3 Biological effects of pH 7.6-7.8 on tropical scleractinian corals

Many factors can modulate the biological response of tropical scleractinian corals to seawater acidification, including the tested species, life-history stage, tested pH or duration of exposure. As shown in the previous section, the effects on various endpoints tend to increase with decreasing pH. To better understand the role of the other modulating factors while minimizing the effect of the pH intensity, we decided to focus on pH class 2 (pH 7.6-7.8) for 2 reasons: (i) it is the class with the most available data (n endpoints=608); and, (ii) this range of pH corresponds to projected scenarios expected for 2100 (SSP3-7.0 and SSP5-8.5; IPCC, 2021).

General trends

Tested endpoints were only moderately affected under pH 7.6-7.8 (Fig. 4). No strong effects of the duration of exposure, coral genus or life stage could be identified, even when endpoints were separated into different categories.

Effects of the duration of exposure

The effect of the duration of the exposure in seawater acidification experiment is still poorly understood (Cumbo et al., 2013). Time-dependent response patterns were previously documented (Castillo et al., 2014; Schoepf et al., 2013). For example, Castillo et al. (2014) observed an increase in the calcification rate of *S. siderea* until the 60th day of exposure to acidified conditions, which thereafter started to decline. Similar results were reported for *A. millepora*, with a decrease of calcification that appeared only during the second half of the experiment (viz. after 1 month; Schoepf et al., 2013). In these specific cases, the parabolic responses to decreased pH would not have been detected if the duration of the experiments was shorter. These results might result from the ability of corals to compensate for the extra energy costs associated with the exposure to decreased pH, until reaching a time-dependent threshold beyond which the organism does not have the energy reserve allowing to counteract the extra costs associated with the stress. In other taxonomic groups, a different pattern was observed with a milder effect observed over longer period of time. For example, Dupont et al. (2013) showed that the strong negative effect of pH on adult sea urchin

fecundity observed after a 4-month exposure disappeared after 16 months. The negative effect observed over the first part of the experiment was interpreted as short-term costs associated with physiological acclimation to the new low pH environment. These examples highlight the value of long-term experiments.

Effects of the coral life stage

Early life stages are generally more vulnerable to environmental conditions than adults (Byrne, 2011), so dismissing the effects of decreasing pH on multiple life stages in current prediction models would represent an important bias for the projection of the fate of coral reef ecosystems. Indeed, if pH disrupts early-life processes, it would compromise sexual recruitment and therefore affect future population dynamics and the ability of coral reefs to recover from a disturbance (Caroselli et al., 2019). This question has received relatively poor attention and experiments that have considered more than one life stage are few (5%). Albright et al. (2010) suggested that OA has the potential to affect multiple, early-life history stages of *Acropora palmata*, with negative effects on gametes fertilization, larval settlement and post settlement growth. More experiment addressing this question and considering multiple life stages are thus crucial.

Effects of the coral genera

The biological responses of corals to decreased pH are highly variable and often considered to be species-specific (Bahr et al., 2016; Bedwell-Ivers et al., 2017; Bove et al., 2019; Brown & Edmunds, 2016; Edmunds et al., 2012; Hoadley et al., 2015; Sekizawa et al., 2017; Strahl et al., 2015). Several other parameters were proposed to explain the diversity of response observed between studies. While methodological differences could contribute to the difference between studies such as light intensities (Crawley et al., 2010; Dufault et al., 2013), feeding regime (Comeau et al., 2013; Edmunds, 2011; Houlbrèque et al., 2015) or water motion (Comeau et al., 2015), acclimation (*i.e.* their life history; Strahl et al., 2015), competition (Brien et al., 2016; Evensen et al., 2015; 2016; Horwitz et al., 2017), or even the size of the fragments used (Edmunds & Burgess, 2016), biological differences between taxa could explain the variability in coral sensitivity to seawater acidification. First, the "biomass plasticity" hypothesis (Edmunds, 2011) supports that corals reduce their sensitivity by increasing tissue thickness at low pH (Comeau et al., 2013; 2014; Edmunds, 2011). Indeed, thicker tissue allow isolating the site of calcification from the surrounding seawater, better protecting the zooxanthellae by changing the intensity and internal properties of the internal light field (Dimond et

al., 2012; Loya et al., 2001), as well as providing more energy reserves, which may be advantageous under low pH (as for Turbinaria reniformis, Hoadley et al., 2015). Massive Porites is known for having an unusually thick tissue layer that penetrates deeply into the skeleton (Edmunds, 2011). Second, the "pH regulation" hypothesis (Holcomb et al., 2015) suggests that scleractinians which have a high control over the regulation of their calcifying fluid's pH will be less affected by low pH conditions. Indeed, as the concentration of H⁺ in seawater increases under OA, the concentration gradient between seawater and the site of calcification increases accordingly (Cohen & McConnaughey, 2003; Jokiel, 2011; Ries, 2011; Von Euw et al., 2017). Transporting H⁺ will require more energy for the coral, which may alter calcification (McCulloch et al., 2012). As such, the coral's capacity/strategy to overcome higher energy requirements (through efficient zooxanthellae photosynthesis, McCulloch et al., 2012) or increased heterotrophic feeding (Towle et al., 2015) under more acidic conditions is also crucial in explaining interspecific differences in sensitivity. Despite this, increasing heterotrophic feeding under low pH conditions was not experimentally proven in all coral genera yet (such as Porites; Grottoli et al., 2006). This goes along with the hypothesis suggesting that differences in skeletal porosity may also explain differences in sensitivity of calcification among species. Indeed, scleractinians with highly porous skeletons require less CaCO₃ to calcify than corals that have low porosity skeletons, so these species may be less susceptible to seawater acidification (Brown & Edmunds, 2016; Jokiel, 2011). Finally, the "growth rate" hypothesis (Comeau et al., 2014) states that species characterized by higher calcification rate will be more vulnerable. Previous studies suggested that different calcification strategies exist across populations (Bove et al., 2019; Leong & Pawlik, 2010; Metcalfe & Monaghan, 2001). It relies on the theory that fast-growing coral species are more likely to be impaired under stressful conditions as their energy requirements for growth are higher than slow-growing species, which mainly store energy when the environment is favorable, allowing them to calcify normally under less-favorable conditions. However, this trend was not verified by Chan & Connolly (2013).

495

496

497

498

499

500

501

502

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

Evolutionary potential of corals under OA: local adaptation, carry over effects, parental effects

Overall, there is very little information on the potential of marine organisms to adapt to OA. To better understand the evolutionary responses of scleractinians to global changes, the best approach is to evaluate the effects of environmental drivers through multiple generations (Putnam et al., 2016). While trans-generational effects of OA has received more attention in the previous years (Allan et al., 2014; Donelson et al., 2012; Dupont et al., 2013; Hettinger et al., 2012; Kurihara et al., 2019; Miller et al., 2012; Munday, 2014; Parker et al., 2012), studies evaluating these effects on

scleractinian corals are few, likely due to their long life span. Among the available studies, evidences suggested that trans-generational acclimatization processes (e.g. epigenetics) have the potential to influence the sensitivity of corals to changing conditions (Putnam et al., 2016). This approach is of great importance to determine whether the exposure as an adult might enhance their descendant tolerance in order to better assess the status of future coral reef ecosystems. A high variability of response between individual corals and a high heritability for corals exposed to low pH conditions was observed, suggesting an adaptation potential (Jury et al., 2019). As highlighted by Putnam (2021), non-genetic, yet heritable phenotypic plasticity, will have significant ecological and evolutionary implications for sessile marine organisms persistence under rapid environmental changes. We support the need to consider the interactions of rapid and evolutionary responses (microbiome, epigenetic effects, parental carry-over effects; see Putnam, 2021), that have been overlooked so far, but deserve more attention as they provide additional avenues for rapid acclimatization.

The future of the reef: past environmental exposure, multiple stressors/drivers interactions

OA acts in conjunction with a myriad of other stressors/drivers, which presence and intensity vary locally. Among those, global ocean warming has received particular attention as it represents the main stressor on the reefs, inducing coral bleaching and widespread mortality in multiple regions. Studies considering a combination of increasing temperatures with seawater acidification often showed additive effects, thus worsening the effects observed when considering a single stressor (e.g. Albright & Mason, 2013; Anlauf et al., 2011; Anthony et al., 2008; Horvath et al., 2016; Kavousi et al., 2016; Reynaud et al., 2003). Then, to predict more realistically the future of coral reefs, their vulnerability must be assessed over long term and consider the stressors experienced by the reef organisms in the past. There is a temporality in the action of stressors on the reef that will condition the response of reef organisms. For example, multiple evidences showed that corals that have gone through repeated heat stress events in the past may have become more tolerant to future heat stress (Guest et al., 2012; Maynard et al., 2008; Thompson & van Woesik, 2009). Similarly, decreased thermal susceptibility was described with corals preconditioned to mild heat stress for a short period (Bay & Palumbi, 2015; Bellantuono et al., 2012a, 2012b; Middlebrook et al., 2008). In addition, OA will impair coral reefs previously altered by several bleaching events. Coral reef communities will be fragilized by the intensity and frequency of bleaching and it is likely that their capacity to endure additional stressor as OA will be highly detrimental.

4. Research gaps and perspectives

536

537538

539

540

Based on all aforementioned points, we selected a list of priorities and actions that should be addressed for future experimental works investigating the effects of seawater acidification on tropical scleractinians (Table 3).

541

542

543

Table 3. List of actions and priorities for future research on the effects of seawater acidification on tropical scleractinians.

Туре	Recommended action(s)
	,
Coral	Investigate corals from less-studied ecoregions, mesophotic depths
characteristics	
Experimental	Consider multigenerational approaches
design	Use multiple life-history stages, including early stages
	Realism: select the diversity and abundance of species based on the local
	assemblage, favour multi-species approach
	Include key species interacting with the tested species
	Avoid short-term exposure, consider multiple duration of exposure when
	feasible

Carbonate chemistry and scenarios

Monitor and consider the local variability of carbonate chemistry while defining scenarios and rely on local projections

Include the natural variability for the carbonate chemistry and other key environmental drivers (multiple stressors)

Apply best practices for carbonate chemistry parameters measurement (Dickson et al., 2007). Report pH in total scale.

Biological endpoints

Focus on endpoints to better understand the acclimatization capacity: epigenetic effects, microbiome and endolithic algae composition

Consider multiple complementary approaches: genetic and physiological tools

545	
546	
547	Funding
548	This work was supported by the "Fondation de France" for a project called "ACID REEFS", by the
549	"Ministère de la Transition Ecologique et Solidaire" and the Foundation for Research on Biodiversity
550	(FRB) for a project entitled "ACID REFES2".

Figure legends

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

551

Figure 1. World map of coral reefs that have been the subject of an experimental study (blue circles) on the effects of seawater acidification. Circle size: number of experimental study. Red colour: Global distribution of coral reefs (UNEP-WCMC, WorldFish Centre, WRI, TNC, 2010; version of March 2021). Figure 2. General trends in the experimental research of the effects of seawater acidification on biological functions of tropical scleractinian corals. (A) Number of experiments per country/location that experimentally examined the effects of seawater acidification on tropical scleractinian corals (total n=169). (B) Number of ex situ (total n= 169) and in situ (total n=27) studies published before 2021 that experimentally examined the effects of seawater acidification on tropical scleractinian corals. Relative proportion of (C) coral life stages (n=180) and (D) coral genera (n=240) studied in the 169 experiments that looked at the effects of seawater acidification on tropical scleractinian corals before 2021. Relative proportion of experimental design by (E) assemblages (n=169; i.e. number of studied species per experiment) and (F) duration of exposure (n=182) to seawater acidification in the same 169 experiments. (G) Relative proportion of the main biological categories studied in the same 169 experiments (n=255), with (H) specific focus on the fitness category. (I) Relative proportion of the pH scale used in the 169 experiments. Some of these bar charts have a (n) superior to 169 because some of the 169 experiments that were selected used more than one life stage, genera,

Figure 3. Effects of pH intensity on tropical scleractinian corals biological functions. Heatmaps indicate the number of endpoints (circle size) and the proportion of affected endpoints (color, in %) in each biological subcategory. Full red (100%) indicates that all endpoints are affected and full green (0%) that no endpoints were affected.

duration of exposure and/or biological endpoint.

Figure 4. Effects of pH 7.6-7.8 on tropical scleractinian corals biological functions according to the duration of exposure, the coral genus and life stage. Heatmaps indicate the number of endpoints

(circle size) and the proportion of affected endpoints (color, in %) in each biological subcategory. Full red (100%) indicates that all endpoints are affected and full green (0%) that no endpoints were affected.

References

581	Agostini, S., Fujimura, H., Higuchi, T., Yuyama, I., Casareto, B. E., Suzuki, Y., & Nakano, Y. (2013). The
582	effects of thermal and high-CO2 stresses on the metabolism and surrounding
583	microenvironment of the coral Galaxea fascicularis. Comptes Rendus Biologies, 336(8), 384-
584	391. https://doi.org/10.1016/j.crvi.2013.07.003
585	Albright, R., Caldeira, L., Hosfelt, J., Kwiatkowski, L., Maclaren, J. K., Mason, B. M., Nebuchina, Y.,
586	Ninokawa, A., Pongratz, J., Ricke, K. L., Rivlin, T., Schneider, K., Sesboüé, M., Shamberger, K.,
587	Silverman, J., Wolfe, K., Zhu, K., & Caldeira, K. (2016). Reversal of ocean acidification
588	enhances net coral reef calcification. <i>Nature</i> , <i>531</i> (7594), 362–365.
589	https://doi.org/10.1038/nature17155
590	Albright, R., & Mason, B. (2013). Projected near-future levels of temperature and pCO2 reduce coral
591	fertilization success. PLoS ONE, 8(2), e56468. https://doi.org/10.1371/journal.pone.0056468
592	Albright, R., Mason, B., Miller, M., & Langdon, C. (2010). Ocean acidification compromises
593	recruitment success of the threatened Caribbean coral Acropora palmata. Proceedings of the
594	National Academy of Sciences, 107(47), 20400–20404.
595	https://doi.org/10.1073/pnas.1007273107
596	Albright, R., Takeshita, Y., Koweek, D. A., Ninokawa, A., Wolfe, K., Rivlin, T., Nebuchina, Y., Young, J.,
597	& Caldeira, K. (2018). Carbon dioxide addition to coral reef waters suppresses net community
598	calcification. <i>Nature</i> , 555(7697), 516–519. https://doi.org/10.1038/nature25968
599	Allan, B. J. M., Miller, G. M., McCormick, M. I., Domenici, P., & Munday, P. L. (2014). Parental effects
600	improve escape performance of juvenile reef fish in a high-CO ₂ world. Proceedings of the
601	Royal Society B: Biological Sciences, 281(1777), 20132179.
602	https://doi.org/10.1098/rspb.2013.2179

603	Anderson, K. D., Cantin, N. E., Casey, J. M., & Pratchett, M. S. (2019). Independent effects of ocean
604	warming versus acidification on the growth, survivorship and physiology of two Acropora
605	corals. Coral Reefs, 38(6), 1225–1240. https://doi.org/10.1007/s00338-019-01864-y
606	Anlauf, H., D'Croz, L., & O'Dea, A. (2011). A corrosive concoction: The combined effects of ocean
607	warming and acidification on the early growth of a stony coral are multiplicative. Journal of
608	Experimental Marine Biology and Ecology, 397(1), 13–20.
609	https://doi.org/10.1016/j.jembe.2010.11.009
610	Anthony, K. R. N., Kline, D. I., Diaz-Pulido, G., Dove, S., & Hoegh-Guldberg, O. (2008). Ocean
611	acidification causes bleaching and productivity loss in coral reef builders. Proceedings of the
612	National Academy of Sciences, 105(45), 17442–17446.
613	https://doi.org/10.1073/pnas.0804478105
614	Asaad, I., Lundquist, C. J., Erdmann, M. V., Van Hooidonk, R., & Costello, M. J. (2018). Designating
615	spatial priorities for marine biodiversity conservation in the coral triangle. Frontiers in Marine
616	Science, 5. https://doi.org/10.3389/fmars.2018.00400
617	Bahr, K. D., Jokiel, P. L., & Rodgers, K. S. (2016). Relative sensitivity of five Hawaiian coral species to
618	high temperature under high-pCO2 conditions. Coral Reefs, 35(2), 729–738.
619	https://doi.org/10.1007/s00338-016-1405-4
620	Barkley, H. C., Cohen, A. L., Golbuu, Y., Starczak, V. R., DeCarlo, T. M., & Shamberger, K. E. F. (2015).
621	Changes in coral reef communities across a natural gradient in seawater pH. Science
622	Advances, 1(5), e1500328. https://doi.org/10.1126/sciadv.1500328
623	Barkley, H. C., Cohen, A. L., McCorkle, D. C., & Golbuu, Y. (2017). Mechanisms and thresholds for pH
624	tolerance in Palau corals. Journal of Experimental Marine Biology and Ecology, 489, 7–14.
625	https://doi.org/10.1016/j.jembe.2017.01.003
626	Barott, K. L., Venn, A. A., Perez, S. O., Tambutté, S., & Tresguerres, M. (2015). Coral host cells acidify
627	symbiotic algal microenvironment to promote photosynthesis. Proceedings of the National
628	Academy of Sciences, 112(2), 607-612. https://doi.org/10.1073/pnas.1413483112

629	Bay, R. A., & Palumbi, S. R. (2015). Rapid acclimation ability mediated by transcriptome changes in
630	reef-building corals. Genome Biology and Evolution, 7(6), 1602–1612.
631	https://doi.org/10.1093/gbe/evv085
632	Bedwell-Ivers, H. E., Koch, M. S., Peach, K. E., Joles, L., Dutra, E., & Manfrino, C. (2017). The role of in
633	hospite zooxanthellae photophysiology and reef chemistry on elevated pCO2 effects in two
634	branching Caribbean corals: Acropora cervicornis and Porites divaricata. ICES Journal of
635	Marine Science, 74(4), 1103–1112. https://doi.org/10.1093/icesjms/fsw026
636	Bellantuono, A. J., Granados-Cifuentes, C., Miller, D. J., Hoegh-Guldberg, O., & Rodriguez-Lanetty, M.
637	(2012). Coral thermal tolerance: tuning gene expression to resist thermal stress. PLoS ONE,
638	7(11), e50685. https://doi.org/10.1371/journal.pone.0050685
639	Bellantuono, A. J., Hoegh-Guldberg, O., & Rodriguez-Lanetty, M. (2012). Resistance to thermal stress
640	in corals without changes in symbiont composition. Proceedings of the Royal Society B:
641	Biological Sciences, 279(1731), 1100–1107. https://doi.org/10.1098/rspb.2011.1780
642	Bellworthy, J., Menoud, M., Krueger, T., Meibom, A., & Fine, M. (2018). Developmental carry over
643	effects of ocean warming and acidification in corals from a potential climate refugium, Gulf
644	of Aqaba. Journal of Experimental Biology. https://doi.org/10.1242/jeb.186940
645	Bielmyer-Fraser, G. K., Patel, P., Capo, T., & Grosell, M. (2018). Physiological responses of corals to
646	ocean acidification and copper exposure. Marine Pollution Bulletin, 133, 781–790.
647	https://doi.org/10.1016/j.marpolbul.2018.06.048
648	Biscéré, T., Zampighi, M., Lorrain, A., Jurriaans, S., Foggo, A., Houlbrèque, F., & Rodolfo-Metalpa, R.
649	(2019). High pCO_2 promotes coral primary production. Biology Letters, 15(7), 20180777.
650	https://doi.org/10.1098/rsbl.2018.0777
651	Bongaerts, P., Ridgway, T., Sampayo, E. M., & Hoegh-Guldberg, O. (2010). Assessing the 'deep reef
652	refugia' hypothesis: Focus on Caribbean reefs. Coral Reefs, 29(2), 309–327.
653	https://doi.org/10.1007/s00338-009-0581-x

- Bove, C. B., Ries, J. B., Davies, S. W., Westfield, I. T., Umbanhowar, J., & Castillo, K. D. (2019).
- 655 Common Caribbean corals exhibit highly variable responses to future acidification and
- warming. Proceedings of the Royal Society B: Biological Sciences, 286(1900), 20182840.
- 657 https://doi.org/10.1098/rspb.2018.2840
- Bove, C. B., Whitehead, R. F., & Szmant, A. M. (2020). Responses of coral gastrovascular cavity pH
- during light and dark incubations to reduced seawater pH suggest species-specific responses
- to the effects of ocean acidification on calcification. Coral Reefs, 39(6), 1675–1691.
- https://doi.org/10.1007/s00338-020-01995-7
- Brading, P., Warner, M. E., Davey, P., Smith, D. J., Achterberg, E. P., & Suggett, D. J. (2011).
- Differential effects of ocean acidification on growth and photosynthesis among phylotypes of
- 664 Symbiodinium (Dinophyceae). Limnology and Oceanography, 56(3), 927–938.
- https://doi.org/10.4319/lo.2011.56.3.0927
- Brien, H. V., Watson, S.-A., & Hoogenboom, M. O. (2016). Presence of competitors influences
- photosynthesis, but not growth, of the hard coral *Porites cylindrica* at elevated seawater CO₂.
- 668 ICES Journal of Marine Science: Journal Du Conseil, 73(3), 659–669.
- https://doi.org/10.1093/icesjms/fsv162
- Brown, D., & Edmunds, P. J. (2016). Differences in the responses of three scleractinians and the
- 671 hydrocoral Millepora platyphylla to ocean acidification. *Marine Biology*, 163(3).
- https://doi.org/10.1007/s00227-016-2837-7
- 673 Byrne, M. (2011). Impact of ocean warming and ocean acidification on marine invertebrate life
- history stages: Vulnerabilities and potential for persistence in a changing ocean.
- 675 Oceanography Marine Science, 49, 1–42.
- 676 Camp, E. F., Nitschke, M. R., Rodolfo-Metalpa, R., Houlbreque, F., Gardner, S. G., Smith, D. J.,
- Zampighi, M., & Suggett, D. J. (2017). Reef-building corals thrive within hot-acidified and
- deoxygenated waters. Scientific Reports, 7(1). https://doi.org/10.1038/s41598-017-02383-y

679	Caroselli, E., Gizzi, F., Prada, F., Marchini, C., Airi, V., Kaandorp, J., Falini, G., Dubinsky, Z., & Goffredo,
680	S. (2019). Low and variable pH decreases recruitment efficiency in populations of a
681	temperate coral naturally present at a CO ₂ vent. Limnology and Oceanography, 64(3), 1059–
682	1069. https://doi.org/10.1002/lno.11097
683	Castillo, K. D., Ries, J. B., Bruno, J. F., & Westfield, I. T. (2014). The reef-building coral Siderastrea
684	siderea exhibits parabolic responses to ocean acidification and warming. Proceedings of the
685	Royal Society B: Biological Sciences, 281(1797), 1–9. https://doi.org/10.1098/rspb.2014.1856
686	Chan, N. C. S., & Connolly, S. R. (2013). Sensitivity of coral calcification to ocean acidification: A meta-
687	analysis. Global Change Biology, 19(1), 282–290. https://doi.org/10.1111/gcb.12011
688	Chauvin, A., Denis, V., & Cuet, P. (2011). Is the response of coral calcification to seawater acidification
689	related to nutrient loading? Coral Reefs, 30(4), 911–923. https://doi.org/10.1007/s00338-
690	011-0786-7
691	Chua, C., Leggat, W., Moya, A., & Baird, A. (2013). Near-future reductions in pH will have no
692	consistent ecological effects on the early life-history stages of reef corals. Marine Ecology
693	Progress Series, 486, 143–151. https://doi.org/10.3354/meps10318
694	Cohen, A. L., & McConnaughey, T. A. (2003). Geochemical Perspectives on Coral Mineralization.
695	Reviews in mineralogy and geochemistry, 54(1), 151-187.
696	Comeau, S., Carpenter, R. C., & Edmunds, P. J. (2013). Effects of feeding and light intensity on the
697	response of the coral Porites rus to ocean acidification. Marine Biology, 160(5), 1127–1134.
698	https://doi.org/10.1007/s00227-012-2165-5
699	Comeau, S., Carpenter, R. C., & Edmunds, P. J. (2014). Effects of irradiance on the response of the
700	coral Acropora pulchra and the calcifying alga Hydrolithon reinboldii to temperature
701	elevation and ocean acidification. Journal of Experimental Marine Biology and Ecology, 453,
702	28–35. https://doi.org/10.1016/j.jembe.2013.12.013

- 703 Comeau, S., Edmunds, P. J., Lantz, C. A., & Carpenter, R. C. (2015). Water flow modulates the
- response of coral reef communities to ocean acidification. Scientific Reports, 4(1).
- 705 https://doi.org/10.1038/srep06681
- 706 Comeau, S., Edmunds, P. J., Spindel, N. B., & Carpenter, R. C. (2013). The responses of eight coral reef
- calcifiers to increasing partial pressure of CO₂ do not exhibit a tipping point. *Limnology and*
- 708 *Oceanography, 58*(1), 388–398. https://doi.org/10.4319/lo.2013.58.1.0388
- 709 Comeau, S., Edmunds, P. J., Spindel, N. B., & Carpenter, R. C. (2014). Fast coral reef calcifiers are
- 710 more sensitive to ocean acidification in short-term laboratory incubations. Limnology and
- 711 *Oceanography*, *59*(3), 1081–1091. https://doi.org/10.4319/lo.2014.59.3.1081
- 712 Crawley, A., Kline, D. I., Dunn, S., Anthony, K., & Dove, S. (2010). The effect of ocean acidification on
- symbiont photorespiration and productivity in Acropora formosa. Global Change Biology,
- 714 16(2), 851–863. https://doi.org/10.1111/j.1365-2486.2009.01943.x
- 715 Crook, E. D., Cohen, A. L., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A. (2013). Reduced
- calcification and lack of acclimatization by coral colonies growing in areas of persistent
- 717 natural acidification. Proceedings of the National Academy of Sciences, 110(27), 11044-
- 718 11049. https://doi.org/10.1073/pnas.1301589110
- 719 Crook, E. D., Potts, D., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A. (2012). Calcifying coral
- 720 abundance near low-pH springs: Implications for future ocean acidification. Coral Reefs,
- 721 31(1), 239–245. https://doi.org/10.1007/s00338-011-0839-y
- 722 Cumbo, V. R., Fan, T. Y., & Edmunds, P. J. (2013). Effects of exposure duration on the response of
- 723 Pocillopora damicornis larvae to elevated temperature and high pCO2. Journal of
- 724 Experimental Marine Biology and Ecology, 439, 100–107.
- 725 https://doi.org/10.1016/j.jembe.2012.10.019
- Cyronak, T., Takeshita, Y., Courtney, T. A., DeCarlo, E. H., Eyre, B. D., Kline, D. I., Martz, T., Page, H.,
- Price, N. N., Smith, J., Stoltenberg, L., Tresguerres, M., & Andersson, A. J. (2020). Diel

- temperature and pH variability scale with depth across diverse coral reef habitats. *Limnology*
- 729 *and Oceanography Letters*, 5(2), 193–203. https://doi.org/10.1002/lol2.10129
- 730 Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P.
- 731 R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings*
- 732 of the National Academy of Sciences, 105(18), 6668–6672.
- 733 https://doi.org/10.1073/pnas.0709472105
- 734 Dickson, A. G. (1993). The measurement of seawater pH. Marine Chemistry, 44(2-4), 131-142.
- 735 https://doi.org/10.1016/0304-4203(93)90198-W
- Dickson, A. G., Sabine, C. L., Christian, J. R., Bargeron, C. P., & North Pacific Marine Science
- 737 Organization (Eds.). (2007). Guide to best practices for ocean CO2 measurements. North
- 738 Pacific Marine Science Organization.
- 739 Dimond, J. L., Holzman, B. J., & Bingham, B. L. (2012). Thicker host tissues moderate light stress in a
- 740 cnidarian endosymbiont. Journal of Experimental Biology, 215(13), 2247–2254.
- 741 https://doi.org/10.1242/jeb.067991
- Dixon, G. B., Davies, S. W., Aglyamova, G. V., Meyer, E., Bay, L. K., & Matz, M. V. (2015). Genomic
- 743 determinants of coral heat tolerance across latitudes. *Science*, 348(6242), 1460–1462.
- 744 https://doi.org/10.1126/science.1261224
- Donelson, J. M., Munday, P. L., McCormick, M. I., & Pitcher, C. R. (2012). Rapid transgenerational
- acclimation of a tropical reef fish to climate change. *Nature Climate Change*, 2(1), 30–32.
- 747 https://doi.org/10.1038/nclimate1323
- Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). Ocean Acidification: The Other CO₂
- 749 Problem. Annual Review of Marine Science, 1(1), 169–192.
- 750 https://doi.org/10.1146/annurev.marine.010908.163834
- 751 Doo, S. S., Edmunds, P. J., & Carpenter, R. C. (2019). Ocean acidification effects on in situ coral reef
- 752 metabolism. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-48407-7

/53	Doropoulos, C., Ward, S., Diaz-Pulido, G., Hoegh-Guldberg, O., & Mumby, P. J. (2012). Ocean
754	acidification reduces coral recruitment by disrupting intimate larval-algal settlement
755	interactions: Elevated CO2 alters CCA-larval interactions. Ecology Letters, 15(4), 338–346.
756	https://doi.org/10.1111/j.1461-0248.2012.01743.x
757	Dufault, A. M., Cumbo, V. R., Fan, TY., & Edmunds, P. J. (2012). Effects of diurnally oscillating pCO 2
758	on the calcification and survival of coral recruits. Proceedings of the Royal Society B:
759	Biological Sciences, 279(1740), 2951–2958. https://doi.org/10.1098/rspb.2011.2545
760	Dufault, A. M., Ninokawa, A., Bramanti, L., Cumbo, V. R., Fan, TY., & Edmunds, P. J. (2013). The role
761	of light in mediating the effects of ocean acidification on coral calcification. Journal of
762	Experimental Biology, 216(9), 1570–1577. https://doi.org/10.1242/jeb.080549
763	Dupont, S., Dorey, N., Stumpp, M., Melzner, F., & Thorndyke, M. (2013). Long-term and trans-life-
764	cycle effects of exposure to ocean acidification in the green sea urchin Strongylocentrotus
765	droebachiensis. <i>Marine Biology</i> , 160(8), 1835–1843. https://doi.org/10.1007/s00227-012-
766	1921-x
767	Edmunds, P. J. (2011). Zooplanktivory ameliorates the effects of ocean acidification on the reef coral
768	Porites spp. <i>Limnology and Oceanography,</i> 56(6), 2402–2410.
769	https://doi.org/10.4319/lo.2011.56.6.2402
770	Edmunds, P. J., Brown, D., & Moriarty, V. (2012). Interactive effects of ocean acidification and
771	temperature on two scleractinian corals from Moorea, French Polynesia. Global Change
772	Biology, 18(7), 2173–2183. https://doi.org/10.1111/j.1365-2486.2012.02695.x
773	Edmunds, P. J., & Burgess, S. C. (2016). Size-dependent physiological responses of the branching coral
774	Pocillopora verrucosa to elevated temperature and pCO2. Journal of Experimental Biology.
775	https://doi.org/10.1242/jeb.146381
776	Edmunds, P. J., Doo, S. S., & Carpenter, R. C. (2019). Changes in coral reef community structure in
777	response to year-long incubations under contrasting pCO2 regimes. Marine Biology, 166(7).
778	https://doi.org/10.1007/s00227-019-3540-2

779	Edmunds, P. J., Doo, S. S., & Carpenter, R. C. (2020). Year-long effects of high pCO2 on the community
780	structure of a tropical fore reef assembled in outdoor flumes. ICES Journal of Marine Science,
781	77(3), 1055–1065. https://doi.org/10.1093/icesjms/fsaa015
782	Edmunds, P. J., & Wall, C. B. (2014). Evidence that high pCO ₂ affects protein metabolism in tropical
783	reef corals. The Biological Bulletin, 227(1), 68–77. https://doi.org/10.1086/BBLv227n1p68
784	Erez, J., Reynaud, S., Silverman, J., Schneider, K., & Allemand, D. (2011). Coral calcification under
785	ocean acidification and global change. In Z. Dubinsky & N. Stambler (Eds.), Coral Reefs: An
786	Ecosystem in Transition (pp. 151–176). Springer Netherlands. https://doi.org/10.1007/978-
787	94-007-0114-4_10
788	Evensen, N., Edmunds, P., & Sakai, K. (2015). Effects of pCO2 on spatial competition between the
789	corals Montipora aequituberculata and Porites lutea. Marine Ecology Progress Series, 541,
790	123–134. https://doi.org/10.3354/meps11512
791	Evensen, N. R., & Edmunds, P. J. (2016). Interactive effects of ocean acidification and neighbouring
792	corals on the growth of Pocillopora verrucosa. Marine Biology, 163(7).
793	https://doi.org/10.1007/s00227-016-2921-z
794	Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., Okazaki, R.,
795	Muehllehner, N., Glas, M. S., & Lough, J. M. (2011). Losers and winners in coral reefs
796	acclimatized to elevated carbon dioxide concentrations. Nature Climate Change, 1(3), 165–
797	169. https://doi.org/10.1038/nclimate1122
798	Fantazzini, P., Mengoli, S., Pasquini, L., Bortolotti, V., Brizi, L., Mariani, M., Di Giosia, M., Fermani, S.,
799	Capaccioni, B., Caroselli, E., Prada, F., Zaccanti, F., Levy, O., Dubinsky, Z., Kaandorp, J. A.,
800	Konglerd, P., Hammel, J. U., Dauphin, Y., Cuif, JP., Goffredo, S. (2015). Gains and losses of
801	coral skeletal porosity changes with ocean acidification acclimation. Nature Communications,
802	6(1). https://doi.org/10.1038/ncomms8785

- Feder, M. E., & Walser, J.-C. (2005). The biological limitations of transcriptomics in elucidating stress
- and stress responses. *Journal of Evolutionary Biology*, 18(4), 901–910.
- 805 https://doi.org/10.1111/j.1420-9101.2005.00921.x
- 806 Feely, R. A. (2004). Impact of Anthropogenic CO2 on the CaCO3 System in the Oceans. Science,
- 807 305(5682), 362–366. https://doi.org/10.1126/science.1097329
- 808 Feil, R., & Fraga, M. F. (2012). Epigenetics and the environment: Emerging patterns and implications.
- 809 Nature Reviews Genetics, 13(2), 97–109. https://doi.org/10.1038/nrg3142
- 810 Gattuso, J.-P., & Hansson, L. (Eds.). (2011). *Ocean acidification*. Oxford University Press, Oxford.
- Gattuso, J. P., Epitalon, J. M., Lavigne, H., Orr, J., Gentili, B., Hofmann, A., ... & Rae, J. (2020). seacarb:
- Seawater Carbonate Chemistry, version 3.2. 14.
- Georgiou, L., Falter, J., Trotter, J., Kline, D. I., Holcomb, M., Dove, S. G., Hoegh-Guldberg, O., &
- 814 McCulloch, M. (2015). Ph homeostasis during coral calcification in a free ocean CO ₂
- enrichment (FOCE) experiment, Heron Island reef flat, Great Barrier Reef. *Proceedings of the*
- 816 *National Academy of Sciences,* 112(43), 13219–13224.
- 817 https://doi.org/10.1073/pnas.1505586112
- 818 Graham, E. R., & Sanders, R. W. (2016). Species-specific photosynthetic responses of symbiotic
- zoanthids to thermal stress and ocean acidification. *Marine Ecology*, 37(2), 442–458.
- 820 https://doi.org/10.1111/maec.12291
- 821 Greenbaum, D., Colangelo, C., Williams, K., & Gerstein, M. (2003). Comparing protein abundance and
- mRNA expression levels on a genomic scale. Genome Biology, 4(9), 117.
- 823 https://doi.org/10.1186/gb-2003-4-9-117
- 824 Grottoli, A. G., Rodrigues, L. J., & Palardy, J. E. (2006). Heterotrophic plasticity and resilience in
- bleached corals. *Nature*, 440(7088), 1186–1189. https://doi.org/10.1038/nature04565
- 826 Guest, J. R., Baird, A. H., Maynard, J. A., Muttaqin, E., Edwards, A. J., Campbell, S. J., Yewdall, K.,
- Affendi, Y. A., & Chou, L. M. (2012). Contrasting patterns of coral bleaching susceptibility in

828	2010 suggest an adaptive response to thermal stress. PLoS ONE, 7(3), e33353.
829	https://doi.org/10.1371/journal.pone.0033353
830	Hendriks, I. E., Duarte, C. M., & Álvarez, M. (2010). Vulnerability of marine biodiversity to ocean
831	acidification: A meta-analysis. Estuarine, Coastal and Shelf Science, 86(2), 157–164.
832	https://doi.org/10.1016/j.ecss.2009.11.022
833	Hettinger, A., Sanford, E., Hill, T. M., Russell, A. D., Sato, K. N. S., Hoey, J., Forsch, M., Page, H. N., &
834	Gaylord, B. (2012). Persistent carry-over effects of planktonic exposure to ocean acidification
835	in the Olympia oyster. <i>Ecology</i> , <i>93</i> (12), 2758–2768. https://doi.org/10.1890/12-0567.1
836	Hoadley, K. D., Pettay, D. T., Grottoli, A. G., Cai, WJ., Melman, T. F., Schoepf, V., Hu, X., Li, Q., Xu, H.,
837	Wang, Y., Matsui, Y., Baumann, J. H., & Warner, M. E. (2015). Physiological response to
838	elevated temperature and pCO2 varies across four Pacific coral species: Understanding the
839	unique host+symbiont response. Scientific Reports, 5(1). https://doi.org/10.1038/srep18371
840	Holcomb, M., Venn, A. A., Tambutté, E., Tambutté, S., Allemand, D., Trotter, J., & McCulloch, M.
841	(2015). Coral calcifying fluid pH dictates response to ocean acidification. Scientific Reports,
842	4(1). https://doi.org/10.1038/srep05207
843	Horvath, K. M., Castillo, K. D., Armstrong, P., Westfield, I. T., Courtney, T., & Ries, J. B. (2016). Next-
844	century ocean acidification and warming both reduce calcification rate, but only acidification
845	alters skeletal morphology of reef-building coral Siderastrea siderea. Scientific Reports, 6(1).
846	https://doi.org/10.1038/srep29613
847	Horwitz, R., Hoogenboom, M. O., & Fine, M. (2017). Spatial competition dynamics between reef
848	corals under ocean acidification. Scientific Reports, 7(1). https://doi.org/10.1038/srep40288
849	Houlbrèque, F., Reynaud, S., Godinot, C., Oberhänsli, F., Rodolfo-Metalpa, R., & Ferrier-Pagès, C.
850	(2015). Ocean acidification reduces feeding rates in the scleractinian coral Stylophora
851	pistillata: Acidification and Stylophora nutrition. Limnology and Oceanography, 60(1), 89–99.
852	https://doi.org/10.1002/lno.10003

853	IPCC 2021. Fox-Kemper, B., H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S. S. Drijfhout, T. L. Edwards, N. R.
854	Golledge, M. Hemer, R. E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I. S. Nurhati, L. Ruiz, J-
855	B. Sallée, A. B. A. Slangen, Y. Yu, 2021, Ocean, Cryosphere and Sea Level Change. In: Climate
856	Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth
857	Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V.,
858	P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis,
859	M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu
860	and B. Zhou (eds.)]. Cambridge University Press. In Press.
861	Jacob, H., Besson, M., Swarzenski, P. W., Lecchini, D., & Metian, M. (2020). Effects of Virgin Micro-
862	and Nanoplastics on Fish: Trends, Meta-Analysis, and Perspectives. Environmental Science &
863	Technology, 54(8), 4733–4745. https://doi.org/10.1021/acs.est.9b05995
864	Jagers, S. C., Matti, S., Crépin, AS., Langlet, D., Havenhand, J. N., Troell, M., Filipsson, H. L., Galaz, V.
865	R., & Anderson, L. G. (2019). Societal causes of, and responses to, ocean acidification. Ambio,
866	48(8), 816–830. https://doi.org/10.1007/s13280-018-1103-2
867	Jiang, L., Guo, ML., Zhang, F., Zhang, YY., Zhou, GW., Lei, XM., Yuan, XC., Sun, YF., Yuan, T.,
868	Cai, L., Lian, JS., Liu, S., Qian, PY., & Huang, H. (2020). Impacts of elevated temperature
869	and pCO2 on the brooded larvae of Pocillopora damicornis from Luhuitou Reef, China:
870	Evidence for local acclimatization. Coral Reefs, 39(2), 331–344.
871	https://doi.org/10.1007/s00338-020-01894-x
872	Jiang, LQ., Carter, B. R., Feely, R. A., Lauvset, S. K., & Olsen, A. (2019). Surface ocean pH and buffer
873	capacity: Past, present and future. Scientific Reports, 9(1). https://doi.org/10.1038/s41598-
874	019-55039-4
875	Jokiel, P. L. (2011). The reef coral two compartment proton flux model: A new approach relating
876	tissue-level physiological processes to gross corallum morphology. Journal of Experimental
877	Marine Biology and Ecology, 409(1–2), 1–12. https://doi.org/10.1016/j.jembe.2011.10.008

878	Jury, C. P., Delano, M. N., & Toonen, R. J. (2019). High heritability of coral calcification rates and
879	evolutionary potential under ocean acidification. Scientific Reports, 9(1).
880	https://doi.org/10.1038/s41598-019-56313-1
881	Kaniewska, P., Campbell, P. R., Kline, D. I., Rodriguez-Lanetty, M., Miller, D. J., Dove, S., & Hoegh-
882	Guldberg, O. (2012). Major cellular and physiological impacts of ocean acidification on a reef
883	building coral. PLoS ONE, 7(4), e34659. https://doi.org/10.1371/journal.pone.0034659
884	Kavousi, J., Parkinson, J. E., & Nakamura, T. (2016). Combined ocean acidification and low
885	temperature stressors cause coral mortality. Coral Reefs, 35(3), 903–907.
886	https://doi.org/10.1007/s00338-016-1459-3
887	Kawahata, H., Fujita, K., Iguchi, A., Inoue, M., Iwasaki, S., Kuroyanagi, A., Maeda, A., Manaka, T.,
888	Moriya, K., Takagi, H., Toyofuku, T., Yoshimura, T., & Suzuki, A. (2019). Perspective on the
889	response of marine calcifiers to global warming and ocean acidification—Behavior of corals
890	and foraminifera in a high CO2 world "hot house". Progress in Earth and Planetary Science,
891	6(1). https://doi.org/10.1186/s40645-018-0239-9
892	Kline, D. I., Teneva, L., Okamoto, D. K., Schneider, K., Caldeira, K., Miard, T., Chai, A., Marker, M.,
893	Dunbar, R. B., Mitchell, B. G., Dove, S., & Hoegh-Guldberg, O. (2019). Living coral tissue slows
894	skeletal dissolution related to ocean acidification. Nature Ecology & Evolution, 3(10), 1438-
895	1444. https://doi.org/10.1038/s41559-019-0988-x
896	Koussounadis, A., Langdon, S. P., Um, I. H., Harrison, D. J., & Smith, V. A. (2015). Relationship
897	between differentially expressed mRNA and mRNA-protein correlations in a xenograft model
898	system. Scientific Reports, 5(1). https://doi.org/10.1038/srep10775
899	Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., & Gattuso,
900	J. (2013). Impacts of ocean acidification on marine organisms: Quantifying sensitivities and
901	interaction with warming. Global Change Biology, 19(6), 1884–1896.
902	https://doi.org/10.1111/gcb.12179

903	Kroeker, K. J., Kordas, R. L., Crim, R. N., & Singh, G. G. (2010). Meta-analysis reveals negative yet
904	variable effects of ocean acidification on marine organisms: Biological responses to ocean
905	acidification. Ecology Letters, 13(11), 1419–1434. https://doi.org/10.1111/j.1461-
906	0248.2010.01518.x
907	Kurihara, H., Wouters, J., & Yasuda, N. (2019). Seasonal calcification of the coral Acropora digitifera
908	from a subtropical marginal Okinawa reef under ocean acidification. Coral Reefs, 38(3), 443-
909	454. https://doi.org/10.1007/s00338-019-01794-9
910	Langdon, C. (2005). Effect of elevated pCO ₂ on photosynthesis and calcification of corals and
911	interactions with seasonal change in temperature/irradiance and nutrient enrichment.
912	Journal of Geophysical Research, 110(C9). https://doi.org/10.1029/2004JC002576
913	Leong, W., & Pawlik, J. (2010). Evidence of a resource trade-off between growth and chemical
914	defenses among Caribbean coral reef sponges. Marine Ecology Progress Series, 406, 71–78.
915	https://doi.org/10.3354/meps08541
916	Liew, Y. J., Zoccola, D., Li, Y., Tambutté, E., Venn, A. A., Michell, C. T., Cui, G., Deutekom, E. S.,
917	Kaandorp, J. A., Voolstra, C. R., Forêt, S., Allemand, D., Tambutté, S., & Aranda, M. (2018).
918	Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-building
919	coral. Science Advances, 10.
920	Loya, Y., Sakai, K., Yamazato, K., Nakano, Y., Sambali, H., & van Woesik, R. (2001). Coral bleaching:
921	The winners and the losers. Ecology Letters, 4(2), 122–131. https://doi.org/10.1046/j.1461-
922	0248.2001.00203.x
923	Marcelino, V. R., Morrow, K. M., Oppen, M. J. H., Bourne, D. G., & Verbruggen, H. (2017). Diversity
924	and stability of coral endolithic microbial communities at a naturally high pCO_2 reef.
925	Molecular Ecology, 26(19), 5344–5357. https://doi.org/10.1111/mec.14268
926	Martinez, A., Crook, E. D., Barshis, D. J., Potts, D. C., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A.
927	(2019). Species-specific calcification response of Caribbean corals after 2-year

928	transplantation to a low aragonite saturation submarine spring. Proceedings of the Royal
929	Society B: Biological Sciences, 286(1905), 20190572. https://doi.org/10.1098/rspb.2019.0572
930	Maynard, J. A., Anthony, K. R. N., Marshall, P. A., & Masiri, I. (2008). Major bleaching events can lead
931	to increased thermal tolerance in corals. <i>Marine Biology</i> , 155(2), 173–182.
932	https://doi.org/10.1007/s00227-008-1015-y
933	McCulloch, M., Falter, J., Trotter, J., & Montagna, P. (2012). Coral resilience to ocean acidification and
934	global warming through pH up-regulation. Nature Climate Change, 2(8), 623–627.
935	https://doi.org/10.1038/nclimate1473
936	McLachlan, R. H., Price, J. T., Solomon, S. L., & Grottoli, A. G. (2020). Thirty years of coral heat-stress
937	experiments: A review of methods. <i>Coral Reefs</i> , <i>39</i> (4), 885–902.
938	https://doi.org/10.1007/s00338-020-01931-9
939	Meron, D., Atias, E., Iasur Kruh, L., Elifantz, H., Minz, D., Fine, M., & Banin, E. (2011). The impact of
940	reduced pH on the microbial community of the coral Acropora eurystoma. The ISME Journal,
941	5(1), 51–60. https://doi.org/10.1038/ismej.2010.102
942	Metcalfe, N. B., & Monaghan, P. (2001). Compensation for a bad start: Grow now, pay later? <i>Trends</i>
943	in Ecology & Evolution, 16(5), 254–260. https://doi.org/10.1016/S0169-5347(01)02124-3
944	Middlebrook, R., Hoegh-Guldberg, O., & Leggat, W. (2008). The effect of thermal history on the
945	susceptibility of reef-building corals to thermal stress. Journal of Experimental Biology,
946	211(7), 1050–1056. https://doi.org/10.1242/jeb.013284
947	Miller, G. M., Watson, SA., Donelson, J. M., McCormick, M. I., & Munday, P. L. (2012). Parental
948	environment mediates impacts of increased carbon dioxide on a coral reef fish. Nature
949	Climate Change, 2(12), 858–861. https://doi.org/10.1038/nclimate1599
950	Morais, J., & Santos, B. A. (2018). Limited potential of deep reefs to serve as refuges for tropical
951	Southwestern Atlantic corals. Ecosphere, 9(7), e02281. https://doi.org/10.1002/ecs2.2281

Morita, M., Suwa, R., Iguchi, A., Nakamura, M., Shimada, K., Sakai, K., & Suzuki, A. (2010). Ocean 952 acidification reduces sperm flagellar motility in broadcast spawning reef invertebrates. 953 954 *Zygote*, 18(2), 103–107. https://doi.org/10.1017/S0967199409990177 Morrow, K. M., Bourne, D. G., Humphrey, C., Botté, E. S., Laffy, P., Zaneveld, J., Uthicke, S., Fabricius, 955 956 K. E., & Webster, N. S. (2015). Natural volcanic CO2 seeps reveal future trajectories for hostmicrobial associations in corals and sponges. The ISME Journal, 9(4), 894-908. 957 958 https://doi.org/10.1038/ismej.2014.188 959 Moya, A., Huisman, L., Forêt, S., Gattuso, J.-P., Hayward, D. C., Ball, E. E., & Miller, D. J. (2015). Rapid 960 acclimation of juvenile corals to CO₂-mediated acidification by upregulation of heat shock 961 protein and Bcl-2 genes. Molecular Ecology, *24*(2), 438–452. 962 https://doi.org/10.1111/mec.13021 Munday, P. L. (2014). Transgenerational acclimation of fishes to climate change and ocean 963 acidification. F1000Prime Reports, 6. https://doi.org/10.12703/P6-99 964 965 Muscatine, L., R. McCloskey, L., & E. Marian, R. (1981). Estimating the daily contribution of carbon 966 from zooxanthellae to coral animal respiration: Muscatine et al. Limnology and 967 Oceanography, 26(4), 601-611. https://doi.org/10.4319/lo.1981.26.4.0601 968 Noonan, S. H. C., & Fabricius, K. E. (2016). Ocean acidification affects productivity but not the severity 969 of thermal bleaching in some tropical corals. ICES Journal of Marine Science: Journal Du Conseil, 73(3), 715–726. https://doi.org/10.1093/icesjms/fsv127 970 Noonan, S. H. C., Kluibenschedl, A., & Fabricius, K. E. (2018). Ocean acidification alters early 971 972 successional coral reef communities and their rates of community metabolism. PLOS ONE, 973 13(5), e0197130. https://doi.org/10.1371/journal.pone.0197130 O'Brien, P. A., Smith, H. A., Fallon, S., Fabricius, K., Willis, B. L., Morrow, K. M., & Bourne, D. G. (2018). 974 975 Elevated CO2 Has Little Influence on the Bacterial Communities Associated With the pH-Porites 976 Tolerant Microbiology, 9. Coral, Massive spp. **Frontiers** in https://doi.org/10.3389/fmicb.2018.02621 977

978 Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. M., Gnanadesikan, A., Gruber, N., 979 Ishida, A., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., 980 Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., ... Yool, A. (2005). Anthropogenic 981 decline in high-latitude ocean carbonate by 2100. Nature, 437(7059), 681-686. 982 Pandolfi, J. M. (2003). Global trajectories of the long-term decline of coral reef ecosystems. Science, 983 301(5635), 955–958. https://doi.org/10.1126/science.1085706 984 Parker, L. M., Ross, P. M., O'Connor, W. A., Borysko, L., Raftos, D. A., & Pörtner, H.-O. (2012). Adult 985 exposure influences offspring response to ocean acidification in oysters. Global Change 986 Biology, 18(1), 82-92. https://doi.org/10.1111/j.1365-2486.2011.02520.x 987 Pierrot, Lewis, D. E., & Wallace, D. W. R. (2006). MS Excel Program Developed for CO2 System Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge 988 989 National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. https://doi.org/10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a 990 991 Pitts, K. A., Campbell, J. E., Figueiredo, J., & Fogarty, N. D. (2020). Ocean acidification partially 992 mitigates the negative effects of warming on the recruitment of the coral, Orbicella 993 faveolata. Coral Reefs, 39(2), 281-292. https://doi.org/10.1007/s00338-019-01888-4 994 Prada, F., Brizi, L., Franzellitti, S., Mengoli, S., Fermani, S., Polishchuk, I., Baraldi, N., Ricci, F., Palazzo, 995 Q., Caroselli, E., Pokroy, B., Giorgini, L., Dubinsky, Z., Fantazzini, P., Falini, G., Goffredo, S., & Fabricius, K. E. (2021). Coral micro- and macro-morphological skeletal properties in response 996 to life-long acclimatization at CO2 vents in Papua New Guinea. Scientific Reports, 11(1). 997 998 https://doi.org/10.1038/s41598-021-98976-9 999 Putnam, H. M. (2021). Avenues of reef-building coral acclimatization in response to rapid 1000 Experimental environmental change. Journal Biology, 224, jeb.239319. of https://doi.org/10.1242/jeb.239319 1001

1002	Putnam, H. M., Davidson, J. M., & Gates, R. D. (2016). Ocean acidification influences host DNA
1003	methylation and phenotypic plasticity in environmentally susceptible corals. Evolutionary
1004	Applications, 9(9), 1165–1178. https://doi.org/10.1111/eva.12408
1005	Putnam, H. M., & Gates, R. D. (2015). Preconditioning in the reef-building coral Pocillopora
1006	damicornis and the potential for trans-generational acclimatization in coral larvae under
1007	future climate change conditions. Journal of Experimental Biology, 218(15), 2365–2372.
1008	https://doi.org/10.1242/jeb.123018
1009	Putnam, H. M., Mayfield, A. B., Fan, T. Y., Chen, C. S., & Gates, R. D. (2013). The physiological and
1010	molecular responses of larvae from the reef-building coral Pocillopora damicornis exposed to
1011	near-future increases in temperature and pCO2. Marine Biology, 160(8), 2157–2173.
1012	https://doi.org/10.1007/s00227-012-2129-9
1013	Raven. (2005). Ocean acidification due to increasing atmospheric carbon dioxide. Policy Document
1014	12/05 (The Royal Society, London).
1015	Reynaud, S., Leclercq, N., Romaine-Lioud, S., Ferrier-Pages, C., Jaubert, J., & Gattuso, JP. (2003).
1016	Interacting effects of CO2 partial pressure and temperature on photosynthesis and
1017	calcification in a scleractinian coral. Global Change Biology, 9(11), 1660–1668.
1018	https://doi.org/10.1046/j.1365-2486.2003.00678.x
1019	Riebesell, U. (2008). Acid test for marine biodiversity. <i>Nature</i> , 454(7200), 46–47.
1020	https://doi.org/10.1038/454046a
1021	Riebesell, U., Fabry, V. J., Hansson, L., & Gattuso, J. P. (2011). Guide to best practices for ocean
1022	acidification research and data reporting. Office for Official Publications of the European
1023	Communities.
1024	Ries, J. B. (2011). A physicochemical framework for interpreting the biological calcification response
1025	to CO2-induced ocean acidification. <i>Geochimica et Cosmochimica Acta, 75</i> (14), 4053–4064.
1026	https://doi.org/10.1016/j.gca.2011.04.025

1027 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F. S., Lambin, E. F., Lenton, T. M., Scheffer, 1028 M., Folke, C., Schellnhuber, H. J., Nykvist, B., de Wit, C. A., Hughes, T., van der Leeuw, S., 1029 Rodhe, H., Sörlin, S., Snyder, P. K., Costanza, R., Svedin, U., ... Foley, J. A. (2009). A safe 1030 operating for humanity. Nature, 461(7263), 472-475. space 1031 https://doi.org/10.1038/461472a 1032 Sayers, A. (2007). Tips and tricks in performing a systematic review. British Journal of General 1033 Practice, 57(538), 425-425. 1034 Schoepf, V., Grottoli, A. G., Warner, M. E., Cai, W. J., Melman, T. F., Hoadley, K. D., Pettay, D. T., Hu, 1035 X., Li, Q., Xu, H., Wang, Y., Matsui, Y., & Baumann, J. H. (2013). Coral energy reserves and 1036 calcification in a high-CO2 world at two temperatures. In S. C. A. Ferse (Ed.), PLoS ONE (Vol. 1037 8, Issue 10, p. e75049). https://doi.org/10.1371/journal.pone.0075049 1038 Sekizawa, A., Uechi, H., Iguchi, A., Nakamura, T., Kumagai, N. H., Suzuki, A., Sakai, K., & Nojiri, Y. 1039 (2017). Intraspecific variations in responses to ocean acidification in two branching coral 1040 species. Marine **Pollution** Bulletin, 122(1-2),282-287. 1041 https://doi.org/10.1016/j.marpolbul.2017.06.061 1042 Smith, J. N., Strahl, J., Noonan, S. H. C., Schmidt, G. M., Richter, C., & Fabricius, K. E. (2016). Reduced 1043 heterotrophy in the stony coral Galaxea fascicularis after life-long exposure to elevated 1044 carbon dioxide. Scientific Reports, 6(1). https://doi.org/10.1038/srep27019 1045 Soriano-Santiago, O. S., Liñán-Cabello, M. A., Delgadillo-Nuño, M. A., Ortega-Ortiz, C., & Cuevas-1046 Venegas, S. (2013). Physiological responses to oxidative stress associated with pH variations in host tissue and zooxanthellae of hermatypic coral Pocillopora capitata. Marine and 1047 1048 Freshwater Behaviour Physiology, and 46(5), 275-286. 1049 https://doi.org/10.1080/10236244.2013.827877 1050 Srednick, G., Bergman, J. L., Doo, S. S., Hawthorn, M., Ferree, J., Rojas, R., Arias, R., Edmunds, P. J., & 1051 Carpenter, R. C. (2020). Shallow coral reef free ocean carbon enrichment: Novel in situ

1052	flumes to manipulate pCO ₂ on shallow tropical coral reef communities. Limnology and
1053	Oceanography: Methods, 18(3), 116–128. https://doi.org/10.1002/lom3.10349
1054	Strahl, J., Stolz, I., Uthicke, S., Vogel, N., Noonan, S. H. C., & Fabricius, K. E. (2015). Physiological and
1055	ecological performance differs in four coral taxa at a volcanic carbon dioxide seep.
1056	Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 184,
1057	179–186. https://doi.org/10.1016/j.cbpa.2015.02.018
1058	Tambutté, E., Venn, A. A., Holcomb, M., Segonds, N., Techer, N., Zoccola, D., Allemand, D., &
1059	Tambutté, S. (2015). Morphological plasticity of the coral skeleton under CO2-driven
1060	seawater acidification. Nature Communications, 6(1). https://doi.org/10.1038/ncomms8368
1061	Thompson, D. M., & van Woesik, R. (2009). Corals escape bleaching in regions that recently and
1062	historically experienced frequent thermal stress. Proceedings of the Royal Society B:
1063	Biological Sciences, 276(1669), 2893–2901. https://doi.org/10.1098/rspb.2009.0591
1064	Towle, E. K., Enochs, I. C., & Langdon, C. (2015). Threatened caribbean coral is able to mitigate the
1065	adverse effects of ocean acidification on calcification by increasing feeding rate. PLOS ONE,
1066	10(4), e0123394. https://doi.org/10.1371/journal.pone.0123394
1067	UNEP-WCMC, WorldFish Centre, WRI, TNC (2010). Global distribution of warm-water coral reefs,
1068	compiled from multiple sources including the Millennium Coral Reef Mapping Project.
1069	Version 4.0. Includes contributions from IMaRS-USF and IRD (2005), IMaRS-USF (2005) and
1070	Spalding et al. (2001). Cambridge (UK): UNEP World Conservation Monitoring Centre. URL:
1071	http://data.unep-wcmc.org/datasets/1
1072	Vargas, C. A., Lagos, N. A., Lardies, M. A., Duarte, C., Manríquez, P. H., Aguilera, V. M., Broitman, B.,
1073	Widdicombe, S., & Dupont, S. (2017). Species-specific responses to ocean acidification should
1074	account for local adaptation and adaptive plasticity. <i>Nature Ecology & Evolution</i> , 1(4).
1075	https://doi.org/10.1038/s41559-017-0084

10/6	Venn, A. A., Bernardet, C., Chabenat, A., Tambutte, E., & Tambutte, S. (2020). Paracellular transport
1077	to the coral calcifying medium: Effects of environmental parameters. The Journal of
1078	Experimental Biology, 223(17), jeb227074. https://doi.org/10.1242/jeb.227074
1079	Venn, A. A., Tambutté, E., Caminiti-Segonds, N., Techer, N., Allemand, D., & Tambutté, S. (2019).
1080	Effects of light and darkness on pH regulation in three coral species exposed to seawater
1081	acidification. Scientific Reports, 9(1). https://doi.org/10.1038/s41598-018-38168-0
1082	Venn, A. A., Tambutte, E., Holcomb, M., Laurent, J., Allemand, D., & Tambutte, S. (2013). Impact of
1083	seawater acidification on pH at the tissue-skeleton interface and calcification in reef corals.
1084	Proceedings of the National Academy of Sciences, 110(5), 1634–1639.
1085	https://doi.org/10.1073/pnas.1216153110
1086	Veron, J., Stafford-Smith, M., DeVantier, L., & Turak, E. (2015). Overview of distribution patterns of
1087	zooxanthellate Scleractinia. Frontiers in Marine Science, 1.
1088	https://doi.org/10.3389/fmars.2014.00081
1089	Viyakarn, V., Lalitpattarakit, W., Chinfak, N., Jandang, S., Kuanui, P., Khokiattiwong, S., & Chavanich,
1090	S. (2015). Effect of lower pH on settlement and development of coral, Pocillopora damicornis
1091	(Linnaeus, 1758). Ocean Science Journal, 50(2), 475-480. https://doi.org/10.1007/s12601-
1092	015-0043-z
1093	Von Euw, S., Zhang, Q., Manichev, V., Murali, N., Gross, J., Feldman, L. C., Gustafsson, T., Flach, C.,
1094	Mendelsohn, R., & Falkowski, P. G. (2017). Biological control of aragonite formation in stony
1095	corals. Science, 356(6341), 933–938. https://doi.org/10.1126/science.aam6371
1096	Wall, C. B., Mason, R. A. B., Ellis, W. R., Cunning, R., & Gates, R. D. (2017). Elevated pCO₂ affects tissue
1097	biomass composition, but not calcification, in a reef coral under two light regimes. Royal
1098	Society Open Science, 4(11), 170683. https://doi.org/10.1098/rsos.170683
1099	Wall, M., Fietzke, J., Crook, E. D., & Paytan, A. (2019). Using B isotopes and B/Ca in corals from low
1100	saturation springs to constrain calcification mechanisms. Nature Communications, 10(1).
1101	https://doi.org/10.1038/s41467-019-11519-9

1102	Webster, N. S., Negri, A. P., Botté, E. S., Laffy, P. W., Flores, F., Noonan, S., Schmidt, C., & Uthicke, S.
1103	(2016). Host-associated coral reef microbes respond to the cumulative pressures of ocean
1104	warming and ocean acidification. Scientific Reports, 6(1). https://doi.org/10.1038/srep19324
1105	Webster, N. S., Negri, A. P., Flores, F., Humphrey, C., Soo, R., Botté, E. S., Vogel, N., & Uthicke, S.
1106	(2013). Near-future ocean acidification causes differences in microbial associations within
1107	diverse coral reef taxa: pH effects on marine microbial associations. Environmental
1108	Microbiology Reports, 5(2), 243–251. https://doi.org/10.1111/1758-2229.12006
1109	Wijgerde, T., Silva, C. I. F., Scherders, V., van Bleijswijk, J., & Osinga, R. (2014). Coral calcification
1110	under daily oxygen saturation and pH dynamics reveals the important role of oxygen. Biology
1111	Open, 3(6), 489–493. https://doi.org/10.1242/bio.20147922
1112	Yang, SY., Lu, CY., Tang, SL., Das, R. R., Sakai, K., Yamashiro, H., & Yang, SH. (2020). Effects of
1113	ocean acidification on coral endolithic bacterial communities in Isopora palifera and Porites
1114	lobata. Frontiers in Marine Science, 7. https://doi.org/10.3389/fmars.2020.603293
1115	Zande, R. M., Achlatis, M., Bender-Champ, D., Kubicek, A., Dove, S., & Hoegh-Guldberg, O. (2020).
1116	Paradise lost: End-of-century warming and acidification under business-as-usual emissions
1117	have severe consequences for symbiotic corals. Global Change Biology, 26(4), 2203–2219.
1118	https://doi.org/10.1111/gcb.14998
1119	Zhou, G., Yuan, T., Cai, L., Zhang, W., Tian, R., Tong, H., Jiang, L., Yuan, X., Liu, S., Qian, P., & Huang, H.
1120	(2016). Changes in microbial communities, photosynthesis and calcification of the coral
1121	Acropora gemmifera in response to ocean acidification. Scientific Reports, 6(1).
1122	https://doi.org/10.1038/srep35971
1123	Zoccola, D., Innocenti, A., Bertucci, A., Tambutté, E., Supuran, C., & Tambutté, S. (2016). Coral
1124	carbonic anhydrases: Regulation by ocean acidification. Marine Drugs, 14(6), 109.
1125	https://doi.org/10.3390/md14060109
1126	







