

Two decades of seawater acidification experiments on tropical scleractinian corals: Overview, meta-analysis and perspectives

Mathilde Godefroid, Sam Dupont, Marc Metian, Laetitia Hédouin

▶ To cite this version:

Mathilde Godefroid, Sam Dupont, Marc Metian, Laetitia Hédouin. Two decades of seawater acidification experiments on tropical scleractinian corals: Overview, meta-analysis and perspectives. Marine Pollution Bulletin, 2022, 178, pp.113552. 10.1016/j.marpolbul.2022.113552. hal-03841318

HAL Id: hal-03841318 https://univ-perp.hal.science/hal-03841318v1

Submitted on 22 Jul 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0025326X2200234X Manuscript_ac4f9c0c1b6eb6109b782cb9a221f638

1	Title			
2	Two decades of seawater acidification experiments on tropical scleractinian corals: overview, meta			
3	analysis and pe	analysis and perspectives		
4				
5	Authors			
6	Mathilde Godefroid ^{1,2,*} , Sam Dupont ^{3,4} , Marc Metian ⁴ , Laetitia Hédouin ^{1,2}			
7				
8	Affiliations			
9	¹ PSL Research University: EPHE-CNRS-UPVD, USR 3278 CRIOBE, BP 1013, 98729 Papetoai, Mo'orea,			
10	French Polynesia			
11	² Laboratoire d'Excellence « CORAIL», Mo'orea, French Polynesia			
12	³ Department of Biological and Environmental Sciences, University of Gothenburg, Kristineberg			
13	Marine Research Station, Kristineberg 566, 45178 Fiskebäckskil, Sweden			
14	⁴ Radioecology	Laboratory International Atomic Energy Agency (IAEA), Marine Laboratories, 4 Quai		
15	Antoine 1er, 98000, Principality of Monaco			
16				
17				
18	*Corresponding author:			
19	Name:	Godefroid Mathilde		
20	Present address:	: Marine biology laboratory,		
21		Université Libre de Bruxelles,		
22		Av. F.D. Roosevelt, 50, CP 160/15		
23		1050 Bruxelles		
24		Belgium		

 25
 Phone:
 (+32)489 36 70 08

 26
 E-mail:
 godefroid.mathilde1@gmail.com

27 Abstract

28 Ocean acidification has emerged as a major concern in the last fifteen years and studies on the 29 impacts of seawater acidification on marine organisms have multiplied accordingly. This review 30 aimed at synthesizing the literature on the effects of seawater acidification on tropical scleractinians 31 under laboratory-controlled conditions. We identified 141 articles (published between 1999-2021) 32 and separated endpoints into 22 biological categories to identify global trends for mitigation and 33 gaps in knowledge and research priorities for future investigators. The relative number of affected 34 endpoints increased with pH intensity (particularly for endpoints associated to calcification and 35 reproduction). When exposed to pH 7.6-7.8 (compared to higher pH), 49% of endpoints were 36 affected. The diversity in experimental designs prevented deciphering the modulating role of coral 37 life stages, genera or duration of exposure. Finally, important bias in research efforts included most 38 experiments on adult corals (68.5%), in 27 out of 150 (18%) coral ecoregions and exclusively from 39 shallow-waters.

40

41 Keywords

42 Ocean acidification, Tropical scleractinian, pH, Calcification, Review

43

44 Acknowledgements

We thank Jacob et al. (2020) for making their R scripts available and Dr Moreau C. for his help in making Fig. 1. This work was funded by the Foundation de France for a project called "ACID REEFS", by Ministère de la Transition Écologique et Solidaire and the Foundation for Research on Biodiversity (FRB) for a project entitled "ACID REEFS". The IAEA is grateful for the support provided to its Environment Laboratories by the Government of the Principality of Monaco.

1. Introduction

51 Due to the increasing emissions of carbon dioxide into the atmosphere, the Ocean is warming and 52 getting more acidic. Ocean acidification (OA) has been identified as a global environmental threat 53 and included as the United Nations' Sustainable Development Goal 14.3, as well as one of the nine 54 planetary boundaries (Jagers et al., 2019; Rockström et al., 2009). OA impacts on marine species and 55 ecosystems are well documented, including effect on marine calcifiers, threatening coral reefs as well 56 as broader marine ecosystems (Doney et al., 2009; Feely, 2004; Gattuso & Hansson, 2011; Hendriks 57 et al., 2010; Jiang et al., 2019; Kroeker et al., 2010, 2013; Orr et al., 2005).

Most of the available knowledge on the effects of OA on marine organisms comes from short-term 58 59 laboratory experiments on isolated organisms (Kroeker et al., 2010). Fixed-term studies have the disadvantage of potentially under/overestimate the effects of OA, as some taxa may show 60 61 vulnerability/acclimatization in the long-term (Fantazzini et al., 2015). Similarly, other life stages than 62 those considered may show different vulnerability and controlled conditions does not consider the 63 indirect effects due to OA-driven ecological changes (Fabricius et al., 2011). In this context, 64 knowledge acquired through field experiments taking advantage of organisms that are naturally 65 exposed to OA (e.g. CO_2 vent systems) are very complementary as they account for the life-long 66 acclimatization of organisms. These studies already showed that OA may change reef community 67 composition and metabolism (Biscéré et al., 2019; Noonan et al., 2018). They also reported effects on 68 skeletal porosity (Prada et al., 2021) and a variety of responses to OA on calcification rate, suggesting 69 species-specific acclimatization to OA (Strahl et al., 2015).

70 Coral reefs have received particular attention as they are among the most severely threatened 71 ecosystem on Earth (Pandolfi, 2003; Raven, 2005). The effects of seawater acidification on corals 72 have been extensively studied (Chan & Connolly, 2013; Erez et al., 2011), but responses often differ 73 between studies depending on parameters such as tested populations, species and life-cycle stages 74 (Kawahata et al., 2019). These apparent conflicting results in the literature could also be attributed to 75 variations in experimental designs and methodologies. In this study, we reviewed the literature that 76 examined the effects of decreased pH on tropical scleractinian corals under laboratory-controlled 77 conditions. A meta-analysis based on 169 experiments conducted in 141 peer-reviewed articles 78 published since 1999 was performed, with four main objectives:

- 79 (1) To provide a semi-quantitative description of the evolution of research efforts testing the
 80 effects of seawater acidification on tropical scleractinian corals,
- 81 (2) To evaluate the effect of seawater acidification on coral biological functions,

82 (3) To investigate the modulating influence of exposure duration, coral life stages and coral83 genera on sensitivity to seawater acidification and,

84 (4) To highlight research gaps and bias.

Based on this review, we provide recommendations and perspectives for future research and anupdated baseline for scientists starting in this field of research.

- 87
- 88

2. Material and methods

89 The code used in this manuscript based Jacob et al. (2020): was on 90 https://doi.org/10.5281/zenodo.3694955.

91

92 <u>2.1 Literature search</u>

A literature search was performed using the following databases: Ocean Acidification – International 93 94 Coordination Centre portal for ocean acidification biological response data, Google Scholar, Scopus 95 (keywords such as "coral" AND "acidification"). We listed all experimental studies published before 96 2021, which mainly focused on the effects of experimental seawater acidification on tropical 97 scleractinian corals. Studies were selected according to the following criteria: 1) the experiment(s) 98 was(were) performed under laboratory-controlled conditions; 2) one treatment (at least) was 99 designed to test the effects of seawater acidification only (without interaction with another stressor); 100 3) a tropical scleractinian coral species (at least) was studied; 4) an acidified treatment was compared 101 with a control treatment (present-day conditions), with constant conditions (i.e. non-oscillating). In 102 situ experiment(s) of the effects of seawater acidification on biological functions were not included in the database but were considered for the discussion. Higher than present-day (control) pH 103 104 conditions (e.g. pre-industrial pH conditions) were not included in the database. When the statistics 105 lacked clarity/transparency or did not allow to be adequately decomposed into the database, the 106 article was excluded. The completeness of the results obtained was considered as satisfactory based 107 on "snowballing" (i.e. checking citations on reference lists of relevant articles until no further 108 relevant article could be found; Sayers, 2007).

109

110 <u>2.2 Main data extraction</u>

For each experiment selected in the database, the following information was extracted: Study location(s), study duration(s), coral genus and species, coral life stage(s), number of studied species, tested pHs, pH scale and endpoints.

114

115 <u>2.3 Data organization</u>

116 For analyses, six pH classes were defined: Class 1 for pH between 7.8 and 8, Class 2 for pH between 117 7.6 and 7.8, Class 3 for pH between 7.4 and 7.6, Class 4 for pH between 7.2 and 7.4, Class 5 for pH 118 between 7 and 7.2, Class 6 for pH below 7. Recent studies highlight the importance of local 119 adaptation to the present natural variability in the carbonate chemistry to define species sensitivity 120 to OA (e.g. Vargas et al., 2017). As a consequence, the definition of pH scenarios or categories 121 relevant in the context of OA should consider this natural variability. As diurnal and seasonal pH 122 variability is low in coral reef environments (e.g. Cyronak et al., 2020), pH within class 1 to 6 are 123 considered relevant in the context of OA. Similarly, duration, genus and life stage were also divided 124 into classes. Four duration classes were defined: Class 1 for very short-term experiments (<1 week), 125 Class 2 for short-term experiments (1-4 week(s)), Class 3 for medium-term experiments (1-3 126 month(s)), Class 4 for long-term experiments (>3 months); Three genus classes: Class 1 for species 127 from the genus Acropora, Class 2 for species from the genus Porites, Class 3 for species from the 128 genus Pocillopora; and three life stage classes were chosen: Class 1 for free-living non-calcifying early 129 life stages (Gametes, Gastrulae, Larvae), Class 2 for settled calcifying early life stages (Recruits, 130 Primary polyps, Juveniles) and Class 3 for the adults. All classes are summarized in Table 1.

131

132

Table 1. pH, Duration, genus and life stage classes used to classify this review.

Parameters	Classes	
	Class 1	7.8-8
	Class 2	7.6-7.8
рH	Class 3	7.4-7.6
p.,	Class 4	7.2-7.4
	Class 5	7.0-7.2
	Class 6	<7
	Class 1	<1 week
Duration	Class 2	1-4 week(s)
	Class 3	1-3 month(s)
	Class 4	>3 months
	Class 1	Acropora
Genus	Class 2	Porites
	Class 3	Pocillopora
	Class 1	Free-living early life stages
Life stage	Class 2	Settled early life stages
	Class 3	Adults

134 <u>2.4 Endpoints</u>

For each study and experiment, all measured endpoints were identified and divided into categories based on their biological significance. When the biological significance of an endpoint was not straightforward and/or when an endpoint could have several biological meanings, we attributed it to a single category based on the focus adopted by authors. In total, eight categories, divided into 22 subcategories, were established (Table 2).

140

141 Table 2. List of the biological categories and subcategories used in the database, with one example of endpoint142 measured.

Biological category	Biological subcatergory	Endpoint measured ¹
Feeding and excretion	Excretion	Ammonia excretion rate
	Feeding	Feeding rate
	Reproductive investment and success	Settlement rate
Fitness	Calcification	Net calcification rate
Filless	Mortality	Total mortality
	Growth and development	Linear extension rate
Cellular stress-response	Apoptosis	p53 expression level
mechanisms	Oxidative stress	Glutathione reductase expression level
mechanisms	Other molecular actors	HSP70
	Carbohydrate	Carbohydrate content
Biomolecules	Lipid	Total lipid content
Biomolecules	Protein	Protein content
	Energy content	Host ATP content
Despiration and	Photophysiology	Net photosynthetic rate
Respiration and	Respiration	Dark respiration rate
photophysiology	Productivity	Net productivity
Microbiome and endolithic algae	Bacterial community composition	Gammaproteobacteria content
Metabolome		Metabolite profile
Epigenetic mechanism		DNA methylation level
	Community structure	Planar cover of the community
Community	Community calcification	Light community calcification
	Respiration	Community net oxygen consumption

143 ¹This is a non-exhaustive list of examples. See the database in Supplementary Material for the full list of endpoints

144 measured.

145

146 For each endpoint, we reported the occurrence of a significant effect of exposure to pH as compared

to the highest tested pH: YES if the endpoint was significantly affected by pH; NO elsewhere (Jacob et

al., 2020). If multiple pH treatments were used and the statistics in the study did not compare the
effect of each treatment with the control, but rather reported the result as a statistical trend, a YES
was attributed to an endpoint when the trend was significant. For each category and subcategory,
the percentage of affected endpoints was calculated by dividing the number of affected endpoints
(*i.e.* number of YES) by the total number of endpoints in this category or subcategory (Jacob et al.,
2020).

154

155 <u>2.5 Database construction</u>

All selected experimental studies were grouped in a common database gathering all relevant information (main data, endpoints and effects as described from 2.2 to 2.4). The database is available as Supplementary Material.

- 159
- 160

3. Results & Discussion

161 <u>3.1 General trends in experimental seawater acidification research on tropical scleractinian corals</u>

162 *Literature trend*

In our database, we identified 169 experiments in 141 articles (listed in Supplementary Material) that 163 164 have evaluated the effects seawater acidification under laboratory-controlled conditions on tropical 165 scleractinian corals. The six best studied regions account for 69% of the studies: Australia (15%), 166 Japan (14%), French Polynesia (14%), Hawaii (10%), the USA (excluding Hawaii, 8%) and Taiwan (8%; 167 Fig. 2A). Six percent of the studies were carried out on corals that have been maintained and 168 acclimated in aquaria for years (referred to as "Aquaria acclimated"; Fig. 2A). These included the 169 species Stylophora pistillata that were held in aquaria in Monaco (Liew et al., 2018; Reynaud et al., 170 2003; Tambutté et al., 2015; Venn et al., 2013, 2019, 2020; Zoccola et al., 2016), but also by other 171 coral species obtained from commercial suppliers (Bove et al., 2020; Wijgerde et al., 2014). 172 Altogether, these data reveal that the effects of experimental seawater acidification on tropical 173 scleractinian corals have only been studied in 27 out of 150 coral ecoregions (18%; Veron et al., 2005, 174 2015). Certain parts of the world remain unexplored, with no study to date in the African reefs nor 175 around the coasts of India or Sri Lanka, and very few studies in the coral triangle (Figure 1), despite 176 being known as the most biodiverse area in the ocean, containing more than 76% of the world's 177 shallow-water reef-building corals (Asaad et al., 2018). Moreover, no study has ever focused on the 178 impact of seawater acidification on tropical mesophotic corals, even though they represent 2/3rd of the world coral reefs (Morais & Santos, 2018) and are expected to act as "deep refugia", by providing
a viable reproductive source for shallow reefs after disturbance (Bongaerts et al., 2010).

181 Experimental seawater acidification research on tropical scleractinians is relatively recent and has 182 increased exponentially since the 21st century (Fig. 2B). Similarly, in situ studies have only started to 183 develop even more recently, with a majority of studies (70%) published over the five past years (Fig. 184 2B). Among these, 80% took advantage of coral communities living under naturally acidified waters 185 such as submarine springs in Mexico (Crook et al., 2012, 2013; Martinez et al., 2019; Wall et al., 186 2019) and CO₂ vents in Papua New Guinea (Biscéré et al., 2019; Fabricius et al., 2011; Fantazini et al., 187 2015 ; Marcelino et al., 2017 ; Morrow et al., 2015 ; Noonan et al., 2016, 2018 ; O'Brien et al., 2018 ; 188 Prada et al., 2021 ; Smith et al., 2016 ; Strahl et al., 2015 ; Wall et al., 2017). Other studies compared 189 adjacent reef sites with distinct pH conditions (Barkley et al., 2015, 2017; Camp et al., 2017). Finally, 190 20% of studies have performed in situ controlled CO₂ enrichment such as FOCE experiments (Free 191 Ocean CO₂ Experiments; Albright et al., 2016, 2018; Doo et al., 2019; Georgiou et al., 2015; Kline et 192 al., 2019; Srednick et al., 2020). In situ studies possess the great advantage of being more ecologically 193 realistic than laboratory-based studies, as they account for the complexity of species interactions of 194 the whole reef community, under natural abiotic conditions (Albright et al., 2018). Yet, naturally 195 acidified sites are not always available and possess their own set of limitations. They are not perfect 196 predictors of future ocean ecology owing to temporal variability in pH, spatial proximity with other 197 populations and changes in other parameters such as seawater temperature (Prada et al., 2021; 198 Riebesell, 2008). Moreover, the manipulation of seawater parameters in situ is technically 199 challenging and comes at great expenses, which greatly limits its reproducibility.

200

201 pH scale

202 Best practices recommend to measure seawater pH using the total hydrogen ion concentration scale 203 (Total scale; Dickson et al., 2007). Yet, only 54.4% of experiments used this scale (Figure 2I). The rest 204 of the experiments measured pH on the NBS scale (US National Bureau of Standards, now referred to 205 as the IUPAC scale, 29.2%) or on the seawater scale (SWS; 5.3%). A minority of experiments (3.5%) 206 did not measure the pH but estimated its value from the other seawater carbonate chemistry 207 parameters, via CO2sys (Pierrot et al., 2006) or the R package seacarb (Gattuso et al., 2020). Finally, a 208 significant proportion (7.6%) did not specify how the pH was measured/calculated. Such heterogeneity in the pH scales used to measure seawater pH probably lies in the fact that it is, 209 210 together with the study of acid-base reactions in seawater, one of the most confused areas of marine 211 chemistry (Dickson, 1993). Above all, it is essential that the pH is defined on the same pH scale as

that of all acid-dissociation constants that are used with it (Riebesell et al., 2011). We highly
recommend future studies to focus on measuring total scale pH following the complete procedure
and best practices in SOP6a and SOP6b (Dickson et al., 2007).

215

216 Coral characteristics

217 The vast majority of studies have evaluated the effects of seawater acidification on adult 218 scleractinians (68.4%; Fig. 2C). 16.7% of studies were devoted to free-living (non-calcified) early life 219 stages (mainly larvae -13.9%-, followed by gastrulae -1.7%- and gametes -1.1%). Then, the most 220 studied settled (calcified) early life stages were recruits (7.8%), followed by primary polyps (4.4%) 221 and juveniles (2.8%; Fig. 2C). This focus on adult corals is a common trend also generally observed for 222 other marine organisms (Kroeker et al., 2010). For corals, the high attention devoted to adult stages 223 is likely due to the convenience of collection and husbandry, compared to early life stages, which 224 require specific knowledge on spawning events and high expertise for a successful rearing. However, 225 future studies should put greater emphasis on early life stages as they are often inferred to be more 226 vulnerable to environmental conditions than adults (Byrne, 2011). However, as only 4.7% of studies 227 considered multiple life history stages, the susceptibility among life stages remains largely 228 underexplored. Multigenerational studies are thus crucial as they allow to compare the sensitivity of 229 a species among life stages and to evaluate transgenerational carry over effects. To the best of our 230 knowledge, very few studies have evaluated these effects with corals, as compared to other taxa 231 (e.g. mollusks; Hettinger et al., 2012; Parker et al., 2012) or other stressors (e.g. global warming; 232 Dixon et al., 2015). Studies on scleractinians suggested that exposure of the parents to stressful 233 conditions had positive influence on coral offspring performances, when subsequently re-exposed 234 (e.g. metabolic acclimation for P. damicornis, Putnam & Gates, 2015; greater larval settlement and 235 survivorship for P. damicornis, Putnam, 2021). Yet, this likely vary between species (e.g. S. pistillata, 236 Bellworthy et al., 2018). Multigenerational studies are thus required to investigate parental (carry-237 over) effects, to decipher whether the exposure to seawater acidification in adults enhance offspring 238 tolerance, and to predict the future of coral reefs more accurately (Putnam et al., 2016, 2021).

Altogether, three genera stand for 65.1% of the experiments, with the most studied being *Acropora* (30.4%), followed by *Porites* (18.3%) and *Pocillopora* (16.3%; Fig. 2D). The most studied species from the database is *Acropora millepora*. Other well studied genera are *Montipora* (7.9%), *Stylophora* (6.3%) and *Seriatopora* (4.6%), followed by studies that have evaluated the effects of seawater acidification on the coral community (3.8%; Fig. 2D). The genera *Siderastrea, Pavona* and *Psammocora* each accounted for 1.7, 1.7 and 1.3% of the database, respectively. Finally, 6.7% of 245 experiments were performed on less studied genera (<1%, referred to as "Others"; Fig. 2D). These 246 include species from the genera Cyphastrea, Duncanopsammia, Favia, Favites, Fungia, Galaxea, 247 Goniopora, Isopora, Leptastrea, Litophyllon, Madracis, Montastrea, Mussismila, Orbicella, 248 Pseudodiploria and Turbinaria (Fig. 2D). Overall, these data show that a small number of coral species 249 are favoured in experimental seawater acidification experiments. As previously reported for heat 250 stress experiments on corals (review by McLachlan et al., 2020), this may be due to the facility of 251 working on highly studied species (*i.e.* with the availability of their genome), the presence of these 252 species within each region or to logistical constraints (e.g. permits, easiness of maintenance in 253 captivity). This finding also means that many species, and so several functional traits which are likely 254 to drive differential responses to acidification, remain uninvestigated.

255

256 Experimental design

257 The majority of experiments (63.9%) considered a single species of coral (Fig. 2E). The remaining 258 experiments used two to four (30.8%) or five to eight species (3.0%) and only 2.3% evaluated the 259 effects of seawater acidification on the coral community (Fig. 2E). Undoubtedly, it limits and 260 oversimplifies our understanding of the effects of OA on the biologically complex assemblage 261 characterizing coral reefs. On top of that, species interactions can also decrease the predictability in 262 species response to seawater acidification, since different pairings and spatial organization of corals 263 might significantly influence the observed endpoints (Brien et al., 2016; Evensen et al., 2015, 2016). 264 For these reasons, designing experiments based on the local species assemblage in terms of diversity, 265 abundance and spatial organization, should be prioritized in future investigations.

266 Furthermore, most experiments lasted between 1 and 4 week(s) (39.6%), with the other experiments 267 lasting less than a week (26.4%), one to three month(s) (24.7%) and more than three months (9.3%), 268 for a minority (Fig. 2F). This trend is likely explained by the constraints associated with long-term 269 studies, in terms of time, logistical or even financial limitations. Nevertheless, the duration of 270 exposure may influence coral sensitivity to seawater acidification. For instance, corals are able to 271 counteract the negative effects of acidification for a specific period through energy-demanding 272 processes, until reaching a time-dependent threshold beyond which it will no longer be able to 273 counter these effects (Castillo et al., 2014; Schoepf et al., 2013). For these reasons, short-term 274 studies should be avoided, and long-term experimental studies prioritized, when possible. In that aspect, field experimentations/observations at CO₂ vents have the benefit to provide natural long-275 276 term studies (e.g. Biscéré et al., 2019; Fabricius et al., 2011; Martinez et al., 2019; Morrow et al., 277 2015; Noonan et al., 2018; Noonan & Fabricius, 2016; Prada et al., 2021; Strahl et al., 2015).

278

279 Endpoints

280 Most experiments worked on the effects of seawater acidification on the fitness (57.3%), respiration 281 and photophysiology (23.9%) and biomolecules (7.1%) of tropical scleractinians (Fig. 2G). The other 282 endpoints studied were involved in cellular stress response mechanism(s) (4.3%), community 283 response(s) (3.5%), microbiome and endolithic algae composition (2.0%), epigenetic effect(s) (0.8%) 284 and/or feeding and excretion (0.8%; Fig. 2G). Among those that have worked on the effects of 285 seawater acidification on coral fitness, calcification was the most studied endpoint (52.6%), followed 286 by growth and development (25.5%), reproductive investment and success (10.9%) and mortality 287 (10.9%; Fig. 2H).

288

289 <u>3.2 Biological effects of pH on tropical scleractinian corals</u>

290 General trends

291 Overall, the relative number of affected endpoints increases with decreasing pH (Fig. 3). Twenty-six 292 percent of endpoints were affected between pH 7.8 and 8, 46.2% between pH 7.6 and 7.8, 48.3% 293 between pH 7.4 and 7.6, 46.3% between pH 7.2 and 7.4, 96.4% between pH 7.0 and 7.2 and 100% 294 below pH 7.0. However, it should be noted that the number of available articles and data is not the 295 same for each pH category. Studies have mainly focused on pH 7.8-7.6 (n endpoints=608), followed 296 by pH 8.0-7.8 (n=519), pH 7.6-7.4 (n=203), pH 7.4-7.2 (n=162), pH 7.2-7.0 (n=56) and below 7.0 (n=4). 297 Similarly, differences in experimental designs within each pH category is potentially a strong 298 confounding factor, mixing results from experiments of multiple duration of exposure, coral life-299 stages, coral species, among others. This may particularly limit our interpretations for the less-300 studied pH categories.

301

302 Fitness

The decrease in pH and $[CO_3^{2-}]$ through OA is believed to increase the energetic cost of calcification for marine calcifiers (Gattuso & Hansson, 2011). Known as one of the biggest threats related to OA, calcification, the ability to build and maintain calcium carbonate structures, was among the most studied parameters (52.6% of studies). Results showed that 43.4% of endpoints related to calcification were affected for pH 7.8-8.0 (n=152), 55.3% for pH 7.6-7.8 (n=244), 66.7% for pH 7.4-7.6 (n=96), 85.7% for pH 7.2-7.4 (n=42), 96.9% for pH 7.0-7.2 (n=33) and 100% for pH below 7.0 (n=2) (Fig. 3). Similarly, growth and development (*i.e.* coral linear extension, biomass, larval size, recruits 310 diameter, etc.) were increasingly affected with decreasing pH, with 29.5% of affected endpoints for 311 pH 7.8-8.0 (n=44), 44.4% for pH 7.6-7.8 (n=63), 61.1% for pH 7.4-7.6 (n=18), 73.7% for pH 7.2-7.4 312 (n=19), 88.9% for pH 7.0-7.2 (n=9) and 100% for pH below 7.0 (n=1). Fewer studies worked on the 313 effects of seawater acidification on reproductive investment and success (10.9%). Yet, they showed that 50.0% of reproductive endpoints were affected for pH 7.8-8.0 (n=20), 75.0% for pH 7.6-7.8 314 315 (n=16), 100% for pH 7.2-7.4 (n=1) as well as for pH below 7 (n=1). Noticeable effects on reproduction 316 included effects on sperm flagellar motility (Morita et al., 2010), time to gastrulation (Chua et al., 317 2013) and larval settlement (Albright et al., 2010; Doropoulos et al., 2012; Jiang et al., 2020; Pitts et 318 al., 2020; Viyakarn et al., 2015). Finally, effects of seawater acidification on coral mortality were only 319 showed under pH 7.6-7.8 (37.5% of affected endpoints, n=24). Mortality on adult colonies included 320 partial mortality (local tissue loss; Bahr et al., 2016; Kavousi et al., 2016) and total mortality (death of 321 the colonies; Yang et al., 2020). Studies also showed mortality of coral larvae (Cumbo et al., 2013; 322 Doropoulos et al., 2012) and recruits (Dufault et al., 2012, 2013) under seawater acidification.

323

324 Respiration and photophysiology

325 Measurement of the coral host respiration rate is often used as an indicator of the metabolism's 326 state/activity level. Moreover, the proper functioning of the algal symbiont photosynthetic activity is 327 fundamental as it provides an energy source as photosynthates to the host (Muscatine et al., 1981). 328 As the host, coral has some control over the photosynthetic activity of the algal symbionts (by 329 contributing to the low pH of the symbiosome via H⁺-ATPase, Barott et al., 2015), maintaining this 330 vital function is thus critical. The endpoints related to coral host respiration rate and algal symbiont 331 photosynthetic activity were poorly affected by seawater acidification, regardless of the pH level (Fig. 332 3). Despite this, studies on the effects of seawater acidification on photosynthetic processes gave 333 ambiguous results, with multiple studies showing an increased photosynthetic rate under low pH 334 (Anderson et al., 2019; Biscéré et al., 2019; Langdon, 2005). Response heterogeneity could be linked 335 to strain-specific responses by distinct Symbiodinium sp. (Brading et al., 2011; Graham & Sanders, 336 2016) or even attributed to differences in experimental parameters such as light or nutrients, which 337 both contributes to photosynthesis (Chauvin et al., 2011). It is expected that the effect of seawater 338 acidification on the coral's metabolism is small as compared to the effect of increasing temperature (Agostini et al., 2013; Cumbo et al., 2013; Edmunds & Burgess, 2016; Van der Zande et al., 2020). 339

340

341 Biomolecules

Quantification of biomolecules such as lipids, carbohydrates and proteins is often used as an 342 343 indicator of energetic reserves of organisms. As such, their depletion under stressful conditions is 344 often related to increased metabolic demands in ectotherms (Deutsch et al., 2008). Endpoints 345 related to lipid, protein and carbohydrate metabolisms seemed to remain relatively stable under decreased pH (Fig. 3). Yet, for lipids, 55.6% of endpoints were affected for pH 7.6-7.8 (n=9) and 33.3% 346 347 for pH 7.4-7.6 (n=6). Reduction of lipid content may show that maintenance of coral performances 348 and calcification under seawater acidification increase energetic costs, which is met through 349 catabolism of energy reserves. The subcategory "energy content" (here, the sum of lipids, 350 carbohydrates and proteins) showed to be greatly affected at pH 7.6-7.8, but this requires further 351 attention as it has only been studied in one experiment (Wall et al., 2017).

352

353 Cellular stress-response mechanism

354 Changes in the environment, like OA, are likely to drive cellular stress response, a universally 355 conserved mechanism used to protect macromolecules within cells from the potential damage that 356 OA may cause (Kaniewska et al., 2012). Here, a number of studies indicated that endpoints involved 357 in cellular stress-response mechanisms were affected by seawater acidification (Fig. 3). In the 358 subcategory "other molecular actors", 19.2% of endpoints were affected for pH 7.8-8.0 (n=26), 16.7% 359 for pH 7.6-7.8 (n=6), 0% for pH 7.4-7.6 (n=2) and 100% for pH 7.2-7.4 (n=1). These were attributed to genes involved in innate immunity (Kaniewska et al., 2012) and heat shock response (Liew et al., 360 361 2018; Moya et al., 2015; Putnam et al., 2013). Thirty-one percent of endpoints associated with 362 oxidative stress were affected at pH 7.8-8.0 (n=26) and 36.6% at pH 7.6-7.8 (n=41). These were 363 essentially affecting the activity level of antioxidant enzymes (Bielmyer-Fraser et al., 2018; Jiang et 364 al., 2019; Soriano-Santiago et al., 2013) or genes involved in the antioxidant response (Kaniewska et 365 al., 2012; Moya et al., 2015). These later studies also indicated up/down regulation of anti-apoptotic 366 genes, as showed with 50% of endpoints affected under pH 7.8-8.0 (n=2) and 100% under pH 7.6-7.8 367 (n=9).

368

369 Other biological category

Coral reefs form complex network of interactions between species. Each of its component plays a role and contributes to the overall equilibrium of the reef as well as its resilience to environmental change. Working on the coral community is therefore necessary as it allows considering the interactions between coral species and therefore provides more realistic responses. Yet, few studies have experimentally worked on the effects of seawater acidification on coral community, likely owing to the difficulty of having a realistic community in an aquarium. For this aspect, we suggest to refer to the numerous *in situ* studies, which naturally consider the coral community. Endpoints indicated that community structure was changed under pH 7.6-8.0 (100% of affected endpoints, n=2; Edmunds et al., 2019, 2020).

As acidifying conditions affect corals, their energy requirements increase and must be compensated in order to sustain the proper biological functioning of the coral. This may be studied by looking at feeding rate (the input of energy *via* heterotrophy) and excretion rate (ammonium ion concentration; the output of energy through protein catabolism) of corals. Towle et al. (2015) indicated that coral can buffer OA effects by increasing feeding rates under pH 7.8-8.0 and Edmunds & Wall (2014) showed that catabolism of proteins and amino acids is accelerated by OA (pH 7.8-8.0), supporting that OA may perturb protein metabolism.

386 Phenotypic plasticity has the potential to facilitate rapid acclimatization of corals, a requirement in 387 view of the rapid rate of change in the environment, driven by human activities. Mechanisms of rapid 388 acclimatization may include changes in microbial community, with 50% of endpoints affected for pH 389 7.8-8.0 (n=4), 50% for pH 7.6-7.8 (n=4), 50% for pH 7.6-7.4 (n=2) and 87.5% for pH 7.2-7.4 (n=16) 390 (Meron et al., 2011; Webster et al., 2013, 2016; Yang et al., 2020; Zhou et al., 2016). Another potential mechanism is epigenetics (i.e. the heritable postsynthesis modification of DNA or DNA-391 392 associated proteins, without a change in the DNA sequence itself, Feil & Fraga, 2012), with 50% of 393 endpoints affected for pH 7.6-7.8 (n=2) and 100% for pH 7.2-7.4 (n=1) (Liew et al., 2018; Putnam et 394 al., 2016). Finally, metabolomic profiling is a useful molecular tool that allows quantifying the 395 intermediates and products of many biochemical processes. Putnam et al. (2016) found lower 396 amount of metabolites under OA, suggesting a general suppression of metabolic activities. These 397 techniques have started to be used on corals recently and deserve more attention in future 398 investigations.

399

400 Gene expression

Understanding the fitness consequences of observed changes in endpoints is not always straightforward. This is particularly evident for changes in gene expression. An over or downregulations of a given gene does not implicitly translate into a positive or negative effect on protein level, function or fitness (Feder & Walser, 2005; Koussounadis et al., 2015; Greenbaum et al., 2003). Moreover, the intensity of an observed change at the gene expression level that is biologically significant is often poorly understood. For example, a 200% change in the expression of genes involved in calcification does not necessarily translate into a 200% change in calcification. For these
 reasons, we did not use the gene expression data to infer biological impacts of seawater acidification.

409

410 <u>3.3 Biological effects of pH 7.6-7.8 on tropical scleractinian corals</u>

Many factors can modulate the biological response of tropical scleractinian corals to seawater acidification, including the tested species, life-history stage, tested pH or duration of exposure. As shown in the previous section, the effects on various endpoints tend to increase with decreasing pH. To better understand the role of the other modulating factors while minimizing the effect of the pH intensity, we decided to focus on pH class 2 (pH 7.6-7.8) for 2 reasons: (i) it is the class with the most available data (n endpoints=608); and, (ii) this range of pH corresponds to projected scenarios expected for 2100 (SSP3-7.0 and SSP5-8.5; IPCC, 2021).

418

419 General trends

Tested endpoints were only moderately affected under pH 7.6-7.8 (Fig. 4). No strong effects of the duration of exposure, coral genus or life stage could be identified, even when endpoints were separated into different categories.

423

424 *Effects of the duration of exposure*

425 The effect of the duration of the exposure in seawater acidification experiment is still poorly 426 understood (Cumbo et al., 2013). Time-dependent response patterns were previously documented 427 (Castillo et al., 2014; Schoepf et al., 2013). For example, Castillo et al. (2014) observed an increase in 428 the calcification rate of S. siderea until the 60th day of exposure to acidified conditions, which 429 thereafter started to decline. Similar results were reported for A. millepora, with a decrease of 430 calcification that appeared only during the second half of the experiment (viz. after 1 month; Schoepf 431 et al., 2013). In these specific cases, the parabolic responses to decreased pH would not have been 432 detected if the duration of the experiments was shorter. These results might result from the ability of 433 corals to compensate for the extra energy costs associated with the exposure to decreased pH, until 434 reaching a time-dependent threshold beyond which the organism does not have the energy reserve 435 allowing to counteract the extra costs associated with the stress. In other taxonomic groups, a 436 different pattern was observed with a milder effect observed over longer period of time. For 437 example, Dupont et al. (2013) showed that the strong negative effect of pH on adult sea urchin

438 fecundity observed after a 4-month exposure disappeared after 16 months. The negative effect 439 observed over the first part of the experiment was interpreted as short-term costs associated with 440 physiological acclimation to the new low pH environment. These examples highlight the value of 441 long-term experiments.

442

443 Effects of the coral life stage

Early life stages are generally more vulnerable to environmental conditions than adults (Byrne, 2011), 444 445 so dismissing the effects of decreasing pH on multiple life stages in current prediction models would 446 represent an important bias for the projection of the fate of coral reef ecosystems. Indeed, if pH 447 disrupts early-life processes, it would compromise sexual recruitment and therefore affect future population dynamics and the ability of coral reefs to recover from a disturbance (Caroselli et al., 448 449 2019). This question has received relatively poor attention and experiments that have considered 450 more than one life stage are few (5%). Albright et al. (2010) suggested that OA has the potential to 451 affect multiple, early-life history stages of Acropora palmata, with negative effects on gametes 452 fertilization, larval settlement and post settlement growth. More experiment addressing this 453 question and considering multiple life stages are thus crucial.

454

455 Effects of the coral genera

The biological responses of corals to decreased pH are highly variable and often considered to be 456 457 species-specific (Bahr et al., 2016; Bedwell-Ivers et al., 2017; Bove et al., 2019; Brown & Edmunds, 458 2016; Edmunds et al., 2012; Hoadley et al., 2015; Sekizawa et al., 2017; Strahl et al., 2015). Several 459 other parameters were proposed to explain the diversity of response observed between studies. 460 While methodological differences could contribute to the difference between studies such as light 461 intensities (Crawley et al., 2010; Dufault et al., 2013), feeding regime (Comeau et al., 2013; Edmunds, 462 2011; Houlbrèque et al., 2015) or water motion (Comeau et al., 2015), acclimation (i.e. their life history; Strahl et al., 2015), competition (Brien et al., 2016; Evensen et al., 2015; 2016; Horwitz et al., 463 464 2017), or even the size of the fragments used (Edmunds & Burgess, 2016), biological differences 465 between taxa could explain the variability in coral sensitivity to seawater acidification. First, the 466 "biomass plasticity" hypothesis (Edmunds, 2011) supports that corals reduce their sensitivity by 467 increasing tissue thickness at low pH (Comeau et al., 2013; 2014; Edmunds, 2011). Indeed, thicker 468 tissue allow isolating the site of calcification from the surrounding seawater, better protecting the zooxanthellae by changing the intensity and internal properties of the internal light field (Dimond et 469

470 al., 2012; Loya et al., 2001), as well as providing more energy reserves, which may be advantageous 471 under low pH (as for Turbinaria reniformis, Hoadley et al., 2015). Massive Porites is known for having 472 an unusually thick tissue layer that penetrates deeply into the skeleton (Edmunds, 2011). Second, the 473 "pH regulation" hypothesis (Holcomb et al., 2015) suggests that scleractinians which have a high 474 control over the regulation of their calcifying fluid's pH will be less affected by low pH conditions. 475 Indeed, as the concentration of H^+ in seawater increases under OA, the concentration gradient 476 between seawater and the site of calcification increases accordingly (Cohen & McConnaughey, 2003; 477 Jokiel, 2011; Ries, 2011; Von Euw et al., 2017). Transporting H⁺ will require more energy for the coral, 478 which may alter calcification (McCulloch et al., 2012). As such, the coral's capacity/strategy to 479 overcome higher energy requirements (through efficient zooxanthellae photosynthesis, McCulloch et 480 al., 2012) or increased heterotrophic feeding (Towle et al., 2015) under more acidic conditions is also 481 crucial in explaining interspecific differences in sensitivity. Despite this, increasing heterotrophic 482 feeding under low pH conditions was not experimentally proven in all coral genera yet (such as 483 Porites; Grottoli et al., 2006). This goes along with the hypothesis suggesting that differences in 484 skeletal porosity may also explain differences in sensitivity of calcification among species. Indeed, 485 scleractinians with highly porous skeletons require less CaCO₃ to calcify than corals that have low 486 porosity skeletons, so these species may be less susceptible to seawater acidification (Brown & 487 Edmunds, 2016; Jokiel, 2011). Finally, the "growth rate" hypothesis (Comeau et al., 2014) states that 488 species characterized by higher calcification rate will be more vulnerable. Previous studies suggested 489 that different calcification strategies exist across populations (Bove et al., 2019; Leong & Pawlik, 490 2010; Metcalfe & Monaghan, 2001). It relies on the theory that fast-growing coral species are more 491 likely to be impaired under stressful conditions as their energy requirements for growth are higher 492 than slow-growing species, which mainly store energy when the environment is favorable, allowing 493 them to calcify normally under less-favorable conditions. However, this trend was not verified by 494 Chan & Connolly (2013).

495

496 Evolutionary potential of corals under OA: local adaptation, carry over effects, parental effects

Overall, there is very little information on the potential of marine organisms to adapt to OA. To better understand the evolutionary responses of scleractinians to global changes, the best approach is to evaluate the effects of environmental drivers through multiple generations (Putnam et al., 2016). While trans-generational effects of OA has received more attention in the previous years (Allan et al., 2014; Donelson et al., 2012; Dupont et al., 2013; Hettinger et al., 2012; Kurihara et al., 2019; Miller et al., 2012; Munday, 2014; Parker et al., 2012), studies evaluating these effects on 503 scleractinian corals are few, likely due to their long life span. Among the available studies, evidences 504 suggested that trans-generational acclimatization processes (e.g. epigenetics) have the potential to 505 influence the sensitivity of corals to changing conditions (Putnam et al., 2016). This approach is of 506 great importance to determine whether the exposure as an adult might enhance their descendant 507 tolerance in order to better assess the status of future coral reef ecosystems. A high variability of 508 response between individual corals and a high heritability for corals exposed to low pH conditions 509 was observed, suggesting an adaptation potential (Jury et al., 2019). As highlighted by Putnam 510 (2021), non-genetic, yet heritable phenotypic plasticity, will have significant ecological and 511 evolutionary implications for sessile marine organisms persistence under rapid environmental 512 changes. We support the need to consider the interactions of rapid and evolutionary responses 513 (microbiome, epigenetic effects, parental carry-over effects; see Putnam, 2021), that have been 514 overlooked so far, but deserve more attention as they provide additional avenues for rapid 515 acclimatization.

516

517 The future of the reef: past environmental exposure, multiple stressors/drivers interactions

518 OA acts in conjunction with a myriad of other stressors/drivers, which presence and intensity vary 519 locally. Among those, global ocean warming has received particular attention as it represents the 520 main stressor on the reefs, inducing coral bleaching and widespread mortality in multiple regions. 521 Studies considering a combination of increasing temperatures with seawater acidification often 522 showed additive effects, thus worsening the effects observed when considering a single stressor (e.g. 523 Albright & Mason, 2013; Anlauf et al., 2011; Anthony et al., 2008; Horvath et al., 2016; Kavousi et al., 524 2016; Reynaud et al., 2003). Then, to predict more realistically the future of coral reefs, their 525 vulnerability must be assessed over long term and consider the stressors experienced by the reef 526 organisms in the past. There is a temporality in the action of stressors on the reef that will condition 527 the response of reef organisms. For example, multiple evidences showed that corals that have gone 528 through repeated heat stress events in the past may have become more tolerant to future heat 529 stress (Guest et al., 2012; Maynard et al., 2008; Thompson & van Woesik, 2009). Similarly, decreased 530 thermal susceptibility was described with corals preconditioned to mild heat stress for a short period 531 (Bay & Palumbi, 2015; Bellantuono et al., 2012a, 2012b; Middlebrook et al., 2008). In addition, OA 532 will impair coral reefs previously altered by several bleaching events. Coral reef communities will be fragilized by the intensity and frequency of bleaching and it is likely that their capacity to endure 533 additional stressor as OA will be highly detrimental. 534

536		4. Research gaps and perspectives			
537 538 539 540 541	Based on all aforementioned points, we selected a list of priorities and actions that should be addressed for future experimental works investigating the effects of seawater acidification on tropical scleractinians (Table 3).				
542 543	Table 3. List of actions and priorities for future research on the effects of seawater acidification on tropical scleractinians.				
	Туре	Recommended action(s)			
	Coral characteristics	Investigate corals from less-studied ecoregions, mesophotic depths			
	Experimental design	Consider multigenerational approaches Use multiple life-history stages, including early stages Realism: select the diversity and abundance of species based on the local assemblage, favour multi-species approach Include key species interacting with the tested species Avoid short-term exposure, consider multiple duration of exposure when feasible			
	Carbonate chemistry and scenarios	Monitor and consider the local variability of carbonate chemistry while defining scenarios and rely on local projections Include the natural variability for the carbonate chemistry and other key environmental drivers (multiple stressors) Apply best practices for carbonate chemistry parameters measurement (Dickson et al., 2007). Report pH in total scale.			
	Biological endpoints	Focus on endpoints to better understand the acclimatization capacity: epigenetic effects, microbiome and endolithic algae composition Consider multiple complementary approaches: genetic and physiological tools			

545

546

547 Funding

- 548 This work was supported by the "Fondation de France" for a project called "ACID REEFS", by the
- 549 "Ministère de la Transition Ecologique et Solidaire" and the Foundation for Research on Biodiversity
- 550 (FRB) for a project entitled "ACID REEFS2".

551 Figure legends

552

Figure 1. World map of coral reefs that have been the subject of an experimental study (blue circles)
on the effects of seawater acidification. Circle size: number of experimental study. Red colour: Global
distribution of coral reefs (UNEP-WCMC, WorldFish Centre, WRI, TNC, 2010; version of March 2021).

556 Figure 2. General trends in the experimental research of the effects of seawater acidification on 557 biological functions of tropical scleractinian corals. (A) Number of experiments per country/location that experimentally examined the effects of seawater acidification on tropical scleractinian corals 558 559 (total n=169). (B) Number of ex situ (total n= 169) and in situ (total n=27) studies published before 560 2021 that experimentally examined the effects of seawater acidification on tropical scleractinian 561 corals. Relative proportion of (C) coral life stages (n=180) and (D) coral genera (n=240) studied in the 169 experiments that looked at the effects of seawater acidification on tropical scleractinian corals 562 563 before 2021. Relative proportion of experimental design by (E) assemblages (n=169; i.e. number of 564 studied species per experiment) and (F) duration of exposure (n=182) to seawater acidification in the 565 same 169 experiments. (G) Relative proportion of the main biological categories studied in the same 566 169 experiments (n=255), with (H) specific focus on the fitness category. (I) Relative proportion of 567 the pH scale used in the 169 experiments. Some of these bar charts have a (n) superior to 169 568 because some of the 169 experiments that were selected used more than one life stage, genera, 569 duration of exposure and/or biological endpoint.

Figure 3. Effects of pH intensity on tropical scleractinian corals biological functions. Heatmaps
indicate the number of endpoints (circle size) and the proportion of affected endpoints (color, in %)
in each biological subcategory. Full red (100%) indicates that all endpoints are affected and full green
(0%) that no endpoints were affected.

574 **Figure 4.** Effects of pH 7.6-7.8 on tropical scleractinian corals biological functions according to the 575 duration of exposure, the coral genus and life stage. Heatmaps indicate the number of endpoints

- 576 (circle size) and the proportion of affected endpoints (color, in %) in each biological subcategory. Full
- 577 red (100%) indicates that all endpoints are affected and full green (0%) that no endpoints were
- 578 affected.

579 References

- Agostini, S., Fujimura, H., Higuchi, T., Yuyama, I., Casareto, B. E., Suzuki, Y., & Nakano, Y. (2013). The effects of thermal and high-CO2 stresses on the metabolism and surrounding microenvironment of the coral Galaxea fascicularis. *Comptes Rendus Biologies*, *336*(8), 384– 391. https://doi.org/10.1016/j.crvi.2013.07.003
- 585 Albright, R., Caldeira, L., Hosfelt, J., Kwiatkowski, L., Maclaren, J. K., Mason, B. M., Nebuchina, Y., Ninokawa, A., Pongratz, J., Ricke, K. L., Rivlin, T., Schneider, K., Sesboüé, M., Shamberger, K., 586 587 Silverman, J., Wolfe, K., Zhu, K., & Caldeira, K. (2016). Reversal of ocean acidification 588 enhances net coral reef calcification. Nature, 531(7594), 362-365. 589 https://doi.org/10.1038/nature17155
- Albright, R., & Mason, B. (2013). Projected near-future levels of temperature and pCO2 reduce coral
 fertilization success. *PLoS ONE*, *8*(2), e56468. https://doi.org/10.1371/journal.pone.0056468
- Albright, R., Mason, B., Miller, M., & Langdon, C. (2010). Ocean acidification compromises
 recruitment success of the threatened Caribbean coral Acropora palmata. *Proceedings of the National Academy of Sciences, 107*(47), 20400–20404.
 https://doi.org/10.1073/pnas.1007273107
- Albright, R., Takeshita, Y., Koweek, D. A., Ninokawa, A., Wolfe, K., Rivlin, T., Nebuchina, Y., Young, J.,
- 597 & Caldeira, K. (2018). Carbon dioxide addition to coral reef waters suppresses net community
 598 calcification. *Nature*, 555(7697), 516–519. https://doi.org/10.1038/nature25968
- Allan, B. J. M., Miller, G. M., McCormick, M. I., Domenici, P., & Munday, P. L. (2014). Parental effects
 improve escape performance of juvenile reef fish in a high-CO₂ world. *Proceedings of the Royal Society B: Biological Sciences, 281*(1777), 20132179.
 https://doi.org/10.1098/rspb.2013.2179

- Anderson, K. D., Cantin, N. E., Casey, J. M., & Pratchett, M. S. (2019). Independent effects of ocean
 warming versus acidification on the growth, survivorship and physiology of two Acropora
 corals. *Coral Reefs*, *38*(6), 1225–1240. https://doi.org/10.1007/s00338-019-01864-y
- Anlauf, H., D'Croz, L., & O'Dea, A. (2011). A corrosive concoction: The combined effects of ocean
 warming and acidification on the early growth of a stony coral are multiplicative. *Journal of Experimental Marine Biology and Ecology, 397*(1), 13–20.
 https://doi.org/10.1016/j.jembe.2010.11.009
- Anthony, K. R. N., Kline, D. I., Diaz-Pulido, G., Dove, S., & Hoegh-Guldberg, O. (2008). Ocean
 acidification causes bleaching and productivity loss in coral reef builders. *Proceedings of the National Academy of Sciences*, 105(45), 17442–17446.
 https://doi.org/10.1073/pnas.0804478105
- Asaad, I., Lundquist, C. J., Erdmann, M. V., Van Hooidonk, R., & Costello, M. J. (2018). Designating
 spatial priorities for marine biodiversity conservation in the coral triangle. *Frontiers in Marine Science*, *5*. https://doi.org/10.3389/fmars.2018.00400
- Bahr, K. D., Jokiel, P. L., & Rodgers, K. S. (2016). Relative sensitivity of five Hawaiian coral species to
 high temperature under high-pCO2 conditions. *Coral Reefs*, 35(2), 729–738.
 https://doi.org/10.1007/s00338-016-1405-4
- 620 Barkley, H. C., Cohen, A. L., Golbuu, Y., Starczak, V. R., DeCarlo, T. M., & Shamberger, K. E. F. (2015).
- 621 Changes in coral reef communities across a natural gradient in seawater pH. *Science* 622 *Advances*, 1(5), e1500328. https://doi.org/10.1126/sciadv.1500328
- Barkley, H. C., Cohen, A. L., McCorkle, D. C., & Golbuu, Y. (2017). Mechanisms and thresholds for pH
 tolerance in Palau corals. *Journal of Experimental Marine Biology and Ecology, 489,* 7–14.
 https://doi.org/10.1016/j.jembe.2017.01.003
- Barott, K. L., Venn, A. A., Perez, S. O., Tambutté, S., & Tresguerres, M. (2015). Coral host cells acidify
 symbiotic algal microenvironment to promote photosynthesis. *Proceedings of the National Academy of Sciences*, *112*(2), 607–612. https://doi.org/10.1073/pnas.1413483112

- Bay, R. A., & Palumbi, S. R. (2015). Rapid acclimation ability mediated by transcriptome changes in
 reef-building corals. *Genome Biology and Evolution*, 7(6), 1602–1612.
 https://doi.org/10.1093/gbe/evv085
- Bedwell-Ivers, H. E., Koch, M. S., Peach, K. E., Joles, L., Dutra, E., & Manfrino, C. (2017). The role of in
 hospite zooxanthellae photophysiology and reef chemistry on elevated pCO2 effects in two
 branching Caribbean corals: Acropora cervicornis and Porites divaricata. *ICES Journal of*
- 635 *Marine Science*, 74(4), 1103–1112. https://doi.org/10.1093/icesjms/fsw026
- Bellantuono, A. J., Granados-Cifuentes, C., Miller, D. J., Hoegh-Guldberg, O., & Rodriguez-Lanetty, M.
- 637 (2012). Coral thermal tolerance: tuning gene expression to resist thermal stress. *PLoS ONE*,
 638 7(11), e50685. https://doi.org/10.1371/journal.pone.0050685
- Bellantuono, A. J., Hoegh-Guldberg, O., & Rodriguez-Lanetty, M. (2012). Resistance to thermal stress
 in corals without changes in symbiont composition. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1731), 1100–1107. https://doi.org/10.1098/rspb.2011.1780
- 642 Bellworthy, J., Menoud, M., Krueger, T., Meibom, A., & Fine, M. (2018). Developmental carry over 643 effects of ocean warming and acidification in corals from a potential climate refugium, Gulf
- 644 of Aqaba. Journal of Experimental Biology. https://doi.org/10.1242/jeb.186940
- Bielmyer-Fraser, G. K., Patel, P., Capo, T., & Grosell, M. (2018). Physiological responses of corals to
 ocean acidification and copper exposure. *Marine Pollution Bulletin*, *133*, 781–790.
 https://doi.org/10.1016/j.marpolbul.2018.06.048
- Biscéré, T., Zampighi, M., Lorrain, A., Jurriaans, S., Foggo, A., Houlbrèque, F., & Rodolfo-Metalpa, R.
 (2019). High pCO₂ promotes coral primary production. *Biology Letters*, *15*(7), 20180777.
 https://doi.org/10.1098/rsbl.2018.0777
- Bongaerts, P., Ridgway, T., Sampayo, E. M., & Hoegh-Guldberg, O. (2010). Assessing the 'deep reef
 refugia' hypothesis: Focus on Caribbean reefs. *Coral Reefs*, 29(2), 309–327.
 https://doi.org/10.1007/s00338-009-0581-x

- Bove, C. B., Ries, J. B., Davies, S. W., Westfield, I. T., Umbanhowar, J., & Castillo, K. D. (2019).
 Common Caribbean corals exhibit highly variable responses to future acidification and
 warming. *Proceedings of the Royal Society B: Biological Sciences, 286*(1900), 20182840.
 https://doi.org/10.1098/rspb.2018.2840
- Bove, C. B., Whitehead, R. F., & Szmant, A. M. (2020). Responses of coral gastrovascular cavity pH
 during light and dark incubations to reduced seawater pH suggest species-specific responses
 to the effects of ocean acidification on calcification. *Coral Reefs, 39*(6), 1675–1691.
 https://doi.org/10.1007/s00338-020-01995-7
- Brading, P., Warner, M. E., Davey, P., Smith, D. J., Achterberg, E. P., & Suggett, D. J. (2011).
 Differential effects of ocean acidification on growth and photosynthesis among phylotypes of *Symbiodinium* (Dinophyceae). *Limnology and Oceanography*, *56*(3), 927–938.
 https://doi.org/10.4319/lo.2011.56.3.0927
- Brien, H. V., Watson, S.-A., & Hoogenboom, M. O. (2016). Presence of competitors influences
 photosynthesis, but not growth, of the hard coral *Porites cylindrica* at elevated seawater CO₂. *ICES Journal of Marine Science: Journal Du Conseil*, 73(3), 659–669.
- 669 https://doi.org/10.1093/icesjms/fsv162
- Brown, D., & Edmunds, P. J. (2016). Differences in the responses of three scleractinians and the
 hydrocoral Millepora platyphylla to ocean acidification. *Marine Biology*, *163*(3).
 https://doi.org/10.1007/s00227-016-2837-7
- Byrne, M. (2011). Impact of ocean warming and ocean acidification on marine invertebrate life
 history stages: Vulnerabilities and potential for persistence in a changing ocean. *Oceanography Marine Science*, 49, 1–42.
- Camp, E. F., Nitschke, M. R., Rodolfo-Metalpa, R., Houlbreque, F., Gardner, S. G., Smith, D. J.,
 Zampighi, M., & Suggett, D. J. (2017). Reef-building corals thrive within hot-acidified and
 deoxygenated waters. *Scientific Reports*, 7(1). https://doi.org/10.1038/s41598-017-02383-y

- Caroselli, E., Gizzi, F., Prada, F., Marchini, C., Airi, V., Kaandorp, J., Falini, G., Dubinsky, Z., & Goffredo,
 S. (2019). Low and variable pH decreases recruitment efficiency in populations of a
 temperate coral naturally present at a CO₂ vent. *Limnology and Oceanography*, 64(3), 1059–
 1069. https://doi.org/10.1002/lno.11097
- Castillo, K. D., Ries, J. B., Bruno, J. F., & Westfield, I. T. (2014). The reef-building coral Siderastrea
 siderea exhibits parabolic responses to ocean acidification and warming. *Proceedings of the*
- 685 *Royal Society B: Biological Sciences, 281*(1797), 1–9. https://doi.org/10.1098/rspb.2014.1856
- Chan, N. C. S., & Connolly, S. R. (2013). Sensitivity of coral calcification to ocean acidification: A metaanalysis. *Global Change Biology*, *19*(1), 282–290. https://doi.org/10.1111/gcb.12011
- Chauvin, A., Denis, V., & Cuet, P. (2011). Is the response of coral calcification to seawater acidification
 related to nutrient loading? *Coral Reefs*, *30*(4), 911–923. https://doi.org/10.1007/s00338011-0786-7
- Chua, C., Leggat, W., Moya, A., & Baird, A. (2013). Near-future reductions in pH will have no
 consistent ecological effects on the early life-history stages of reef corals. *Marine Ecology Progress Series*, 486, 143–151. https://doi.org/10.3354/meps10318
- Cohen, A. L., & McConnaughey, T. A. (2003). Geochemical Perspectives on Coral Mineralization.
 Reviews in mineralogy and geochemistry, 54(1), 151-187.
- Comeau, S., Carpenter, R. C., & Edmunds, P. J. (2013). Effects of feeding and light intensity on the
 response of the coral Porites rus to ocean acidification. *Marine Biology*, *160*(5), 1127–1134.
 https://doi.org/10.1007/s00227-012-2165-5
- Comeau, S., Carpenter, R. C., & Edmunds, P. J. (2014). Effects of irradiance on the response of the
 coral Acropora pulchra and the calcifying alga Hydrolithon reinboldii to temperature
 elevation and ocean acidification. *Journal of Experimental Marine Biology and Ecology*, 453,
- 702 28–35. https://doi.org/10.1016/j.jembe.2013.12.013

- Comeau, S., Edmunds, P. J., Lantz, C. A., & Carpenter, R. C. (2015). Water flow modulates the
 response of coral reef communities to ocean acidification. *Scientific Reports*, *4*(1).
 https://doi.org/10.1038/srep06681
- Comeau, S., Edmunds, P. J., Spindel, N. B., & Carpenter, R. C. (2013). The responses of eight coral reef
 calcifiers to increasing partial pressure of CO₂ do not exhibit a tipping point. *Limnology and Oceanography*, 58(1), 388–398. https://doi.org/10.4319/lo.2013.58.1.0388
- Comeau, S., Edmunds, P. J., Spindel, N. B., & Carpenter, R. C. (2014). Fast coral reef calcifiers are
 more sensitive to ocean acidification in short-term laboratory incubations. *Limnology and Oceanography*, *59*(3), 1081–1091. https://doi.org/10.4319/lo.2014.59.3.1081
- Crawley, A., Kline, D. I., Dunn, S., Anthony, K., & Dove, S. (2010). The effect of ocean acidification on
 symbiont photorespiration and productivity in Acropora formosa. *Global Change Biology*, *16*(2), 851–863. https://doi.org/10.1111/j.1365-2486.2009.01943.x
- Crook, E. D., Cohen, A. L., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A. (2013). Reduced
 calcification and lack of acclimatization by coral colonies growing in areas of persistent
 natural acidification. *Proceedings of the National Academy of Sciences, 110*(27), 11044–
- 718 11049. https://doi.org/10.1073/pnas.1301589110
- Crook, E. D., Potts, D., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A. (2012). Calcifying coral
 abundance near low-pH springs: Implications for future ocean acidification. *Coral Reefs*,

721 *31*(1), 239–245. https://doi.org/10.1007/s00338-011-0839-y

- Cumbo, V. R., Fan, T. Y., & Edmunds, P. J. (2013). Effects of exposure duration on the response of
 Pocillopora damicornis larvae to elevated temperature and high pCO2. *Journal of Experimental Marine Biology and Ecology,* 439, 100–107.
 https://doi.org/10.1016/j.jembe.2012.10.019
- Cyronak, T., Takeshita, Y., Courtney, T. A., DeCarlo, E. H., Eyre, B. D., Kline, D. I., Martz, T., Page, H.,
 Price, N. N., Smith, J., Stoltenberg, L., Tresguerres, M., & Andersson, A. J. (2020). Diel

- temperature and pH variability scale with depth across diverse coral reef habitats. *Limnology and Oceanography Letters*, 5(2), 193–203. https://doi.org/10.1002/lol2.10129
- 730 Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P.
- 731 R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings*
- 732 of the National Academy of Sciences, 105(18), 6668–6672.
 733 https://doi.org/10.1073/pnas.0709472105
- Dickson, A. G. (1993). The measurement of seawater pH. *Marine Chemistry*, 44(2–4), 131–142.
 https://doi.org/10.1016/0304-4203(93)90198-W
- Dickson, A. G., Sabine, C. L., Christian, J. R., Bargeron, C. P., & North Pacific Marine Science
 Organization (Eds.). (2007). *Guide to best practices for ocean CO2 measurements*. North
 Pacific Marine Science Organization.
- Dimond, J. L., Holzman, B. J., & Bingham, B. L. (2012). Thicker host tissues moderate light stress in a
 cnidarian endosymbiont. *Journal of Experimental Biology*, *215*(13), 2247–2254.
 https://doi.org/10.1242/jeb.067991
- 742Dixon, G. B., Davies, S. W., Aglyamova, G. V., Meyer, E., Bay, L. K., & Matz, M. V. (2015). Genomic743determinants of coral heat tolerance across latitudes. *Science*, *348*(6242), 1460–1462.
- 744 https://doi.org/10.1126/science.1261224
- Donelson, J. M., Munday, P. L., McCormick, M. I., & Pitcher, C. R. (2012). Rapid transgenerational
 acclimation of a tropical reef fish to climate change. *Nature Climate Change*, 2(1), 30–32.
 https://doi.org/10.1038/nclimate1323
- Doney, S. C., Fabry, V. J., Feely, R. A., & Kleypas, J. A. (2009). Ocean Acidification: The Other CO₂
 Problem. *Annual Review of Marine Science*, 1(1), 169–192.
 https://doi.org/10.1146/annurev.marine.010908.163834
- Doo, S. S., Edmunds, P. J., & Carpenter, R. C. (2019). Ocean acidification effects on in situ coral reef
 metabolism. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598-019-48407-7

- Doropoulos, C., Ward, S., Diaz-Pulido, G., Hoegh-Guldberg, O., & Mumby, P. J. (2012). Ocean
 acidification reduces coral recruitment by disrupting intimate larval-algal settlement
 interactions: Elevated CO2 alters CCA-larval interactions. *Ecology Letters*, *15*(4), 338–346.
 https://doi.org/10.1111/j.1461-0248.2012.01743.x
- Dufault, A. M., Cumbo, V. R., Fan, T.-Y., & Edmunds, P. J. (2012). Effects of diurnally oscillating pCO 2
 on the calcification and survival of coral recruits. *Proceedings of the Royal Society B: Biological Sciences*, 279(1740), 2951–2958. https://doi.org/10.1098/rspb.2011.2545
- Dufault, A. M., Ninokawa, A., Bramanti, L., Cumbo, V. R., Fan, T.-Y., & Edmunds, P. J. (2013). The role
 of light in mediating the effects of ocean acidification on coral calcification. *Journal of Experimental Biology*, *216*(9), 1570–1577. https://doi.org/10.1242/jeb.080549
- Dupont, S., Dorey, N., Stumpp, M., Melzner, F., & Thorndyke, M. (2013). Long-term and trans-lifecycle effects of exposure to ocean acidification in the green sea urchin Strongylocentrotus
 droebachiensis. *Marine Biology*, *160*(8), 1835–1843. https://doi.org/10.1007/s00227-0121921-x
- Edmunds, P. J. (2011). Zooplanktivory ameliorates the effects of ocean acidification on the reef coral
 Porites spp. *Limnology and Oceanography*, 56(6), 2402–2410.
 https://doi.org/10.4319/lo.2011.56.6.2402
- Edmunds, P. J., Brown, D., & Moriarty, V. (2012). Interactive effects of ocean acidification and
 temperature on two scleractinian corals from Moorea, French Polynesia. *Global Change Biology*, *18*(7), 2173–2183. https://doi.org/10.1111/j.1365-2486.2012.02695.x
- Edmunds, P. J., & Burgess, S. C. (2016). Size-dependent physiological responses of the branching coral
 Pocillopora verrucosa to elevated temperature and *p*CO2. *Journal of Experimental Biology*.
 https://doi.org/10.1242/jeb.146381
- Edmunds, P. J., Doo, S. S., & Carpenter, R. C. (2019). Changes in coral reef community structure in
 response to year-long incubations under contrasting pCO2 regimes. *Marine Biology*, *166*(7).
 https://doi.org/10.1007/s00227-019-3540-2

- Edmunds, P. J., Doo, S. S., & Carpenter, R. C. (2020). Year-long effects of high pCO2 on the community
 structure of a tropical fore reef assembled in outdoor flumes. *ICES Journal of Marine Science*,
 77(3), 1055–1065. https://doi.org/10.1093/icesjms/fsaa015
- Edmunds, P. J., & Wall, C. B. (2014). Evidence that high *p*CO₂ affects protein metabolism in tropical
 reef corals. *The Biological Bulletin*, 227(1), 68–77. https://doi.org/10.1086/BBLv227n1p68
- Erez, J., Reynaud, S., Silverman, J., Schneider, K., & Allemand, D. (2011). Coral calcification under
 ocean acidification and global change. In Z. Dubinsky & N. Stambler (Eds.), *Coral Reefs: An Ecosystem in Transition* (pp. 151–176). Springer Netherlands. https://doi.org/10.1007/97894-007-0114-4_10
- Evensen, N., Edmunds, P., & Sakai, K. (2015). Effects of pCO2 on spatial competition between the
 corals Montipora aequituberculata and Porites lutea. *Marine Ecology Progress Series, 541*,
 123–134. https://doi.org/10.3354/meps11512
- Evensen, N. R., & Edmunds, P. J. (2016). Interactive effects of ocean acidification and neighbouring
 corals on the growth of Pocillopora verrucosa. *Marine Biology*, *163*(7).
 https://doi.org/10.1007/s00227-016-2921-z
- Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., Okazaki, R.,
 Muehllehner, N., Glas, M. S., & Lough, J. M. (2011). Losers and winners in coral reefs
 acclimatized to elevated carbon dioxide concentrations. *Nature Climate Change*, 1(3), 165–
 169. https://doi.org/10.1038/nclimate1122
- Fantazzini, P., Mengoli, S., Pasquini, L., Bortolotti, V., Brizi, L., Mariani, M., Di Giosia, M., Fermani, S.,
 Capaccioni, B., Caroselli, E., Prada, F., Zaccanti, F., Levy, O., Dubinsky, Z., Kaandorp, J. A.,
 Konglerd, P., Hammel, J. U., Dauphin, Y., Cuif, J.-P., ... Goffredo, S. (2015). Gains and losses of
 coral skeletal porosity changes with ocean acidification acclimation. *Nature Communications*,
- 802 *6*(1). https://doi.org/10.1038/ncomms8785

- Feder, M. E., & Walser, J.-C. (2005). The biological limitations of transcriptomics in elucidating stress
 and stress responses. *Journal of Evolutionary Biology*, 18(4), 901–910.
 https://doi.org/10.1111/j.1420-9101.2005.00921.x
- Feely, R. A. (2004). Impact of Anthropogenic CO2 on the CaCO3 System in the Oceans. *Science*,
 305(5682), 362–366. https://doi.org/10.1126/science.1097329
- Feil, R., & Fraga, M. F. (2012). Epigenetics and the environment: Emerging patterns and implications.
 Nature Reviews Genetics, *13*(2), 97–109. https://doi.org/10.1038/nrg3142
- 810 Gattuso, J.-P., & Hansson, L. (Eds.). (2011). *Ocean acidification*. Oxford University Press, Oxford.
- Gattuso, J. P., Epitalon, J. M., Lavigne, H., Orr, J., Gentili, B., Hofmann, A., ... & Rae, J. (2020). seacarb:
 Seawater Carbonate Chemistry, version 3.2. 14.
- 813 Georgiou, L., Falter, J., Trotter, J., Kline, D. I., Holcomb, M., Dove, S. G., Hoegh-Guldberg, O., &
 814 McCulloch, M. (2015). Ph homeostasis during coral calcification in a free ocean CO 2

815 enrichment (FOCE) experiment, Heron Island reef flat, Great Barrier Reef. Proceedings of the

- 816
 National
 Academy
 of
 Sciences,
 112(43),
 13219–13224.

 817
 https://doi.org/10.1073/pnas.1505586112
 https://doi.org/10.1073/pnas.1505586112
 https://doi.org/10.1073/pnas.1505586112
- Graham, E. R., & Sanders, R. W. (2016). Species-specific photosynthetic responses of symbiotic
 zoanthids to thermal stress and ocean acidification. *Marine Ecology*, *37*(2), 442–458.
 https://doi.org/10.1111/maec.12291
- Greenbaum, D., Colangelo, C., Williams, K., & Gerstein, M. (2003). Comparing protein abundance and
 mRNA expression levels on a genomic scale. *Genome Biology*, 4(9), 117.
 https://doi.org/10.1186/gb-2003-4-9-117
- Grottoli, A. G., Rodrigues, L. J., & Palardy, J. E. (2006). Heterotrophic plasticity and resilience in
 bleached corals. *Nature*, 440(7088), 1186–1189. https://doi.org/10.1038/nature04565
- Guest, J. R., Baird, A. H., Maynard, J. A., Muttaqin, E., Edwards, A. J., Campbell, S. J., Yewdall, K.,
 Affendi, Y. A., & Chou, L. M. (2012). Contrasting patterns of coral bleaching susceptibility in

- 828 2010 suggest an adaptive response to thermal stress. *PLoS ONE*, 7(3), e33353.
 829 https://doi.org/10.1371/journal.pone.0033353
- Hendriks, I. E., Duarte, C. M., & Álvarez, M. (2010). Vulnerability of marine biodiversity to ocean
 acidification: A meta-analysis. *Estuarine, Coastal and Shelf Science, 86*(2), 157–164.
 https://doi.org/10.1016/j.ecss.2009.11.022
- Hettinger, A., Sanford, E., Hill, T. M., Russell, A. D., Sato, K. N. S., Hoey, J., Forsch, M., Page, H. N., &
 Gaylord, B. (2012). Persistent carry-over effects of planktonic exposure to ocean acidification
 in the Olympia oyster. *Ecology*, *93*(12), 2758–2768. https://doi.org/10.1890/12-0567.1
- Hoadley, K. D., Pettay, D. T., Grottoli, A. G., Cai, W.-J., Melman, T. F., Schoepf, V., Hu, X., Li, Q., Xu, H.,
 Wang, Y., Matsui, Y., Baumann, J. H., & Warner, M. E. (2015). Physiological response to
 elevated temperature and pCO2 varies across four Pacific coral species: Understanding the
 unique host+symbiont response. *Scientific Reports*, *5*(1). https://doi.org/10.1038/srep18371
- 840 Holcomb, M., Venn, A. A., Tambutté, E., Tambutté, S., Allemand, D., Trotter, J., & McCulloch, M.
- 841 (2015). Coral calcifying fluid pH dictates response to ocean acidification. *Scientific Reports*,

842 4(1). https://doi.org/10.1038/srep05207

- Horvath, K. M., Castillo, K. D., Armstrong, P., Westfield, I. T., Courtney, T., & Ries, J. B. (2016). Next century ocean acidification and warming both reduce calcification rate, but only acidification
 alters skeletal morphology of reef-building coral Siderastrea siderea. *Scientific Reports, 6*(1).
- 846 https://doi.org/10.1038/srep29613
- Horwitz, R., Hoogenboom, M. O., & Fine, M. (2017). Spatial competition dynamics between reef
 corals under ocean acidification. *Scientific Reports*, 7(1). https://doi.org/10.1038/srep40288
- 849 Houlbrèque, F., Reynaud, S., Godinot, C., Oberhänsli, F., Rodolfo-Metalpa, R., & Ferrier-Pagès, C.
- 850 (2015). Ocean acidification reduces feeding rates in the scleractinian coral Stylophora
- pistillata: Acidification and Stylophora nutrition. *Limnology and Oceanography, 60*(1), 89–99.
- 852 https://doi.org/10.1002/lno.10003

IPCC 2021. Fox-Kemper, B., H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S. S. Drijfhout, T. L. Edwards, N. R.
Golledge, M. Hemer, R. E. Kopp, G. Krinner, A. Mix, D. Notz, S. Nowicki, I. S. Nurhati, L. Ruiz, JB. Sallée, A. B. A. Slangen, Y. Yu, 2021, Ocean, Cryosphere and Sea Level Change. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V.,
P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis,
M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu

- and B. Zhou (eds.)]. Cambridge University Press. In Press.
- Jacob, H., Besson, M., Swarzenski, P. W., Lecchini, D., & Metian, M. (2020). Effects of Virgin Micro and Nanoplastics on Fish: Trends, Meta-Analysis, and Perspectives. *Environmental Science & Technology*, *54*(8), 4733–4745. https://doi.org/10.1021/acs.est.9b05995
- Jagers, S. C., Matti, S., Crépin, A.-S., Langlet, D., Havenhand, J. N., Troell, M., Filipsson, H. L., Galaz, V.
 R., & Anderson, L. G. (2019). Societal causes of, and responses to, ocean acidification. *Ambio*,
 48(8), 816–830. https://doi.org/10.1007/s13280-018-1103-2
- 867 Jiang, L., Guo, M.-L., Zhang, F., Zhang, Y.-Y., Zhou, G.-W., Lei, X.-M., Yuan, X.-C., Sun, Y.-F., Yuan, T., 868 Cai, L., Lian, J.-S., Liu, S., Qian, P.-Y., & Huang, H. (2020). Impacts of elevated temperature 869 and pCO2 on the brooded larvae of Pocillopora damicornis from Luhuitou Reef, China: 870 Evidence 331-344. for local acclimatization. Coral Reefs, 39(2), https://doi.org/10.1007/s00338-020-01894-x 871
- Jiang, L.-Q., Carter, B. R., Feely, R. A., Lauvset, S. K., & Olsen, A. (2019). Surface ocean pH and buffer
 capacity: Past, present and future. *Scientific Reports*, *9*(1). https://doi.org/10.1038/s41598019-55039-4
- Jokiel, P. L. (2011). The reef coral two compartment proton flux model: A new approach relating
 tissue-level physiological processes to gross corallum morphology. *Journal of Experimental Marine Biology and Ecology*, 409(1–2), 1–12. https://doi.org/10.1016/j.jembe.2011.10.008

- Jury, C. P., Delano, M. N., & Toonen, R. J. (2019). High heritability of coral calcification rates and
 evolutionary potential under ocean acidification. *Scientific Reports*, *9*(1).
 https://doi.org/10.1038/s41598-019-56313-1
- Kaniewska, P., Campbell, P. R., Kline, D. I., Rodriguez-Lanetty, M., Miller, D. J., Dove, S., & HoeghGuldberg, O. (2012). Major cellular and physiological impacts of ocean acidification on a reef
 building coral. *PLoS ONE*, 7(4), e34659. https://doi.org/10.1371/journal.pone.0034659
- Kavousi, J., Parkinson, J. E., & Nakamura, T. (2016). Combined ocean acidification and low
 temperature stressors cause coral mortality. *Coral Reefs*, *35*(3), 903–907.
 https://doi.org/10.1007/s00338-016-1459-3
- Kawahata, H., Fujita, K., Iguchi, A., Inoue, M., Iwasaki, S., Kuroyanagi, A., Maeda, A., Manaka, T.,
 Moriya, K., Takagi, H., Toyofuku, T., Yoshimura, T., & Suzuki, A. (2019). Perspective on the

889

and foraminifera in a high CO2 world "hot house". *Progress in Earth and Planetary Science*,
6(1). https://doi.org/10.1186/s40645-018-0239-9

response of marine calcifiers to global warming and ocean acidification—Behavior of corals

- Kline, D. I., Teneva, L., Okamoto, D. K., Schneider, K., Caldeira, K., Miard, T., Chai, A., Marker, M.,
 Dunbar, R. B., Mitchell, B. G., Dove, S., & Hoegh-Guldberg, O. (2019). Living coral tissue slows
 skeletal dissolution related to ocean acidification. *Nature Ecology & Evolution*, *3*(10), 1438–
 1444. https://doi.org/10.1038/s41559-019-0988-x
- Koussounadis, A., Langdon, S. P., Um, I. H., Harrison, D. J., & Smith, V. A. (2015). Relationship
 between differentially expressed mRNA and mRNA-protein correlations in a xenograft model
 system. *Scientific Reports*, *5*(1). https://doi.org/10.1038/srep10775
- 899 Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., & Gattuso,
- J. (2013). Impacts of ocean acidification on marine organisms: Quantifying sensitivities and
 interaction with warming. *Global Change Biology*, *19*(6), 1884–1896.
 https://doi.org/10.1111/gcb.12179

- Wroeker, K. J., Kordas, R. L., Crim, R. N., & Singh, G. G. (2010). Meta-analysis reveals negative yet
 variable effects of ocean acidification on marine organisms: Biological responses to ocean
 acidification. *Ecology Letters*, *13*(11), 1419–1434. https://doi.org/10.1111/j.14610248.2010.01518.x
- Kurihara, H., Wouters, J., & Yasuda, N. (2019). Seasonal calcification of the coral Acropora digitifera
 from a subtropical marginal Okinawa reef under ocean acidification. *Coral Reefs*, *38*(3), 443–
 454. https://doi.org/10.1007/s00338-019-01794-9
- Langdon, C. (2005). Effect of elevated pCO₂ on photosynthesis and calcification of corals and
 interactions with seasonal change in temperature/irradiance and nutrient enrichment.
 Journal of Geophysical Research, 110(C9). https://doi.org/10.1029/2004JC002576
- Leong, W., & Pawlik, J. (2010). Evidence of a resource trade-off between growth and chemical
 defenses among Caribbean coral reef sponges. *Marine Ecology Progress Series, 406,* 71–78.
 https://doi.org/10.3354/meps08541
- Liew, Y. J., Zoccola, D., Li, Y., Tambutté, E., Venn, A. A., Michell, C. T., Cui, G., Deutekom, E. S.,
 Kaandorp, J. A., Voolstra, C. R., Forêt, S., Allemand, D., Tambutté, S., & Aranda, M. (2018).
 Epigenome-associated phenotypic acclimatization to ocean acidification in a reef-building
- 919 coral. *Science Advances*, 10.
- Loya, Y., Sakai, K., Yamazato, K., Nakano, Y., Sambali, H., & van Woesik, R. (2001). Coral bleaching:
 The winners and the losers. *Ecology Letters*, 4(2), 122–131. https://doi.org/10.1046/j.1461 0248.2001.00203.x
- Marcelino, V. R., Morrow, K. M., Oppen, M. J. H., Bourne, D. G., & Verbruggen, H. (2017). Diversity
 and stability of coral endolithic microbial communities at a naturally high pCO₂ reef.
 Molecular Ecology, 26(19), 5344–5357. https://doi.org/10.1111/mec.14268
- Martinez, A., Crook, E. D., Barshis, D. J., Potts, D. C., Rebolledo-Vieyra, M., Hernandez, L., & Paytan, A.
 (2019). Species-specific calcification response of Caribbean corals after 2-year

transplantation to a low aragonite saturation submarine spring. *Proceedings of the Royal Society B: Biological Sciences, 286*(1905), 20190572. https://doi.org/10.1098/rspb.2019.0572
Maynard, J. A., Anthony, K. R. N., Marshall, P. A., & Masiri, I. (2008). Major bleaching events can lead
to increased thermal tolerance in corals. *Marine Biology, 155*(2), 173–182.
https://doi.org/10.1007/s00227-008-1015-y

- McCulloch, M., Falter, J., Trotter, J., & Montagna, P. (2012). Coral resilience to ocean acidification and
 global warming through pH up-regulation. *Nature Climate Change*, 2(8), 623–627.
 https://doi.org/10.1038/nclimate1473
- 936 McLachlan, R. H., Price, J. T., Solomon, S. L., & Grottoli, A. G. (2020). Thirty years of coral heat-stress
- 937 experiments: A review of methods. *Coral Reefs*, *39*(4), 885–902.
 938 https://doi.org/10.1007/s00338-020-01931-9
- Meron, D., Atias, E., Iasur Kruh, L., Elifantz, H., Minz, D., Fine, M., & Banin, E. (2011). The impact of
 reduced pH on the microbial community of the coral Acropora eurystoma. *The ISME Journal*,
 5(1), 51–60. https://doi.org/10.1038/ismej.2010.102
- 942 Metcalfe, N. B., & Monaghan, P. (2001). Compensation for a bad start: Grow now, pay later? *Trends* 943 *in Ecology & Evolution*, *16*(5), 254–260. https://doi.org/10.1016/S0169-5347(01)02124-3
- Middlebrook, R., Hoegh-Guldberg, O., & Leggat, W. (2008). The effect of thermal history on the
 susceptibility of reef-building corals to thermal stress. *Journal of Experimental Biology*, *211*(7), 1050–1056. https://doi.org/10.1242/jeb.013284
- Miller, G. M., Watson, S.-A., Donelson, J. M., McCormick, M. I., & Munday, P. L. (2012). Parental
 environment mediates impacts of increased carbon dioxide on a coral reef fish. *Nature Climate Change*, 2(12), 858–861. https://doi.org/10.1038/nclimate1599
- Morais, J., & Santos, B. A. (2018). Limited potential of deep reefs to serve as refuges for tropical
 Southwestern Atlantic corals. *Ecosphere*, *9*(7), e02281. https://doi.org/10.1002/ecs2.2281

- Morita, M., Suwa, R., Iguchi, A., Nakamura, M., Shimada, K., Sakai, K., & Suzuki, A. (2010). Ocean
 acidification reduces sperm flagellar motility in broadcast spawning reef invertebrates.
 Zygote, 18(2), 103–107. https://doi.org/10.1017/S0967199409990177
- Morrow, K. M., Bourne, D. G., Humphrey, C., Botté, E. S., Laffy, P., Zaneveld, J., Uthicke, S., Fabricius,
 K. E., & Webster, N. S. (2015). Natural volcanic CO2 seeps reveal future trajectories for host–
 microbial associations in corals and sponges. *The ISME Journal, 9*(4), 894–908.
 https://doi.org/10.1038/ismej.2014.188
- Moya, A., Huisman, L., Forêt, S., Gattuso, J.-P., Hayward, D. C., Ball, E. E., & Miller, D. J. (2015). Rapid
 acclimation of juvenile corals to CO₂-mediated acidification by upregulation of heat shock
 protein and Bcl-2 genes. *Molecular Ecology*, 24(2), 438–452.
 https://doi.org/10.1111/mec.13021
- 963 Munday, P. L. (2014). Transgenerational acclimation of fishes to climate change and ocean
 964 acidification. *F1000Prime Reports, 6*. https://doi.org/10.12703/P6-99
- Muscatine, L., R. McCloskey, L., & E. Marian, R. (1981). Estimating the daily contribution of carbon
 from zooxanthellae to coral animal respiration: Muscatine et al. *Limnology and Oceanography*, 26(4), 601–611. https://doi.org/10.4319/lo.1981.26.4.0601
- Noonan, S. H. C., & Fabricius, K. E. (2016). Ocean acidification affects productivity but not the severity
 of thermal bleaching in some tropical corals. *ICES Journal of Marine Science: Journal Du Conseil, 73*(3), 715–726. https://doi.org/10.1093/icesjms/fsv127
- Noonan, S. H. C., Kluibenschedl, A., & Fabricius, K. E. (2018). Ocean acidification alters early
 successional coral reef communities and their rates of community metabolism. *PLOS ONE*, *13*(5), e0197130. https://doi.org/10.1371/journal.pone.0197130
- O'Brien, P. A., Smith, H. A., Fallon, S., Fabricius, K., Willis, B. L., Morrow, K. M., & Bourne, D. G. (2018). 974 975 Elevated CO2 Has Little Influence on the Bacterial Communities Associated With the pH-Porites 976 Tolerant Microbiology, 9. Coral, Massive spp. Frontiers in https://doi.org/10.3389/fmicb.2018.02621 977

- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. M., Gnanadesikan, A., Gruber, N.,
 Ishida, A., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A.,
 Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., ... Yool, A. (2005). Anthropogenic
 decline in high-latitude ocean carbonate by 2100. Nature, *437*(7059), 681-686.
- Pandolfi, J. M. (2003). Global trajectories of the long-term decline of coral reef ecosystems. *Science*,
 301(5635), 955–958. https://doi.org/10.1126/science.1085706
- Parker, L. M., Ross, P. M., O'Connor, W. A., Borysko, L., Raftos, D. A., & Pörtner, H.-O. (2012). Adult
 exposure influences offspring response to ocean acidification in oysters. *Global Change Biology*, *18*(1), 82–92. https://doi.org/10.1111/j.1365-2486.2011.02520.x
- 987 Pierrot, Lewis, D. E., & Wallace, D. W. R. (2006). MS Excel Program Developed for CO2 System Calculations. ORNL/CDIAC-105a. Carbon Dioxide Information Analysis Center, Oak Ridge 988 989 National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. https://doi.org/10.3334/CDIAC/otg.CO2SYS_XLS_CDIAC105a 990
- Pitts, K. A., Campbell, J. E., Figueiredo, J., & Fogarty, N. D. (2020). Ocean acidification partially
 mitigates the negative effects of warming on the recruitment of the coral, Orbicella
 faveolata. *Coral Reefs*, *39*(2), 281–292. https://doi.org/10.1007/s00338-019-01888-4
- Prada, F., Brizi, L., Franzellitti, S., Mengoli, S., Fermani, S., Polishchuk, I., Baraldi, N., Ricci, F., Palazzo,
 Q., Caroselli, E., Pokroy, B., Giorgini, L., Dubinsky, Z., Fantazzini, P., Falini, G., Goffredo, S., &
- Fabricius, K. E. (2021). Coral micro- and macro-morphological skeletal properties in response
 to life-long acclimatization at CO2 vents in Papua New Guinea. *Scientific Reports*, *11*(1).
 https://doi.org/10.1038/s41598-021-98976-9
- 999 Putnam, H. M. (2021). Avenues of reef-building coral acclimatization in response to rapid
 1000 environmental change. *Journal of Experimental Biology*, 224, jeb.239319.
 1001 https://doi.org/10.1242/jeb.239319

- Putnam, H. M., Davidson, J. M., & Gates, R. D. (2016). Ocean acidification influences host DNA
 methylation and phenotypic plasticity in environmentally susceptible corals. *Evolutionary Applications*, 9(9), 1165–1178. https://doi.org/10.1111/eva.12408
- Putnam, H. M., & Gates, R. D. (2015). Preconditioning in the reef-building coral Pocillopora
 damicornis and the potential for trans-generational acclimatization in coral larvae under
 future climate change conditions. *Journal of Experimental Biology*, *218*(15), 2365–2372.
 https://doi.org/10.1242/jeb.123018
- Putnam, H. M., Mayfield, A. B., Fan, T. Y., Chen, C. S., & Gates, R. D. (2013). The physiological and
 molecular responses of larvae from the reef-building coral Pocillopora damicornis exposed to
 near-future increases in temperature and pCO2. *Marine Biology*, *160*(8), 2157–2173.
 https://doi.org/10.1007/s00227-012-2129-9
- 1013 Raven. (2005). Ocean acidification due to increasing atmospheric carbon dioxide. Policy Document
 1014 12/05 (The Royal Society, London).
- 1015 Reynaud, S., Leclercq, N., Romaine-Lioud, S., Ferrier-Pages, C., Jaubert, J., & Gattuso, J.-P. (2003).
 1016 Interacting effects of CO2 partial pressure and temperature on photosynthesis and
 1017 calcification in a scleractinian coral. *Global Change Biology*, *9*(11), 1660–1668.
 1018 https://doi.org/10.1046/j.1365-2486.2003.00678.x
- 1019 Riebesell, U. (2008). Acid test for marine biodiversity. *Nature*, 454(7200), 46–47.
 1020 https://doi.org/10.1038/454046a
- Riebesell, U., Fabry, V. J., Hansson, L., & Gattuso, J. P. (2011). *Guide to best practices for ocean acidification research and data reporting*. Office for Official Publications of the European
 Communities.
- 1024 Ries, J. B. (2011). A physicochemical framework for interpreting the biological calcification response
 1025 to CO2-induced ocean acidification. *Geochimica et Cosmochimica Acta*, 75(14), 4053–4064.
 1026 https://doi.org/10.1016/j.gca.2011.04.025

1027 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F. S., Lambin, E. F., Lenton, T. M., Scheffer, 1028 M., Folke, C., Schellnhuber, H. J., Nykvist, B., de Wit, C. A., Hughes, T., van der Leeuw, S., 1029 Rodhe, H., Sörlin, S., Snyder, P. K., Costanza, R., Svedin, U., ... Foley, J. A. (2009). A safe 1030 operating for humanity. Nature, 461(7263), 472-475. space 1031 https://doi.org/10.1038/461472a

- Sayers, A. (2007). Tips and tricks in performing a systematic review. *British Journal of General Practice*, *57*(538), 425-425.
- 1034 Schoepf, V., Grottoli, A. G., Warner, M. E., Cai, W. J., Melman, T. F., Hoadley, K. D., Pettay, D. T., Hu,

1035 X., Li, Q., Xu, H., Wang, Y., Matsui, Y., & Baumann, J. H. (2013). Coral energy reserves and

calcification in a high-CO2 world at two temperatures. In S. C. A. Ferse (Ed.), PLoS ONE (Vol.

1037 8, Issue 10, p. e75049). https://doi.org/10.1371/journal.pone.0075049

1036

- Sekizawa, A., Uechi, H., Iguchi, A., Nakamura, T., Kumagai, N. H., Suzuki, A., Sakai, K., & Nojiri, Y.
 (2017). Intraspecific variations in responses to ocean acidification in two branching coral
 species. *Marine Pollution Bulletin*, 122(1–2), 282–287.
 https://doi.org/10.1016/j.marpolbul.2017.06.061
- Smith, J. N., Strahl, J., Noonan, S. H. C., Schmidt, G. M., Richter, C., & Fabricius, K. E. (2016). Reduced
 heterotrophy in the stony coral Galaxea fascicularis after life-long exposure to elevated
 carbon dioxide. *Scientific Reports*, 6(1). https://doi.org/10.1038/srep27019

Soriano-Santiago, O. S., Liñán-Cabello, M. A., Delgadillo-Nuño, M. A., Ortega-Ortiz, C., & CuevasVenegas, S. (2013). Physiological responses to oxidative stress associated with pH variations
in host tissue and zooxanthellae of hermatypic coral Pocillopora capitata. *Marine and Freshwater Behaviour and Physiology, 46*(5), 275–286.
https://doi.org/10.1080/10236244.2013.827877

Srednick, G., Bergman, J. L., Doo, S. S., Hawthorn, M., Ferree, J., Rojas, R., Arias, R., Edmunds, P. J., &
 Carpenter, R. C. (2020). Shallow coral reef free ocean carbon enrichment: Novel in situ

1052 flumes to manipulate pCO₂ on shallow tropical coral reef communities. *Limnology and* 1053 *Oceanography: Methods, 18*(3), 116–128. https://doi.org/10.1002/lom3.10349

- Strahl, J., Stolz, I., Uthicke, S., Vogel, N., Noonan, S. H. C., & Fabricius, K. E. (2015). Physiological and
 ecological performance differs in four coral taxa at a volcanic carbon dioxide seep.
 Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 184,
- 1057 179–186. https://doi.org/10.1016/j.cbpa.2015.02.018
- Tambutté, E., Venn, A. A., Holcomb, M., Segonds, N., Techer, N., Zoccola, D., Allemand, D., &
 Tambutté, S. (2015). Morphological plasticity of the coral skeleton under CO2-driven
 seawater acidification. *Nature Communications*, 6(1). https://doi.org/10.1038/ncomms8368
- 1061 Thompson, D. M., & van Woesik, R. (2009). Corals escape bleaching in regions that recently and 1062 historically experienced frequent thermal stress. *Proceedings of the Royal Society B:* 1063 *Biological Sciences*, 276(1669), 2893–2901. https://doi.org/10.1098/rspb.2009.0591
- 1064Towle, E. K., Enochs, I. C., & Langdon, C. (2015). Threatened caribbean coral is able to mitigate the1065adverse effects of ocean acidification on calcification by increasing feeding rate. PLOS ONE,

1066 *10*(4), e0123394. https://doi.org/10.1371/journal.pone.0123394

- UNEP-WCMC, WorldFish Centre, WRI, TNC (2010). Global distribution of warm-water coral reefs,
 compiled from multiple sources including the Millennium Coral Reef Mapping Project.
 Version 4.0. Includes contributions from IMaRS-USF and IRD (2005), IMaRS-USF (2005) and
 Spalding et al. (2001). Cambridge (UK): UNEP World Conservation Monitoring Centre. URL:
- 1071 http://data.unep-wcmc.org/datasets/1
- Vargas, C. A., Lagos, N. A., Lardies, M. A., Duarte, C., Manríquez, P. H., Aguilera, V. M., Broitman, B.,
 Widdicombe, S., & Dupont, S. (2017). Species-specific responses to ocean acidification should
 account for local adaptation and adaptive plasticity. *Nature Ecology & Evolution*, 1(4).
 https://doi.org/10.1038/s41559-017-0084

- 1076 Venn, A. A., Bernardet, C., Chabenat, A., Tambutté, E., & Tambutté, S. (2020). Paracellular transport 1077 to the coral calcifying medium: Effects of environmental parameters. The Journal of 1078 Experimental Biology, 223(17), jeb227074. https://doi.org/10.1242/jeb.227074
- Venn, A. A., Tambutté, E., Caminiti-Segonds, N., Techer, N., Allemand, D., & Tambutté, S. (2019). 1079
- 1080 Effects of light and darkness on pH regulation in three coral species exposed to seawater 1081 acidification. Scientific Reports, 9(1). https://doi.org/10.1038/s41598-018-38168-0
- 1082 Venn, A. A., Tambutte, E., Holcomb, M., Laurent, J., Allemand, D., & Tambutte, S. (2013). Impact of 1083 seawater acidification on pH at the tissue-skeleton interface and calcification in reef corals. 1084 Proceedings the National Academy Sciences, 110(5), 1634-1639. of of https://doi.org/10.1073/pnas.1216153110
- 1086 Veron, J., Stafford-Smith, M., DeVantier, L., & Turak, E. (2015). Overview of distribution patterns of 1087 zooxanthellate Scleractinia. Frontiers Marine Science, 1. in 1088 https://doi.org/10.3389/fmars.2014.00081
- 1089 Viyakarn, V., Lalitpattarakit, W., Chinfak, N., Jandang, S., Kuanui, P., Khokiattiwong, S., & Chavanich, 1090 S. (2015). Effect of lower pH on settlement and development of coral, Pocillopora damicornis 1091 (Linnaeus, 1758). Ocean Science Journal, 50(2), 475-480. https://doi.org/10.1007/s12601-1092 015-0043-z
- 1093 Von Euw, S., Zhang, Q., Manichev, V., Murali, N., Gross, J., Feldman, L. C., Gustafsson, T., Flach, C., 1094 Mendelsohn, R., & Falkowski, P. G. (2017). Biological control of aragonite formation in stony 1095 corals. Science, 356(6341), 933–938. https://doi.org/10.1126/science.aam6371
- 1096 Wall, C. B., Mason, R. A. B., Ellis, W. R., Cunning, R., & Gates, R. D. (2017). Elevated pCO₂ affects tissue 1097 biomass composition, but not calcification, in a reef coral under two light regimes. Royal 1098 Society Open Science, 4(11), 170683. https://doi.org/10.1098/rsos.170683
- Wall, M., Fietzke, J., Crook, E. D., & Paytan, A. (2019). Using B isotopes and B/Ca in corals from low 1099 1100 saturation springs to constrain calcification mechanisms. Nature Communications, 10(1).
- 1101 https://doi.org/10.1038/s41467-019-11519-9

Webster, N. S., Negri, A. P., Botté, E. S., Laffy, P. W., Flores, F., Noonan, S., Schmidt, C., & Uthicke, S.
(2016). Host-associated coral reef microbes respond to the cumulative pressures of ocean

1104 warming and ocean acidification. *Scientific Reports*, *6*(1). https://doi.org/10.1038/srep19324


- 1105 Webster, N. S., Negri, A. P., Flores, F., Humphrey, C., Soo, R., Botté, E. S., Vogel, N., & Uthicke, S.
- (2013). Near-future ocean acidification causes differences in microbial associations within
 diverse coral reef taxa: pH effects on marine microbial associations. *Environmental Microbiology Reports*, 5(2), 243–251. https://doi.org/10.1111/1758-2229.12006
- Wijgerde, T., Silva, C. I. F., Scherders, V., van Bleijswijk, J., & Osinga, R. (2014). Coral calcification
 under daily oxygen saturation and pH dynamics reveals the important role of oxygen. *Biology*


1111 *Open, 3*(6), 489–493. https://doi.org/10.1242/bio.20147922

- Yang, S.-Y., Lu, C.-Y., Tang, S.-L., Das, R. R., Sakai, K., Yamashiro, H., & Yang, S.-H. (2020). Effects of
 ocean acidification on coral endolithic bacterial communities in Isopora palifera and Porites
 lobata. *Frontiers in Marine Science*, *7*. https://doi.org/10.3389/fmars.2020.603293
- 1115 Zande, R. M., Achlatis, M., Bender-Champ, D., Kubicek, A., Dove, S., & Hoegh-Guldberg, O. (2020).

1116 Paradise lost: End-of-century warming and acidification under business-as-usual emissions

- 1117 have severe consequences for symbiotic corals. *Global Change Biology*, *26*(4), 2203–2219.
- 1118 https://doi.org/10.1111/gcb.14998
- Zhou, G., Yuan, T., Cai, L., Zhang, W., Tian, R., Tong, H., Jiang, L., Yuan, X., Liu, S., Qian, P., & Huang, H.
 (2016). Changes in microbial communities, photosynthesis and calcification of the coral
 Acropora gemmifera in response to ocean acidification. *Scientific Reports*, 6(1).
 https://doi.org/10.1038/srep35971
- Zoccola, D., Innocenti, A., Bertucci, A., Tambutté, E., Supuran, C., & Tambutté, S. (2016). Coral
 carbonic anhydrases: Regulation by ocean acidification. *Marine Drugs*, *14*(6), 109.
 https://doi.org/10.3390/md14060109
- 1126
- 1127







			Duration			Genus			Life stage					
ALL			• - ⁷ / ₄₀₀	• ⁻ ⁻ ⁻	• - 7.3 noot	(S)41001, CT - •		• 00 1168		· Setto Wij	- 40 ¹¹¹ ^{Can}			
Fitness	Calcification Growth & development Reproductive investment & success Mortality		• • • •	• • •	• .	•		•	•		• • •			
Respiration & Photophysic	Dlogy Photophysiology Respiration Productivity	-	•	•	•	·		•			• • •		10	oints (%) 0
Biomolecules	Lipid Carbohydrate Protein Energy content	-		• •	•						•) ; per of
Cellular stress response mechanism	Other molecular actors Oxidative stress Apoptosis	-	•	•	•			•			•	e	• 2	oints 100 200 300
Community	Community structure	-											٠ د	400
Microbiome	Bacterial community composition]-[•	•						•		•	500
Epigenetic mechanism	DNA methylation level]-[·			•			•			
Metabolome	Metabolite profile	}			•			•			•			